



A Baseline Study on Groundwater Quality of the Tourist Island, Pulau Tiga, Sabah, Malaysia

Chin Yik Lin (Corresponding Author)

School of Science and Technology, Universiti Malaysia Sabah
Beg Berkunci 2073, 88999, Kota Kinabalu, Sabah, Malaysia
Tel: 60-88-325-808 E-mail: cy_lin_ars@hotmail.com

Mohd Harun Abdullah

School of Science and Technology, Universiti Malaysia Sabah
Beg Berkunci 2073, 88999, Kota Kinabalu, Sabah, Malaysia
Tel: 60-88-325-808 E-mail: harunabd@ums.edu.my

Ahmad Zaharin Aris

Faculty of Environmental Studies, Universiti Putra Malaysia
43400 UPM, Serdang, Selangor, Malaysia
Tel: 60-88-325-808 E-mail: zaharin@env.upm.edu.my

Sarva Mangala Praveena

School of Science and Technology, Universiti Malaysia Sabah
Beg Berkunci 2073, 88999, Kota Kinabalu, Sabah, Malaysia
Tel: 60-88-325-808 E-mail: smpraveena@gmail.com

This research was financially supported by the Ministry of Higher Education, Malaysia through the Fundamental Research Grant No. FRG0050-ST-1/2006. Permission from the Sabah Park Trustees for the study site is highly acknowledged. Besides, the authors would also opt to acknowledge Chua Li Ying, Lim Kuan Leang and Ooi Chun Wei for their assistance during the fieldworks.

Abstract

Pulau Tiga is a group of small islands located at Kimanis Bay off the western coast of Sabah. This preliminary study on groundwater of Pulau Tiga is undertaken to provide guidance and baseline data for future references. An understanding of the chemistry and behavior of water in the island aquifer is crucial for the determination of the availability of freshwater, and also to create a platform for considering proper management and successful remediation of groundwater resources on island to prevent over exploitation of such limited resources. Groundwater samples were collected from five wells on Pulau Tiga to understand the groundwater chemistry based on various ion composition (Ca^{2+} , Mg^{2+} , Na^+ , K^+ , HCO_3^- , SO_4^{2-} , Cl^- , NO_3^-), *in situ* parameters such as DO (Dissolved oxygen), EC (Electrical conductivity), TDS (Total dissolved solids), pH, salinity and temperature. Even though groundwater in Pulau Tiga was constantly subject to abstraction activities for its tourism purposes, preliminary results showed that the groundwater still remains freshwater type. The Piper diagram suggested that the groundwater facies generally range from Ca- HCO_3 to Ca-Cl waters type. The processes influencing the groundwater chemistry are mainly dissolution of minerals.

Keywords: Groundwater, Pulau Tiga, Hydrochemical, Hydrochemical Facies, Groundwater Chemistry