

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/297728790>

Complete genome sequence of a giant Vibrio phage ValKK3 infecting *Vibrio alginolyticus*

Article in *Genomics Data* - March 2016

DOI: 10.1016/j.gdata.2016.03.002

CITATIONS

8

READS

137

4 authors, including:



Mohammad Tamrin Mohamad Lal
Universiti Malaysia Sabah (UMS)

23 PUBLICATIONS 99 CITATIONS

[SEE PROFILE](#)



Kishio Hatai
Nippon Veterinary and Life Science University

266 PUBLICATIONS 3,436 CITATIONS

[SEE PROFILE](#)



Julian Ransangan
Universiti Malaysia Sabah (UMS)

118 PUBLICATIONS 422 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Probiotic properties of fish intestinal bacteria to fish growth, survival and immunity with and without bacterial infection [View project](#)



Marine oomycetes infection and bacterial infection of mud crab larvae in Universiti Malaysia Sabah [View project](#)



Data in Brief

Complete genome sequence of a giant Vibrio phage ValKK3 infecting *Vibrio alginolyticus*



Tamrin M. Lal^a, Motohiko Sano^b, Kishio Hatai^a, Julian Ransangan^{a,*}

^a Microbiology and Fish Disease Laboratory, Borneo Marine Research Institute, Universiti Malaysia Sabah, Jalan UMS, 88400, Kota Kinabalu, Sabah, Malaysia

^b Laboratory of Fish Pathology, Department of Marine Biosciences, Tokyo University of Marine Science and Technology, Room 223, Building No. 2 2F, Konan 4-5-7, Minato-ku, Tokyo 108-8477, Japan

ARTICLE INFO

Article history:

Received 29 February 2016

Accepted 8 March 2016

Available online 10 March 2016

Keywords:

Vibrio alginolyticus

Whole genome sequencing

Giant marine myophage

ABSTRACT

This paper describes the complete sequence of a giant lytic marine myophage, Vibrio phage ValKK3 that is specific to *Vibrio alginolyticus* ATCC[®] 17749[™]. Vibrio phage ValKK3 was subjected to whole genome sequencing on MiSeq sequencing platform and annotated using Blast2Go. The complete sequence of ValKK3 genome was deposited in DDBJ/EMBL/GenBank under accession number KP671755.

© 2016 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Specifications	
Organism	Myoviridae
Genus	T4-Like
Sequencer	Illumina (MiSeq)
Data format	Analyzed
Experimental factors	Bacteriophage strain
Experimental features	Whole genome analysis and gene annotation of Vibrio phage ValKK3
Consent	N/A
Sample source location	Marine sediment, Kota Kinabalu, N 06° 02.270', E 116° 06.710'

1. Direct link to deposited data

www.ncbi.nlm.nih.gov/genome/?term=KP671755

2. Materials and methods

Vibrio phage ValKK3 was isolated from marine sediment of Kota Kinabalu, Malaysia that was capable to lyse fish and human pathogen, *Vibrio alginolyticus* ATCC[®] 17749[™] [1]. The phage morphology was determined by a transmission electron microscope. The genome was extracted and purified using DNeasy Blood and Tissue Kit (Qiagen). The genome sequencing and assembly was performed by AITBiotech Pte. Ltd. (Singapore). The DNA library was prepared by Illumina Nextera XT kit. The samples were dual barcoded for multiplexing on the MiSeq instrument. The paired of 250 bp reads (2 × 250,000 samples) were

generated and demultiplexed corresponding to the barcodes. The FASQ files were generated on board of MiSeq Instrument. The sequence assembly was performed using de novo assembly using Velvet assembly software version 1.1 (Zerbino, European Bioinformatics Institute, UK). The assembly producing the lowest number of contigs with the largest n50 value was chosen for further assembly with a > 30-fold sequence coverage into single contig. The complete contig sequence was subjected to BLASTn [2] and the sequence was annotated using Blast2GO [3].




Fig. 1. Electron micrograph of negative-stained ValKK3. Bars = 100 nm.

* Corresponding author.

E-mail address: liandra@ums.edu.my (J. Ransangan).

Table 1
Genome comparison among giant vibriophages.

Bacteriophage strain	Vibrio phage ValKK3	Vibrio phage VH7D	Bacteriophage KVP40	Vibrio phage phi-pp2	Vibrio phage nt-1
Host species	<i>V. alginolyticus</i>	<i>Vibrio</i> strain 7D	<i>V. parahaemolyticus</i>	<i>V. parahaemolyticus</i>	<i>V. natriegens</i>
GenBank accession no.	KP671755	KC131129	AY283928	JN849462	HQ317393
Head length (nm)	140	155	140	150	120
Head diameter (nm)	79	75	70	90	70
Tail length (nm)	120	Not reported	130	120	110
Genome size (bp)	248,088	246,964	244,834	246,421	247,511
Blastn identity (%)	100	99	81	81	85
ORF number	390	378	386	383	379
G + C level (%)	41.2	41.3	42.6	42.7	41.3
Reference	This study	Luo et al. [4]	Miller et al. [5], Matsuzaki et al. [6]	Lin and Lin [7]	Comeau et al. [8], Zachary [9]

3. Data description

The morphological characteristics of ValKK3 (Fig. 1) showed that it belongs to the family of *Myoviridae*. To the best of our knowledge, among the giant T4-like phages reported so far (Table 1), the Vibrio phage ValKK3 is the only strain that can infect *V. alginolyticus*. The genome annotation analysis predicted 390 open reading frames which are numbered consecutively from ORF1 to ORF390 which represented 92.4% of the total genome. The putative genes varied from 111 bp (ORF268) to 4200 bp (ORF349). The protein function analysis showed that some CDSs involve in repression of host gene synthesis (ORF197, ORF202), defense system of own gene synthesis (ORF170, ORF176, ORF207) and quality control (ORF255), DNA synthesis (ORF61, ORF199, ORF367, ORF370, ORF241, ORF256, ORF381), DNA replication and recombination (ORF371, ORF372, ORF320, ORF178, ORF204, ORF106, ORF242, ORF257), structural protein: the head (ORF281–286), tail (ORF287–319) and tail fiber (ORF346–349), DNA packaging protein (ORF289, ORF290) and host lysis protein (ORF304). In this study, the putative functions of ValKK3 genome only represent 15.6% of the genes in the ValKK3. The other 84.4% in the ValKK3 are consisted of the genes which encoded proteins with unknown function or hypothetical proteins. The ValKK3 is similar to the Vibrio phage VH7D, Bacteriophage KVP40, Vibrio phage phi-pp2 and Vibrio phage nt-1 (Table 1). This assigns the ValKK3 to the T4-like phage genus of the *Myoviridae* family which belongs to the giant vibriophage group [7,8] with even larger genome size. The blast search revealed that ValKK3 was homologous to vibriophage that is known to infect some members of *Harveyi* clade bacteria. It is also revealed that T4-like phage is widely spread in the marine environment infecting different *Vibrio* species. Finally, an additional genome information of ValKK3 is hoped to provide positive contributions to the study of Vibrio bacteriophage.

4. Nucleotide accession number

This complete genome sequence has been deposited at DDBJ/EMBL/GenBank under accession no. KP671755.

Conflict of interest

The authors clarified that this work and writing has no conflict of interest.

Acknowledgments

This study was financially supported by the Universiti Malaysia Sabah's Research Priority Area Scheme (SBK0110-STWN-2013) and Malaysia Ministry of Education's Fundamental Research Grant Scheme (FRG0338-STWN-1/2013).

References

- [1] X.F. Liu, Y. Cao, H.L. Zhang, Y.J. Chen, C.J. Hu, Complete genome sequence of *Vibrio alginolyticus* ATCC 17749T. *Genome Announc.* 3 (1) (2015) e01500–e01514 (2015).
- [2] S.F. Altschul, W. Gish, W. Miller, E.W. Myers, D.J. Lipman, Basic local alignment search tool. *J. Mol. Biol.* 215 (1990) 403–410.
- [3] A. Conesa, S. Götz, J.M. García-Gómez, J. Terol, M. Talón, M. Robles, Blast2GO: universal tool for annotation, visualization and analysis in functional genomics research. *Bioinformatics* 21 (18) (2005) 3674–3676.
- [4] Z.H. Luo, Y.P. Yu, G. Jost, W. Xu, X.L. Huang, Complete genome sequence of a giant Vibrio bacteriophage VH7D. *Mar. Genomics* 3 (2015) 293–295.
- [5] E.S. Miller, J.F. Heidelberg, J.A. Eisen, W.C. Nelson, A. Scott Durkin, A. Ciecko, T.V. Feldblyum, O. White, I.T. Paulsen, W.C. Nierman, J. Lee, B. Szczypinski, C.M. Fraser, Complete genome sequence of the broad-host-range vibriophage KVP40: comparative genomics of a T4-related bacteriophage. *J. Bacteriol.* 185 (17) (2003) 5220–5233.
- [6] S. Matsuzaki, S. Tanaka, T. Koga, T. Kawata, A broad-host-range vibriophage, KVP40, isolated from sea water. *Microbiol. Immunol.* 36 (1992) 93–97.
- [7] Y.R. Lin, C.S. Lin, Genome-wide characterization of vibrio phage dpp2 with unique arrangements of the mob-like genes. *BMC Genomics* 13 (2012) 224.
- [8] A.M. Comeau, C. Arbiol, H.M. Krish, Composite conserved promoter-terminator motifs (PeSLs) that mediate modular shuffling in the diverse T4-like myovirus. *Genome Biol. Evol.* 6 (7) (2014) 1611–1619 (2014).
- [9] A. Zachary, Physiology and ecology of bacteriophages of marine bacterium *Beneckea natriegens*: salinity. *Appl. Environ. Microbiol.* 31 (3) (1976) 415–422.