

SEASONAL VARIATIONS AND POTENTIAL
SOURCES OF PM₁₀ IN MALAYSIA

JOYCE NGU SHU HUI

ENVIRONMENTAL SCIENCE PROGRAMME
FACULTY OF SCIENCE AND NATURAL RESOURCE
UNIVERSITI MALAYSIA SABAH

2014



UMS
UNIVERSITI MALAYSIA SABAH

SEASONAL VARIATIONS AND POTENTIAL SOURCES OF PM₁₀ IN MALAYSIA

JOYCE NGU SHU HUI
(BS 11110253)

THIS DISSERTATION IS SUBMITTED AS A PARTIAL REQUIREMENT TO OBTAIN
BACHELOR OF SCIENCE (Hons) DEGREE

ENVIRONMENTAL SCIENCE PROGRAM
FACULTY OF SCIENCE AND NATURAL RESOURCES
UNIVERSITI MALAYSIA SABAH

2014

DECLARATION

I affirm that this dissertation is of my own effort except the excerpt and summary which each of them has been referred to the sources mentioned in the reference section.

JOYCE NGU SHU HUI

BS11110253

26 JUNE 2014

AUTHENTICATION

VERIFICATION BY

SIGNATURE

SUPERVISOR

(DR. JUSTIN SENTIAN)

ACKNOWLEDGEMENT

I would like to express greatest gratitude to DR. JUSTIN SENTIAN for his professional advices and supports. Without his patient guidance, I would probably unable to get rid of the obstacles and solve problems in my research. Thanks to his great knowledge and always get the simple but useful solution to me.

I was also fortunate to have a postgraduate, Alan Yat who provided me assistance the most other than my supervisor. He was the person that I can refer to when my supervisor was outstation or busy.

Last but not least, I owe my profound gratitude to my family for supporting in my university life for all courses. It has encouraged me a lot.

VARIASI BERMUSIM DAN SUMBER BERPOTENSI PM10 DI MALAYSIA

ABSTRAK

Jerebu adalah kehadiran partikel halus dengan diameter 0.1 hingga 10 μm di udara dan menyebarkan dalam kepekatan yang tinggi dan majoritinya mengandungi PM10. Episod jerebu dan insiden serius ke atas Asia Tenggara termasuk Malaysia berlaku sejak akhir 1980-an. Kajian ini adalah untuk menganalisis kejadian jerebu di Malaysia dan mengesan punca pada tempoh yang disyaki tertentu. Hotspot pencemaran dan monsoon bermusim digunakan untuk menerangkan dan menjelaskan tempoh jerebu. Model HYSPLIT telah digunakan sebagai perisian trajektori jerebu dan empat stesen yang berbeza dengan lokasi geografi telah dipilih iaitu Kuala Lumpur, Pahang, Kuching, dan Kota Kinabalu. Dalam kajian ini, jerebu didapati jarang berlaku di Kota Kinabalu tetapi ia sering dilihat di Semenanjung Malaysia. Tempoh Jerebu dan corak di Semenanjung Malaysia adalah sama dan biasanya berlaku semasa tempoh monsoon musim panas. Sumber-sumber PM10 Semenanjung biasanya dari Sumatra, Indonesia. Kuching telah diperhatikan dengan beberapa kejadian jerebu pada Januari kepada Mac (2008) dan bulan Julai hingga September (2009 & 2010). Untuk Kota Kinabalu, tempoh masa didapati dengan peningkatan kepekatan PM10 besar. Ia berlaku pada bulan Februari untuk bulan April 2010. Ia disumbangkan oleh El Nino dan kemarau yang berpanjangan dari tahun 2009 hingga 2010. Monsun dan El Nino adalah kedua-dua faktor penting dalam menentukan arah tuju jerebu dan intensiti sebagai monsun perubahan laluan perjalanan bahan pencemar dan El Nino meningkatkan kekerapan pembakaran biomass.

ABSTRACT

Haze is defined as the presence of fine particles with diameter of 0.1 to 10 μm in the air which dispersed at high concentration caused white or brown vision (Soleiman *et al.*, 2003) and majority containing particulate matter (Rahman, 2013). The haze episodes and incidents were severe over the Southeast Asia including Malaysia since the late 1980s. This research is to analyze occurrence of haze in Malaysia and backward tracking its sources at the specific suspected period. Pollution hotspot and seasonal monsoon were used to further explain and clarify the haze period. HYSPLIT model was used as backward trajectory software and four different stations with geographical locations were selected which are Kuala Lumpur, Pahang, Kuching, and Kota Kinabalu. In my findings, haze seldom happened in Kota Kinabalu but it was frequently seen in Peninsular Malaysia. Haze period and pattern in Peninsular Malaysia was similar where it usually happened during period of summer monsoon. The sources of PM_{10} to Peninsular are usually from Sumatra, Indonesia. Kuching was observed with few haze outbreaks which is in January to March (2008) and July to September (2009 & 2010). For Kota Kinabalu, a period of time was found with tremendous increased PM_{10} concentration. It happened in February to April in 2010. It is contributed by occurrence of *El Nino* and prolonged dry spells from year 2009 to 2010. Monsoon and *El Nino* are both important factors in determining the direction of haze and its intensity as monsoon changes the travelling path of pollutants and *El Nino* increases frequency of forest burning.

CONTENTS

	Page
DECLARATION	ii
CERTIFICATION	iii
ACKNOWLEDGEMENT	iv
ABSTRAK	v
ABSTRACT	vi
LIST OF CONTENTS	vii
LIST OF TABLES	ix
LIST OF FIGURES	x
LIST OF SYMBOLS AND UNITS	xii
LIST OF ABBREVIATIONS	xiii
LIST OF APPENDICES	xv
CHAPTER 1 INTRODUCTION	1
1.1 Overview on Haze Pollution	1
1.2 Research Question	6
1.3 Objectives	6
1.4 Scope of Study	7
CHAPTER 2 LITERATURE REVIEW	
2.1 Air Quality Status in Malaysia	8
2.2 Biomass Burning	9
2.3 Haze	10
2.4 Regional Haze Transportation	14
2.4.1 Outflow of Haze into SEA	14
2.4.2 Inflow of Haze into SEA	14
2.5 Climate and Atmospheric Processes	15



2.5.1	Monsoon	15
2.5.2	Weather Patterns and its seasonal cycle	16
2.5.3	Maritime Continent Processes	17
2.5.4	<i>El Nino</i>	19
CHAPTER 3 METHODOLOGY		
3.1	Background	20
3.2	Data Collections	21
3.3	HYPPLIT Model Description	22
3.4	Operation of HYSPLIT Model	25
3.4.1	ARL's Ready Website	25
3.4.2	Graphical User Interface (GUI)	26
CHAPTER 4 RESULTS AND DISCUSSIONS		
4.1	Overview	27
4.2	Status of PM ₁₀ in Malaysia	28
4.3	Potential Sources of PM ₁₀	31
4.3.1	Kota Kinabalu	31
4.3.2	Kuching	33
4.3.3	Kuala Lumpur & Pahang	35
4.4	Monsoonal Variation on Haze	39
4.4.1	Monsoons and PM ₁₀ Concentration	39
4.4.2	<i>El Nino</i> Southern Oscillation (ENSO) Events	46
CHAPTER 5 CONCLUSION		
5.1	Conclusion	50
5.2	Future research	51
	REFERENCES	52
	APPENDICES	

LIST OF TABLES

Table No.		Page
Table 3.1	Overview of HYSPLIT (Source: UNIGE, 2010)	23
Table 3.2	Process and parameters accounted in HYSPLIT (Source: UNIGE, 2010)	23
Table 3.3	Particle granulometric in HYSPLIT model (Source: UNIGE, 2010)	23
Table 3.4	Meteorological data in HYSPLIT model (Source: UNIGE, 2010)	24
Table 3.5	Eruptive column (source term) of HYSPLIT model (Source: UNIGE, 2010)	24
Table 3.6	Possible output of HYSPLIT model (Source: UNIGE, 2010)	24

LIST OF FIGURES

Figure No.		Page
3.1	Malaysia map and locations selected were pointed with arrow	20
3.2	Flow diagram of operating web-based HYSPLIT model	25
3.3	Flow diagram of operating PC version HYSPLIT	26
4.1	Seasonal variation cycle of PM ₁₀ in four different stations from 2008 to 2010	30
4.2	Backward trajectories from Kota Kinabalu and map of fire hotspots in SEA from February to April 2010	32
4.3	Backward trajectories from Kuching and fire map hotspots in Southeast Asia during February 2008 and from July to September 2009	34
4.4	Backward trajectories from Kuala Lumpur and Pahang, and fire map hotspots in Southeast Asia from June to September 2009	36
4.5	Wind vectors from COADS climatology in Southeast Asia from January to June	37
4.6	Wind vectors from COADS climatology in Southeast Asia from July to December	38
4.7	Daily PM ₁₀ concentration during Summer Monsoons and Winter Monsoons from year 2007 to 2011 in Kota Kinabalu	42
4.8	Daily PM ₁₀ concentration during Summer Monsoons and Winter Monsoons from year 2007 to 2011 in Kuching	43
4.9	Daily PM ₁₀ concentration during Summer Monsoons and Winter Monsoons from year 2007 to 2011 in Pahang	44

4.10	Daily PM ₁₀ concentration during Summer Monsoons and Winter Monsoons from year 2007 to 2011 in Kuala Lumpur	45
4.11	Comparison of humidity and temperature among four years for Pahang station and Kuala Lumpur station	48
4.12	Comparison of humidity and temperature among four years for Kuching station and Kota Kinabalu station	49

LIST OF SYMBOLS AND UNITS

$^{\circ}\text{C}$	Degree Celsius
%	Percentage
$\mu\text{g}/\text{m}^3$	Microgram per cubic metre
μm	Micrometer
hrs	Hours
km	Kilometre
m	Metre



LIST OF ABBREVIATIONS

AOD	Aerosol optical depth
API	Air Pollution Index
Apr	April
ARL's Ready	Real-time Environmental Applications and Display system
ASMA	Alam Sekitar Sdn. Bhd.
Aug	August
CO	Carbon monoxide
COADS	Comprehensive Ocean -Atmosphere Data Set
Dec	December
DJF	December-January-February
DOE	Department of Environmental
ENSO	<i>El Nino</i> Southern Oscillation
Feb	February
GUI	Graphical user interface
HYSPLIT	Hybrid Single-Particle Lagrangian Integrated Trajectory
IOD	Indian Ocean Dipole
ITCZ	Intertropical convergence zone
Jan	January
Jul	July
Jun	June
LAS	Live Access Sever
Mar	March

MC	Maritime Continent
MJO	Madden Julian Oscillation
MLD	Mixing level depth
NO ₂	Nitrogen dioxide
NOAA	National Oceanic and Atmospheric Administration
Nov	November
Oct	October
PC	Personal Computer
PM	Particulate matter
PM ₁₀	Particulate matter < 10 µg/m ³
PM _{2.5}	Particulate matter < 2.5 µg/m ³
PSD	Particle size distribution
RM	Ringgit Malaysia
SEA	South-east Asia
Sept	September
SON	September-October-November
SST	Sea surface temperature
USD	United States Dollar
UV	Ultraviolet
VOC	Volatile organic compounds
WWF	World Wildlife Fund for Nature

LIST OF APPENDICES

Appendix No.		Pages
A	Methodology HYSPLIT	59
B	Table: Average concentration of PM ₁₀ at four different stations in year 2008, 2009, 2010, and 2011	63
C	Table: PM ₁₀ variation in monsoons at Kuching	64
D	Table: PM ₁₀ variation in monsoons at Pahang	65
E	Table: PM ₁₀ variation in monsoons at Kota Kinabalu	66
F	Table: PM ₁₀ variation in monsoons at Kuala Lumpur	67
G	Table: Monthly data in Kuching, Pahang, Kota Kinabalu and Kuala Lumpur	68

CHAPTER 1

INTRODUCTION

1.1 Overview on Haze Pollution

Air pollution is the presence of chemicals in atmosphere in sufficient quantities and duration which cause harm to abiotic and biotic (Miller, 2004). These chemical gases or particles in air are called air pollutants (Kaufmann and Cleveland, 2008). It can be defined as primary pollutants which are emitted directly from combustion of fuels and industrial operations and secondary pollutants form where the primary pollutants react with one another or with basic components in atmosphere. Long-lived primary and secondary air pollutants might spread by prevailing winds from its origin areas to country-side and to downwind urban and rural areas. These pollutants are influenced not only by the rate of emission but also interference by wind speed, turbulence level, air temperature and precipitation (Dominick *et al.*, 2012). Man-made urban air pollution is a complex mixture with toxic compounds which are considered as particulate matter (PM) usually comes from combustion processes (Cohen *et al.*, 2004). Particulate matter with less than 10 μm of its diameter (PM_{10}) has been known as an important atmospheric pollutant and act as main contributor of haze. This particulate matter pollutant in air is the combination

of solid and liquid particles. Majority sources of particulate matter were contributed by biomass burning.

Haze is air pollution which can be carried by wind from its origin and often to hundreds or thousands of miles (EPA, 1999). Haze is defined as the presence of fine particles with diameter of 0.1 to 1.0 μm in the air which dispersed at high concentration caused white or brown visible in view (Soleiman *et al.*, 2003). It contains dust and smoke particles together with air pollutants such as sulphur dioxide, nitrogen oxide, ozone, carbon monoxide and majority containing particulate matter (Rahman, 2013). It can be as fine as droplets of liquids in fogs and mists or as solid particles such as soot (Mahajan, 2003). Haze particles not only influence visibility of atmosphere, but also solar radiation and even can involve further chemical reactions in atmosphere to produce secondary pollutants (Roosli *et al.*, 2001). When sunlight (solar radiation) encounters tiny pollution particles in the air, some of the light is reflected, scattered, or absorbed by haze particles before it reaches an observer (EPA, 1999). Light absorption by particles which also called light absorbing particles is primary contains elemental carbon.

Particles of haze generally come from many sources, some from natural and some from anthropogenic. Anthropogenic activities especially biomass burning including forest fires, open burning and agricultural land clearings are the main causative factors for deteriorating of air quality and visibility (Rahman, 2002). Fire can influence air quality of both local and regional (McKenzie *et al.*, 2006). During the 20th century, biomass combustion due to man-kind activities has been more vigorous and fire activities have been annual phenomena over the last 30 years (Field *et al.*, 2009). Biomass burning releases more air pollutants compare to well control combustion sources on a mass of pollutants per mass of fuel basis (Lin *et al.*, 2007; Shih *et al.*, 2008). Beside, fires activities not only cause haze pollutions, but also release large amounts of carbon into the atmosphere (Page *et al.*, 2002). In western region of the United State, it was found that total carbon was the major contributor to $\text{PM}_{2.5}$ concentrations and elevated contributions at the upper extremes of aerosol fine mass probably correspond to biomass burning (Ames and Malm, 2001).

Therefore, biomass burning is known as the major source of global greenhouse gases and hazardous aerosols (Andreae and Merlet, 2001; Lee *et al.*, 2011).

Meteorological conditions are important role in transportation, diffusion and deposition the air pollutants (Chen *et al.*, 2007; Zhu *et al.*, 2010). The major weather system such as stable continental high-pressure and low-pressure systems will make favorable for meso-scale and large scale of heavy haze pollutions form. Beside, as haze concentration is high, it means that the particular area is with higher pollutants and humidity level. Relative humidity affect the light extinction effect, which either by hydration of dry particles when the humidity is high or by condensation of water vapor to droplets which often occur of fog (Elias *et al.*, 2009; Winkler, 1988).The aerosols component are hygroscopic especially sulfates and nitrate particles, usually grow in size as humidity increases by accumulating water from the atmosphere and cause further impairing visibility. Besides, several factors such as prolonged dry weather and stable atmosphere are the supreme ingredients for the haze formation (Rahman, 2013). Wind and weather are playing an important part on transporting the pollution from one place to another. A lower surface wind speed and stably stratified atmosphere can lead to accumulation of air pollutants. Somewhere, the most important reasons of long-lasting haze pollution process happen due to accumulation of air pollutants near the surface caused by low mixing level depth (MLD). However, usually, as a cold front passed through the region polluted place, the pollutants can be induced due to strong wind and precipitation (Kang *et al.*, 2013). Particles in haze are eliminated from atmosphere mainly due to the precipitation. The interaction between rainwater and haze caused chemical degradation and influences the pH values (Tanner *et al.*, 1997). Deposition of the particles to the soil and water were increase during the period of haze (Norela *et al.*, 2013).

Haze is an indicator of high concentration of particulate matter, therefore, it has potential to harm and cause impact to public health, ecological systems and climate (Tie *et al.*, 2009; Chameides *et al.*, 1999; Solomon *et al.*, 2007). The particles

in haze can increase physical health problems, allergies, cancer as well as pre-term delivery. It can influence the health either local or systemic effect. Local effect can result in eye, nose and throat irritation while systemic effect can be range from respiratory conditions to worsening of heart diseases. Transportation accidents increase and lower vegetation productivity occur during haze periods (Radojevic and Hassan, 1999). Smoke aerosols are the particulate matter in air which is mainly released by biomass burning influence directly and indirectly to temperature, atmospheric circulation, clouds formation, precipitation and climate (Xian *et al.*, 2013).

The haze episodes and incidents were severe over the Southeast Asia including Malaysia since the late 1980s. Most of the air quality studies in Malaysia are related to the occurrence of haze episodes (Cheang, 1991; Awang *et al.*, 1997, Juneng *et al.*, 2011). Malaysia is one the world's twelve areas of mega-biodiversity and act as home to a number of species disproportionate to its geographical size (CIJ, 2007). Land use change in Malaysia from forest to plantations, mainly oil palm remains the major threat to biodiversity. Activities such as forest fires, industrial emissions and transport emissions worsen the quality of air. Occurrence of haze has threatened precious natural environment of Malaysia. Particulate matter is one of the major air pollutants and is conclusive in the computation of Air Pollution Index (API) (Afroz *et al.*, 2003). API is simple, generalized way and provides easily understandable information to describe the air quality used in Malaysia, Singapore and China (Hajah, 1997). API level less than 100 consider good and safe air while API level higher than 100 consider the haze pollution process (Kang *et al.*, 2013). API level in the range of 200 to 300 is consider very unhealthy air and API level higher than 300 is consider hazardous air.

According to Dominick (2012), Malaysia is known as a tropical climate country which experience constant temperatures and continuous high relative humidity. High humidity usually gives a result of high precipitation. Wind is generally light and flexible throughout the country as Malaysia located near the equator. The wind flow

patterns change with four seasons which are the Northeast Monsoon from November to March, a transitional period from April and May, the Southwest Monsoon from June to September and a second transitional period from October to November. The air pollutants usually move with winds flow base on northeast monsoon and southwest monsoon. High amount of particulate matter from biomass burning in Sumatera and Kalimantan, Indonesia follow wind pattern of southwest monsoon to Malaysia while during northeast monsoon, wind bring particulate matter from Indochina region to Peninsular Malaysia (Juneng *et al.*, 2009).

Transportation of particulate matter can be calculated in various scales by using various trajectory models such as Ash3d, PUFF, ATHAM, and HYSPLIT. Hybrid Single-Particle Lagrangian Integrated Trajectory (HYSPLIT) model is known as a complete system for computing both simple air parcel trajectories and complex dispersion and deposition simulations (NOAA, 2011). Through National Oceanic and Atmospheric Administration (NOAA) and Australia's Bureau of Meteorology cooperation, this model uses advection algorithms, updated stability and dispersion equations, a graphical user interface, and the option to include modules for chemical transformations. Air Resources Laboratory's HYSPLIT model is a tool that helps to explain how, where, and when chemicals and materials are atmospherically transported, dispersed, and deposited. It is designed to support a wide range of simulations which associated to the atmospheric transport and dispersion of pollutants and hazardous materials along with its deposition to the Earth's surface. Trajectory can show local emission and regional transport (Kang *et al.*, 2013). The transport pathway of air parcels pass through the region places can be detected by HYSPLIT model. Backwards trajectory analysis can use this model as well to estimate the central path from the original air mass to the receptor (Chang *et al.*, 2013).

1.2 Research Question

Haze episodes over the Southeast Asia region has been recurring in unprecedented scale. It happened in greater extent and intensity, longer duration and impacted larger community. Malaysia is one of the countries which experiencing the haze pollution disaster. Malaysia is made up of Peninsula, Sarawak and Sabah which characterized with different geographic locations. There are several questions to ponder in this research. Firstly, where is the haze pollution coming from? Is it originating from within Malaysia or neighboring countries for example Indonesia, Thailand and etc.? How do Peninsula and Borneo vary in timing during regional scale haze incident? Which places in Malaysia are considered the pollution hotspots? How is the transportation route of air pollutants differ in both dominant monsoons?

1.3 Objectives

In order to ponder and study the uncertainties stated in the research questions, this research is designed to achieve the following objectives:

- To develop backward trajectory of haze in Malaysia in different locations and during different monsoon seasons
- To determine potential sources of PM_{10}
- To determine the role of monsoon seasons to the haze coverage and transport over Malaysia region

1.4 Scope of Study

In this research, two locations in Peninsula Malaysia and two locations in East Malaysia are chosen. In Peninsula Malaysia, the selected locations were Kuala Lumpur (3.1357° N, 101.6880° E) and Pahang (3.7500° N, 102.5000°N). While in East Malaysia, Kuching (1.5600° N, 110.3450° E) and Kota Kinabalu (5.9714° N, 116.0953° E) were selected. In order to perform trajectory analysis, Hybrid Single-Particle Lagrangian Integrated Trajectory (HYSPPLIT) model will be employed to determine the long range transport of pollutants over Malaysia.

CHAPTER 2

LITERATURE REVIEW

2.1 Air Quality Status in Malaysia

In an undisturbed environment with no human interference, fire activity is a rare occurrence in humid tropical region of insular South-east Asia (SEA) (Goldammer, 2006). Fire can influence air quality of both local and regional (McKenzie *et al.*, 2006). Regional haze form as area burned and fire severity increase cause the increase of biomass consumption and smoke emissions which consequently release more atmospheric dispersion of particulates and aerosols (McKenzie *et al.*, 2006). The haze episode and incident is severe over the Southeast Asia including Malaysia since the late 1980s. Haze incident in the 1997 is the most extreme due to association of *El Nino* event. Most of the air quality studies in Malaysia are related to the occurrence of haze incident (Cheang, 1991; Awang *et al.*, 1997, Juneng *et al.*, 2011). The haze incident 1997 is enormous and its extent has influenced over the region. About 70% of total emissions are vehicular emission during the non-haze period. Awang *et al.* (2000) states that the smoke haze is mainly consist of abundant suspended microparticulate matter and also associated with other gaseous pollutant. During the haze occasion in 1997, study shows the PM₁₀ concentration has increased four times higher in Klang and 20 times more in Kuching in comparable to the Malaysian Air Quality Guideline 1989 (Awang *et al.*, 2000).



References

- Afroz, R., Hassan, M.N. & Ibrahim, N. 2003. Review of air pollution and health impacts in Malaysia. *Environmental Research*, **92**: 71-77.
- Ames, R.B. & Malm, W.C. 2001. Chemical species' contributions to the upper extremes of aerosol fine mass. *Atmos. Environ.*, **35**: 5193–5204.
- Andreae, M.O. & Merlet, P. 2001. Emission of trace gases and aerosols from biomass burning. *Global Biogeochem.*, **15**: 955-966.
- Asian Development Bank (ADB). 1999. Planning for Fire Prevention and Drought Management Project, Final Report. Jakarta, Indonesia.
- Chan, C.Y., Chan, L.Y., Zheng, Y.G., Harris, J.M., Oltmans, S.J. & Christopher, S.2001. Effects of 1997 Indonesian forest fires on tropospheric ozone enhancement, radiative forcing, and temperature change over the Hong Kong region. *Journal of Geophysical Research D: Atmospheres*, **106**: 14875-14885.
- Chang, S.S., Lee, W.J., Wang, L.C., Lin, N.H. & Chang, C.G.P. 2013. Influence of the Southeast Asian biomass burnings on the atmospheric persistent organic pollutants observed at near sources and receptor site. *Atmospheric Environment*, **78**: 184-194.
- Chen, Z.H., Cheng, S.Y., Su, F.Q., Gao, Q.X., Yu, T. & Ren, Z.H. 2007. Analysis of large-scale weather pattern during heavy air pollution process in Beijing. *Res. Environ. Sci.*, **20** (2): 99–105 (in Chinese).
- CIJ. 2007. A haze of secrecy: Access to Environmental Information in Malaysia. The Centre for Independent Journalism, Malaysia (CIJ), ISBN: 978-1-902598-86-4.
- Cohen, A. J., Ross, H. A., Bart, O. Pandey, K. D., Krzyzanowski, M., Künzli, N., Gutschmidt, K., Arden Pope III, C., Isabelle, R., Samet, J. M. & Smith, K. R. 2004. Urban air pollution. Chap. 17, pp 1353 - 1434.
- Dominick, D., Latif, M. T., Juahir, H., Aris, A. A. & Zain, S. M. 2012. An assessment of influence of meteorological factors on PM10 and NO₂ at selected stations in Malaysia. *Sustain. Environ. Res*, **22**(5), 305-315.

- Eck, T.F., Holben, B.N., Reid, J.S., O'Neill, N.T., Schafer, J.S., Dubovik, O., Smirnov, A., Yamasoe, M.A. & Artaxo, P. 2003. High aerosol optical depth biomass burning events: a comparison of optical properties for different source regions. *Geophys. Res. Lett.*, **30** (20): 2035.
- Edward, J.H. & Boon, N.C. 2010. Aerosol transport model evaluation of an extreme smoke episode in Southeast Asia. *Atmospheric Environment*, **44**: 1422-1427.
- Elias, T., Haeffelin, M., Drobinski, P., Gomes, L., Rangognio, J., Bergot, T., Chazette, P., Raut, J.C. & Colomb, M. 2009. Particulate contribution to extinction of visible radiation: pollution, haze, and fog. *Atmospheric Research*, **92**: 443-454
- EPA. 1999. Haze: How air pollution affects the view. United States Environmental Protection Agency (EPA). EPA-456/F-99-001.
- Field, R.D., Werf, W.D.G.R. & Shen, S.S.P. 2009. Human amplification of drought-induced biomass burning in Indonesia since 1960. *Nat. Geosci.*, **2**: 185–188.
- Goldammer, J.G. 2006. History of equatorial vegetation fires and fire research in Southeast Asia before the 1997/98 episode: a reconstruction of creeping environmental changes. *Mitig. Adapt. Strateg. Glob. Chang.*, **12**: 13–32.
- Hajah, R.I. 1997. A Guide to Air Pollutant Index in Malaysia (API). Department of Environment.
- Ho, C. H., Baik, J. J., Kim, J. H., Gong, D. Y. & Sui, C. H. 2004. Interdecadal changes in summertime typhoon tracks. *Journal of Climate*, **17**, 1767-1776.
- Hong, S.B., Kim, D.S., Ryu, S.Y., Kim, Y.J. & Lee, J.H. 2008. Chemical characteristics of PM_{2.5} ions measured by a semicontinuous measurement system during the fall season at a suburban site, Gwangju, Korea. *Atmos. Res.*, **89**: 62–75.
- Hyer, E.J. & Chew, B.N. 2010. Aerosol transport model evaluation of an extreme smoke episode in Southeast Asia. *Atmospheric Environment* **44**, 1422-1427.
- Juneng, L., Latif, M.T., Tangang, F.T. & Mansor, H. 2009. Spatio-temporal characteristics of PM₁₀ concentration across Malaysia. *Atmos. Environ.*, **43** (30): 4584-4594.

- Juneng, L., Latif, M. T. & Tangang, F. 2011. Factors influencing the variations of PM₁₀ aerosol dust in Klang Valley, Malaysia during the summer. *Atmospheric Environment*, **45**: 4370-4378
- Jung, J., Lee, H., Kim, Y. J., Liu, X., Zhang, Y., Gu, J. & Fan, S. 2009. Aerosol chemistry and the effect of aerosol water content on visibility impairment and radiative forcing in Guangzhou during the 2006 Pearl River Delta campaign. *J. Environ. Manage.*, **90**: 3231–3244.
- Kang, H., Zhu, B., Su, J., Wang, H., Zhang, Q. & Wang, F. 2013. Analysis of a long-lasting haze episode in Nanjing, China, Department of Atmospheric Physics, Nanjing University of Information Science & Technology, Nanjing 210044, China. *Atmospheric Research*, **120–121**: 78–87.
- Kaufmann, R. K. & Cleverland, C. J. 2008. *Environmental Science*, McGraw-Hill. ISBN 978-0-07-298429-3.
- Keyword, M.D., Ayers, G.P., Gras, J.L., Boers, R. & Leong, C.P. 2003. Haze in the Klang Valley of Malaysia. *Atmos. Chem. Phys.* 3, 591–605.
- Kim, O.N.T. & Leelasakultum, K. 2011. Analysis of meteorology and emission in haze episode prevalence over mountain-bounded region for early warning. *Science of the Total Environment*, **409**: 2261-2271.
- Lam, K. S., Wang, T. J., Chan, L. Y., Wang, T. & Harris, J. 2001. Flow pattern influencing the seasonal behavior of surface ozone and carbon monoxide at a coastal site near Hong Kong. *Atmospheric Environment*, 35, 3121-3135.
- Latif, M. T., Lim, S. H. & Juneng, L. 2012. Variations of surface ozone concentration across the Klang Valley, Malaysia. *Atmospheric Environment*, 61, 434-435.
- Lee, C.T., Chuang, M.T., Lin, N.H., Wang, J.L., Sheu, G.R., Chang, S.C., Wang, S.H., Huang, H., Chen, H.W., Liu, Y.L., Weng, G.H., Lai, H.Y. & Hsu, S.P. 2011. The enhancement of PM_{2.5} mass and water-soluble ions of biosmoke transported from Southeast Asia over the Mountain Lulin site in Taiwan. *Atmos. Environ.*, **45**: 5784-5794.

- Lee, K.H., Kim, Y.J. & Kim, M.J. 2006. Characteristics of aerosol observed during two severe haze events over Korea in June and October 2004. *Atmos. Environ.*, **40**: 5146–5155.
- Lin, L.F., Lee, W.J., Li, H.W., Wang, M.S. & Chang, C.G.P. 2007. Characterization and inventory of PCDD/F emissions from coal-fired power plants and other sources in Taiwan. *Chemosphere*, **68**: 1642-1649.
- Ling, A.H. 2012. Weather effects on palm oil production: supply outlook 2012/2013. Malaysia Palm Oil Conference. Kuala Lumpur.
- Liousse, C., ByrneDevaux, C. & Cachier, H. 1995. Aging of savannah biomass burning aerosols: consequences on their optical properties. *J. Atmos. Chem.*, **22**, 1–17.
- Mahajan, S.P. 2003. Air Pollution Control. Capital publishing Company. Centre for Ecological Sciences Indian Institute of Science, Bangalore.
- McBride, J., Drosdowsky, W., Kirono, D, Gunawan, D., Soetamato & Winarso, A.P. 1998. Inter-annual variability of the Indonesian Monsoon, Proc. International Conference on Monsoon and Hydrologic Cycle. Korean Met. Soc. 39–42.
- McKenzie, D., O'Neill, S.M., Larkin, N.K. & Norheim, R.A. 2006. Integrating models to predict regional haze from wildland fire. *Ecological modelling*, **199**: 278–288.
- MdYusoh, N.F.F., Ghazali, N.A., Ramli, N.A., Yahaya, A.S., Sansuddin, N. & Al Madhoun,W. 2008. Correlation of PM10 concentration and weather parameters in conjunction with haze event in Seberang Perai, Penang. Proceedings International Conference on Construction and Building Technology, D(20), pp. 211-220.
- Miettinen, J. & Liew, S.C. 2009. Burn-scar patterns and their effect on regional burnt-area mapping in insular South-east Asia. *Int. J. Wildland Fire*, **18**: 837–847.
- Miller, G.T. 2004. *Working with the earth, Environmental Science*, 10th Ed. Thompson Brooks/Cole, ISBN 0-534-42407-4.
- Nichol, J. 1998. Smoke Haze in Southeast Asia: A predictable recurrence. *Atmospheric Environment*, PII S1352-2310, 00086-7

- NOAA. 2011. Air Resources Laboratory HYSPLIT Model Research. National Oceanic and Atmospheric Administration (NOAA), U.S Department of Commerce.
- Noh, Y.M., Muller, D., Shin, D.H., Lee, H., Jung, J.S., Lee, K.H., Cribb, M., Li, Z. & Kim, Y. J. 2009. Optical and microphysical properties of severe haze and smoke aerosol measured by integrated remote sensing techniques in Gwangju, Korea. *Atmos. Environ.*, **43**: 879–888.
- Norela, S., Saidah, M.S. & Mahmud, M. 2013. Chemical composition of the haze in Malaysia 2005. *Atmospheric Environment*, **77**: 1005-1010.
- Page, S.E., Siegert, F., Rieley, J.O., Boehm, H.D.V., Jaya, A. & Limin, S. 2002. The amount of carbon released from peat and forest fires in Indonesia during 1997. *Nature*, **420**: 61–65.
- Radojevic, M. & Hassan, H. 1999. Air quality in Brunei Darussalam during the 1998 haze episode. *Atmospheric Environment*, **33**: 3651-3658.
- Radojevic, M. 1997. The haze in Southeast Asia. *Environmental Scientist*, **6** (6): 1-3.
- Rahman, A. K. 2002. Observation of PM₁₀ readings in relation to forest fire events from ASMA's continuous air quality monitoring stations. In: World Conference on Land and Forest Fire Hazards 2002, Kuala Lumpur. 155-170.
- Rahman, A.H. 2013. Haze Phenomenon in Malaysia: Domestic or Transboundary Factor?. 3rd International Journal Conference on Chemical Engineering and its Applications (ICCEA'13), 28-29 September 2013, Phuket, Thailand.
- RAND. 2002. The Dangers of Smoke Haze: Mortality in Malaysia from Indonesian Forest Fires. *Labor and Population Program*, RB-5066.
- Reid, J.S. & Hobbs, P.V. 1998. Physical and optical properties of young smoke from individual biomass fire in Brazil. *J. Geophys. Res.*, **103**: 30013–30031.
- Reid, J.S., Hobbs, P.V., Ferek, R.J., Blake, D.R., Martins, J.V., Dunlap, M.R. & Liousse, C. 1998. Physical, chemical, and optical properties of regional hazes dominated by smoke in Brazil. *J. Geophys. Res.*, **103**: 32059–32080.
- Reid, J.S., Hyer, E.J. & Prins, E.M. 2009. Global monitoring and forecasting of biomass burning smoke: description and lessons from the Fire locating and

- Modeling of Burning Emission (FLAMBE) program. *J. Sel. Top. Appl. Earth Obs. Rem. Sens*, **2**: 144-162.
- Roosli, M., Theis, G., Kunzli, N., Staehelin, J., Mathys, P., Oglesby, L., Camenzind, M. & Braun-Fahrlander, C. 2001. Temporal and spatial variation of the chemical composition of PM₁₀ at urban and rural sites in the Basel area, Switzerland. *Atmos. Environ.*, **35**: 3701-3713.
- See, S.W., Balasubramanian, R. & Wang, W. 2006. A study of the physical, chemical, and optical properties of ambient aerosol particles in Southeast Asia during hazy and non-hazy days. *J. Geophys. Res.-Atmos.*, **111**: 10.
- Shahwahid, M.H.O. & Othman., J. 1999. "Malaysia" in Indonesia's Fire and Haze, the cost of catastrophe, Glover D and Jessup. The Institute of Southeast Asia Studies (ISEAS), 22-50.
- Shih, S.I., Lee, W.J., Lin, L.F., Huang, J.Y., Su, J.W. & Chang-Chien, G.P. 2008. Significance of biomass open burning on the levels of polychlorinated dibenzo-p-dioxins and dibenzofurans in the ambient air. *J. Hazard. Mater.*, **153**: 276-284.
- Sirinanda, K.U. 1990. Rainfall seasonality in Negara Brunei Darussalam. *Asian Geographer*, **9**: 39-52.
- Soleiman, A., Ohtman, M., Samah, A.A., Sulaiman, N.M. & Radojevic.M. 2003. The occurrence of haze in Malaysia: a case study in an urban industrial area. *Pure Appl. Geophys.*, **160**: 221-238.
- Sundarambal, P., Balasubramanian, R., Tklich, P. & He, J. 2010. Impact of biomass burning on ocean water quality in Southeast Asia through atmospheric deposition: field observations. *Atmos. Chem. Phys.*, **10**: 11323-11336.
- Tacconi, L., Jotzo, F. & Grafton, R.Q. 2006. Local causes, regional co-operation and global financing for environmental problems: the case of Southeast Asian haze pollution.
- Tanner, P.A. & Wong, Y.S. 1997. Atmospheric gases, particulates and rainfall concentrations during summer rain events. *Inter. J. Environ. Anal. Chem.*, **67**: 185-202.

- Tayeh, S.M.O. & Ramli, N.A. 2012. High particulate events in Southeast Asia and their impacts: Review. The 4th International Engineering Conference.
- Tosca, M.G., Randerson, J.T., Zender, C.S., Nelson, D.L., Diner, D.J. & Logan, J.A. 2011. Dynamics of fire plumes and smoke clouds associated with peat and deforestation fires in Indonesia. *J. Geophys. Res.*, **116**:D08207.
- UNEP. 2003. Asia Toolkit Project on Inventories of Dioxin and Furan Releases National PCDD/PCDF Inventories, p. 18 of Vietnam 2002.
- Vedal, S. & Dutton, S.J. 2006. Wildfire air pollution and daily mortality in a large urban area. *Environ. Res.*, **102**: 29-35.
- Westphal, D.L. & Toon, O.B., 1991. Simulations of microphysical, radiative, and dynamical processes in a continental-scale forest fire smoke plume. *J. Geophys. Res.*, **96**: 22379–22400.
- Winkler, P. 1988. The growth of atmospheric aerosol particles with relative humidity. *Physica Scripta*, **37**: 223.
- Xian, P., Reid, J.S., Atwood, A.S., Johnson, R.S., Hyer, E.J., Westphal, D.L. & Sessions, W. 2012. Smoke aerosol transport patterns over the Maritime Continent. *Atmospheric Research*, **122**: 469-485.
- Yusof, N.F.F., Ghazali, N.A., Ramli, N.A., Yahya, A.S. & Al Madhoun, W. 2008. Correlation of PM₁₀ concentration and weather parameters in conjunction with haze event in Seberang Perai, Penang. International Conference on Construction and Building Technology **20**, 211-220
- Zheng, Y.G., Zhu, P.J., Chan, C.Y., Chang, L.Y., Cui, H., Zheng, X.D., Zhao, Q. & Qin, Y. 2004. Influence of biomass burning in Southeast Asia on the lower tropospheric ozone distribution over South China. *Acta Geophysica Sinica*, **47**: 767-775.
- Zhu, B., Su, J.F., Han, Z.W., Yin, C. & Wang, T.J. 2010. Analysis of a serious air pollution event resulting from crop residue burning over Nanjing and surrounding regions, China. *Environ. Sci.*, **30** (5): 585–592 (in Chinese).