

**ISOLATION AND CHARACTERIZATION OF  
cDNA SEQUENCES ENCODING FOR MADS-  
BOX TRANSCRIPTIONAL FACTORS  
FROM RICE LEAVES**

**HELENA BIUN**

**SCHOOL OF SCIENCE AND TECHNOLOGY  
UNIVERSITI MALAYSIA SABAH  
2007**



**UMS**  
UNIVERSITI MALAYSIA SABAH

**ISOLATION AND CHARACTERIZATION OF  
cDNA SEQUENCES ENCODING FOR MADS-  
BOX TRANSCRIPTIONAL FACTORS  
FROM RICE LEAVES**

**HELENA BIUN**

**THESIS SUBMITTED IN PARTIAL  
FULFILLMENT FOR THE DEGREE OF  
MASTER OF SCIENCE**

**SCHOOL OF SCIENCE AND TECHNOLOGY  
UNIVERSITI MALAYSIA SABAH  
2007**



**UMS**  
UNIVERSITI MALAYSIA SABAH

**UNIVERSITI MALAYSIA SABAH**

**BORANG PENGESAHAN STATUS TESIS®**

**JUDUL : ISOLATION AND CHARACTERIZATION OF cDNA SEQUENCES  
ENCODING FOR MADS-BOX TRANSCRIPTIONAL FACTORS FROM  
RICE LEAVES**

**IJAZAH : SARJANA SAINS (BIOTEKNOLOGI TUMBUHAN)**

**SESI PENGAJIAN : 2000-2007**

Saya, HELENA BIUN mengaku membenarkan tesis Sarjana ini disimpan di Perpustakaan Universiti Malaysia Sabah dengan syarat-syarat kegunaan seperti berikut:

1. Tesis adalah hakmilik Universiti Malaysia Sabah
2. Perpustakaan Universiti Malaysia Sabah dibenarkan membuat salinan untuk tujuan pengajian saya
3. Perpustakaan dibenarkan membuat salinan tesis ini sebagai bahan pertukaran antara institusi pengajian tinggi
4. TIDAK TERHAD

Disahkan oleh

\_\_\_\_\_  
(Penulis: HELENA BIUN)

\_\_\_\_\_  
(TANDATANGAN PUSTAKAWAN)

Alamat:  
Peti Surat 293,  
89507 Penampang.

\_\_\_\_\_  
(Penyelia: Lt. Kol. Prof. Datuk Dr. Kamaruzaman Ampon)

Tarikh: 01 Julai 2008

Tarikh: \_\_\_\_\_

\_\_\_\_\_  
(Penyelia: Dr. Lee Ping Chin)

Tarikh: \_\_\_\_\_



\_\_\_\_\_  
(Penyelia: Dr. Zaleha Abdul Aziz)

Tarikh: \_\_\_\_\_

\_\_\_\_\_  
(Penyelia: Dr. Douglas B. Furtek)

Tarikh: \_\_\_\_\_

**CATATAN:** <sup>®</sup> Tesis dimaksudkan sebagai tesis Ijazah Doktor Falsafah dan Sarjana secara penyelidikan atau disertasi bagi pengajian secara kerja kursus dan penyelidikan, atau laporan Projek Sarjana Muda (LPSM)



## DECLARATION

I hereby declare that the material in this thesis is my own except for quotations, excerpts, equations, summaries and references, which have been duly acknowledged.

30 April 2007

---

Helena Biun  
PS00-001-226



## CERTIFICATION

TITLE : **ISOLATION AND CHARACTERIZATION OF cDNA SEQUENCES ENCODING FOR MADS-BOX TRANSCRIPTIONAL FACTORS FROM RICE LEAVES**

DEGREE : **MASTER OF SCIENCE**

VIVA DATE : **31 DECEMBER 2007**

### DECLARED BY

- 1. CHAIRMAN OF SUPERVISORY COMMITTEE**  
(LT. KOL. PROF. DATUK DR. KAMARUZAMAN AMPON)
- 2. MEMBER OF SUPERVISORY COMMITTEE**  
(DR. LEE PING CHIN)
- 3. MEMBER OF SUPERVISORY COMMITTEE**  
(DR. ZALEHA ABDUL AZIZ)
- 4. MEMBER OF SUPERVISORY COMMITTEE**  
(DR. DOUGLAS B. FURTEK)



## ACKNOWLEDGEMENT

First and foremost, I thank God for the completion of the thesis. I would like to give my thanks to Dr. Douglas Bruce Furtek for teaching me with a lot of patience and for valuable technical advice and discussions throughout this study. I'm also very grateful to him for letting me to work at the Malaysian Cocoa Board. I would like to give my sincere thanks to Dr. Lee Ping Chin, Dr. Zaleha Abdul Aziz, Prof. Datuk Dr. Kamaruzzaman Ampon and Dr. Jualang Azlan Gansau for many helpful suggestions and advice during the course of this work, especially in the acquisition of the research grant. I'm also indebted to Prof K. Paranjothy for his valuable technical advice and discussions on any topic when I was in doubt of and not forgetting his faith in me to complete this study. I am grateful to the members of the MADS group, Jagdish Kaur Chahil, Fiana Yong, Roslina Mohd. Shah, Norasekin Tamchek and Tan Su Hui, for their support and encouragement during our ups and downs in the lab. My sincere thanks and appreciation also goes out to other lab members especially Adrian Ng Wui Ming and Awang Mohammad Sagaf, for all their assistance and free supplies. I also want to thank Helda Souki for helping me with the thesis preparation. I would like to thank my dear husband, Carlos and my dear children, Clevelence and Hayden, for their understanding, encouragement and financial support. To my parents, my brother and my sisters, thank you for your strong support and constant encouragement. Finally, I would like to thank Universiti Malaysia Sabah for providing me the opportunity to work in the School of Science and Technology.



## ABSTRACT

### ISOLATION AND CHARACTERIZATION OF cDNA SEQUENCES ENCODING FOR MADS-BOX TRANSCRIPTIONAL FACTORS FROM RICE LEAVES

MADS-box genes are characterized by the conserved MADS domain and found in a wide range of eukaryotes including metazoans, fungi, slime mold, and green plants. Throughout plant evolution, MADS-box genes have been recruited as transcriptional regulators active in the development of diverse plant structures. In monocots, MADS-box genes have been identified in maize, rice, and orchids which play important roles during reproductive and vegetative developmental processes. Although rice MADS-box genes involved in flower and fruit development have been well characterized, rice MADS-box genes expressed in vegetative structures have yet to be explored. The aim of this project is to isolate and analyze the cDNA sequences encoding for putative MADS-box transcription factors from MR84 leaves. MR84 is currently the popular local cultivar which produced good quality and yield as well as confers resistance to Karah and Merah diseases in Malaysia. Besides that, this project also aimed to characterize leaf-expressed MADS-box genes in MR84 and to examine their expression in developing leaves. Poly-A<sup>+</sup> RNA was isolated from developing leaves at different stages using oligo-dT magnetic beads, and first strand cDNA was synthesized directly on the beads. A degenerate 5' (forward) primer was designed against the conserved amino acid sequences in the MADS-box domain and an anchored dT<sub>18</sub> as the reverse primer for the three-prime (3')-RACE procedure. PCR using sscDNA as template resulted in an expected fragment size of approximately 1,013 bp. This PCR product was cloned in pGEM-T Easy plasmid vector and sequenced using forward and reverses universal M13 sequencing primers. The sequence analyses of the cloned product showed 94% similarity to other rice MADS-box cDNA. However, in order to obtain the full sequence, five-prime (5')-RACE using inner 5'-RACE as forward primer and MADS-specific as reverse primer were used to obtain the remaining 284 bp of the cDNA. A BLAST search of the complete 1,297 bp full sequence confirmed that this was the MADS-box gene and it was designated as *OsMADS\_UMS1*. The deduced amino acid sequence of *OsMADS\_UMS1* contains the typical MADS-box cDNA namely, the M, I and K boxes as well as a C terminal variable domain. Phylogenetic analyses using the PHYLIP software showed that *OsMADS\_UMS1* belongs to the class F (AGL20 or TM3) subfamily. Real-time RT-PCR analysis shows gradual mRNA steady-state levels increment during leaf development. This preliminary result indicates a possible role for *OsMADS\_UMS1* in leaf initiation and its further development. In the course of this study, another partial MADS-box cDNA was isolated designated *OsMADS\_UMS2*. This cDNA was not further studied.





## **ABSTRAK**

*Gen-gen MADS-box dicirikan oleh kehadiran domain MADS dan terdapat pada eukariot termasuk metazoa, kulat dan tumbuhan hijau. Sepanjang evolusi tumbuhan, gen-gen MADS-box berfungsi sebagai pengawal transkripsi yang aktif dalam perkembangan struktur tumbuhan. Dalam tumbuhan monokot, gen-gen MADS-box telah dikenalpasti dalam jagung, padi dan orkid. Gen-gen MADS-box mempunyai fungsi penting ketika proses perkembangan vegetatif dan reproduktif. Walaupun gen-gen MADS-box padi yang terlibat dalam perkembangan bunga dan buah telah dicirikan dengan teliti, gen-gen MADS-box padi yang diekspreskan di struktur vegetatif masih perlu diterokai. Tujuan projek ini adalah untuk mengasingkan dan menganalisis jujukan cDNA yang mengkodkan faktor transkripsi MADS-box daripada daun MR84. MR84 adalah varieti padi tempatan yang popular dan mempunyai hasil dan kualiti yang baik serta tahan kepada serangan penyakit karah dan merah. Selain itu, ia juga bertujuan untuk mencirikan gen MADS-box yang diekspreskan pada daun MR84 dan seterusnya meneliti tahap pengekspresan dalam daun yang berkembang. RNA poly-A<sup>+</sup> diasingkan daripada daun yang berkembang menggunakan manik-manik magnet oligo-dT. Lima (5') primer hadapan yang degenerat dibina daripada jujukan asid amino dalam domain MADS-box dan dT<sub>18</sub> sebagai primer belakang untuk prosedur tiga (3')-RACE. PCR daripada satu bebenang DNA komplementari menghasilkan kira-kira 1,013 pasangan bes seperti yang dijangka. Produk PCR ini diklonkan pada pGEM-T Easy vektor plasmid dan diujukkan menggunakan primer penjujukan M13 universal hadapan dan belakang. Analisis jujukan klon menunjukkan 94% persamaan dengan cDNA MADS-box padi. Untuk mendapatkan jujukan lengkap, lima (5')-RACE digunakan untuk mendapatkan baki 284 pasangan bes komplementari DNA tersebut menggunakan primer dalaman 5'-RACE sebagai primer hadapan manakala primer spesifik untuk MADS-box sebagai primer belakang. Pencarian jujukan lengkap 1,297 pasangan bes tersebut menggunakan BLAST telah mengesahkan bahawa jujukan tersebut adalah gen MADS-box dan dinamakan sebagai OsMADS\_UMS1. Jujukan asid amino OsMADS\_UMS1 mengandungi domain tipikal gen MADS-box iaitu domain MADS, I, K dan C. Keputusan filogenetik menggunakan perisian PHYLIP menunjukkan OsMADS\_UMS1 berada dalam kelas F (AGL20 atau TM3). Analisis RT-PCR secara langsung menunjukkan peningkatan tahap mRNA secara berperingkat ketika perkembangan daun. Keputusan awal ini menunjukkan kemungkinan OsMADS\_UMS1 terlibat dalam permulaan pembentukan daun dan perkembangan daun yang seterusnya. Dalam kajian ini juga, MADS-box cDNA separa lengkap telah diasingkan dan dinamakan sebagai OsMADS\_UMS2. Namun demikian, cDNA ini tidak terus dianalisis dalam kajian ini.*



## TABLE OF CONTENTS

	Page
<b>TITLE</b>	i
<b>DECLARATION</b>	iv
<b>ACKNOWLEDGEMENT</b>	vi
<b>ABSTRACT</b>	vii
<b>ABSTRAK</b>	viii
<b>LIST OF FIGURES</b>	xii
<b>LIST OF TABLES</b>	xv
<b>LIST OF APPENDICES</b>	xvi
<b>LIST OF ABBREVIATIONS</b>	xvii
<b>LIST OF SYMBOLS</b>	xix
<b>CHAPTER 1: INTRODUCTION</b>	1
<b>CHAPTER 2: LITERATURE REVIEW</b>	4
2.1 THE MADS-BOX GENE FAMILY	4
2.1.1 The Modular Structure of Plant MADS-Box Proteins	4
a. DNA Binding	4
i. DNA-Sequence Recognition	5
b. Protein-Protein Interactions	6
i. Protein Dimerization	6
ii. Ternary Complex Formation	10
c. Transcription Regulation	11
2.2 EVOLUTION OF THE PLANT MADS-BOX GENE FAMILY	12
2.2.1 Classification of Plant MADS-Box Genes	13
2.3 DIVERSE FUNCTIONS OF MADS-BOX GENES IN PLANTS	18
2.3.1 Control of Flowering Time	18
2.3.2 Control of Floral Meristem Identity	19
2.3.3 Control of Floral Organ Identity	20
a. Genetic Control of Floral Organ Identity: The ABCDE Model	20
2.3.4 Functions of MADS-Box Genes Beyond Floral Homeotic Function	24
2.4 REGULATION OF MADS-BOX GENE EXPRESSION	26
<b>CHAPTER 3: MATERIALS AND METHODS</b>	28
3.1 ISOLATION OF MADS-BOX CDNA	28
3.1.1 Plant Material	29



3.1.2	Three Prime (3')- RACE	31
a.	Poly-A <sup>+</sup> RNA purification	31
i.	Preparation of Lysate from Rice Leaves	31
ii.	Preparation of Dynabeads Oligo (dT) <sub>25</sub>	31
iii.	Direct mRNA Isolation	32
b.	Reverse Transcription	33
c.	Primer Design for Three (3')-RACE	33
d.	PCR Reaction	36
3.1.3	Five Prime (5')-RACE	37
a.	Total RNA Extraction (Salt-Precipitation Method)	37
b.	DNAse1 Treatment	38
c.	Calf Intestine Alkaline Phosphatase (CIAP) Treatment	38
d.	Tobacco Acid Pyrophosphatase (TAP) Treatment	39
e.	Five (5')-RACE Adapter Ligation	39
f.	Reverse Transcription	39
g.	Primer Design for Five (5')-RACE	40
h.	PCR Reaction	43
3.1.4	Isolation Of Full-Length Target cDNA	44
a.	Full-length cDNA Primer Design	44
b.	PCR Reaction	47
3.2	CLONING	48
3.2.1	Gel Purification of PCR Products	48
3.2.2	Ligation of PCR Products	50
3.2.3	Preparation of Competent JM109 Cells	50
a.	Method A (Hanahan Method)	50
b.	Method B (Inoue Method)	50
3.2.4	Rapid Colony Transformation of JM109	51
3.2.5	Plasmid Miniprep	51
3.2.6	Restriction Digestion	52
3.3	SEQUENCING	52
3.4	SEQUENCE ANALYSES	52
3.4.1	Blast Search and Clustalw Alignment	52
3.4.2	Exon/Intron and RNA Regulatory Motifs and Elements Prediction	53
3.5	REAL-TIME PCR	53
3.5.1	Primer Design and Specificity Confirmation	53
3.5.2	Template Preparation	57
3.5.3	Real-Time PCR Set Up	57
3.5.4	Thermal Cycling	57
3.5.5	Data Processing	57
3.6	PHYLOGENETIC TREE CONSTRUCTION	58
3.6.1	Sequence Sources and Database Searches	58
3.6.2	Sequence Alignment	60
3.6.3	Phylogenetic Trees Construction	60



<b>CHAPTER 4: RESULTS</b>	61
4.1 ISOLATION OF RICE MADS-BOX cDNA	61
4.1.1 Isolation of <i>OsMADS_UMS1</i> cDNA	62
4.1.2 Isolation of <i>OsMADS_UMS2</i> cDNA	75
4.2 ANALYSIS OF <i>OSMADS_UMS1</i> mRNA STRUCTURE	79
4.3 COMPARISON OF <i>OSMADS_UMS1</i> PROTEIN TO OTHER PLANT MADS-BOX PROTEINS	83
4.4 mRNA STEADY STATE EXPRESSION ANALYSIS OF <i>OSMADS_UMS1</i> BY REAL-TIME RT-PCR	85
4.5 PHYLOGENETIC RELATIONSHIPS	92
<b>CHAPTER 5: DISCUSSION</b>	97
5.1 ISOLATION OF RICE MADS-BOX cDNA	97
5.1.1 Analysis of the Strategy Applied to Isolate New MADS-Box Factors	97
5.1.2 <i>OSMADS_UMS1</i> and <i>OsMADS_UMS2</i> are Likely to Represent Previously Isolated Rice MADS-Box cDNA	97
5.2 TWO TYPES OF <i>OSMADS_UMS1</i> ; <i>OSMADS_UMS1a</i> AND <i>OSMADS_UMS1b</i> , DIFFERING BY cDNA SEQUENCE	99
5.2.1 <i>OSMADS_UMS1b</i> Might Be an Allele of <i>OSMADS_UMS1a</i>	99
5.2.2 <i>OSMADS_UMS1b</i> Might Be Another Different MADS-Box Gene	100
5.2.3 <i>OSMADS_UMS1b</i> Might Be an Aberrant Splice Variant Product of <i>OSMADS_UMS1a</i> Gene	100
5.3 COMPARISON OF <i>OSMADS_UMS1</i> cDNA TO OTHER PLANT GENES mRNA IN GENERAL	103
5.4 COMPARISON OF <i>OSMADS_UMS1</i> PROTEIN TO OTHER PLANT MADS-BOX PROTEINS	105
5.5 EXPRESSION AND PHYLOGENETIC ANALYSES: HINTS FOR THE FUNCTION OF MADS-BOX GENES IN DEVELOPMENT	107
5.5.1 Analysis of the Real-Time RT-PCR Procedure	107
5.5.2 Analysis of the MADS-Box Genes Phylogenetic Tree Construction Procedure	109
5.5.3 A Possible Role for <i>OSMADS_UMS1</i> in Leaf Initiation and Development	111
<b>CHAPTER 6: CONCLUSION</b>	114
<b>REFERENCES</b>	116
<b>APPENDICES</b>	133



## LIST OF FIGURES

		Page
Figure 2.1	K domain sequence alignment	9
Figure 2.2	Schematic representation of the domain structures of types I and II MADS-Box genes in plants and animals	17
Figure 2.3	Schematic representation of the ABCDE model	23
Figure 3.1	Primer sets location for 3'-RACE, 5'-RACE and full-length cDNA transcript	28
Figure 3.2	Germination of rice seeds, MR84	30
Figure 3.3	Locations of the 3' RACE primers	34
Figure 3.4	Locations of the 5' RACE primers	41
Figure 3.5	Locations of the full-length target cDNA primers	45
Figure 3.6	A diagrammatic representation of gel extraction procedure using Qiaquick Gel Extraction Kit (QIAGEN)	49
Figure 3.7	Locations of the Real-Time RT-PCR MADS-specific primers	55
Figure 4.1	Agarose gel electrophoresis with products from 3'-RACE experiment using degenerate MADS domain as forward and anchored oligo-dt <sub>17</sub> as reverse primers	62
Figure 4.2	Agarose gel electrophoresis with PCR product to confirm the presence of MADS-Box gene in the 3'-RACE experiment	63
Figure 4.3	Agarose gel electrophoresis with products from 5'-RACE experiment using nested 5'-RACE as forward primer and nested MADS-specific as reverse primer	64
Figure 4.4	Agarose gel electrophoresis with products from end-to-end PCR experiment	65
Figure 4.5	Agarose gel electrophoresis with PCR product which served to confirm the presence of MADS-box gene in the end-to-end two-step PCR experiment	66
Figure 4.6	Nucleotide and deduced amino acid sequences of <i>OsMADS_UMS1a</i> cDNA	67
Figure 4.7	Difference in <i>OsMADS_UMS1b</i> as compared to <i>OsMADS_UMS1a</i>	68
Figure 4.8	The predicted intron/exon junctions between rice genomic DNA and <i>OsMADS_UMS1</i> cDNA sequence	69
Figure 4.9	Graphical representation of the alignment	70



Figure 4.10	Alignment between the genomic DNA and <i>OsMADS_UMS1a</i> cDNA at exon 8	70
Figure 4.11	Comparison of the direct repeat sites in full-length <i>OsMADS_UMS1a</i> cDNA with the location of the direct repeat sites of <i>OsMADS_UMS1b</i> cDNA	71
Figure 4.12	Alignment of <i>OsMADS_UMS1</i> cDNA nucleotides sequence with <i>FDRMADS8</i> and <i>OsMADS56</i> cDNA sequences	72
Figure 4.13	Alignment of <i>OsMADS_UMS1</i> deduced amino acids sequence with <i>FDRMADS8</i> and <i>OsMADS56</i> protein sequences	74
Figure 4.14	<i>OsMADS_UMS2</i> PCR product using degenerate MADS domain as forward and anchored oligo-dt <sub>36</sub> as reverse primers	75
Figure 4.15	Nucleotide and deduced amino acid sequences of <i>OsMADS_UMS2</i> cDNA	76
Figure 4.16	Alignment of <i>OsMADS_UMS2</i> cDNA nucleotides sequence with <i>RAGL20</i> and <i>OsMADS50</i> cDNA sequences	77
Figure 4.17	Alignment of <i>OsMADS_UMS2</i> deduced amino acids sequence with <i>RAGL20</i> and <i>OsMADS50</i> protein sequences	79
Figure 4.18	Representative secondary structure model of <i>OsMADS_UMS1</i> mRNA predicted by Mfold	80
Figure 4.19	<i>OsMADS_UMS1</i> putative Terminal Oligopyrimidine Tract (TOP).	81
Figure 4.20	<i>OsMADS_UMS1</i> putative polyadenylation signals	81
Figure 4.21	Representative secondary structure model of <i>OsMADS_UMS1</i> 3'-UTR predicted by Mfold	82
Figure 4.22	<i>OsMADS_UMS1</i> MADS-Box protein structure and the predicted positions of NLS	83
Figure 4.23	<i>OsMADS_UMS1</i> MADS-Box protein structure and the predicted positions of potential phosphorylation site	83
Figure 4.24	Helical wheel model of <i>OsMADS_UMS1</i> K-domain	84
Figure 4.25	Alignment of the <i>OsMADS_UMS1</i> K domain amino acid with reported rice MADS-Box proteins	85
Figure 4.26	Confirmations of the products of <i>OsMADS_UMS1</i> and <i>Actin</i> primers specificity	86

Figure 4.27	Preliminary testing of the steady state expression level of <i>OsMADS_UMS1</i> by end-point RT-PCR	87
Figure 4.28	Melt curve profile in developing leaf tissues by real time RT-PCR analysis	88
Figure 4.29	Threshold cycle (Ct) profile in developing leaf tissues by real-time RT-PCR analysis	89
Figure 4.30	Graphical representation of the steady state expression levels of rice developing leaves	91
Figure 4.31	Phylogenetic tree of <i>OsMADS_UMS1</i> protein constructed by PROTPARS	93
Figure 4.32	Phylogenetic tree of <i>OsMADS_UMS1</i> protein constructed by PROTDIST and NJ	94
Figure 4.33	Multiple alignments of <i>OsMADS_UMS1</i> overall amino acid sequences to <i>OsMADS56</i> , <i>FDRMADS8</i> , <i>OsMADS50</i> and <i>AGL19</i> overall amino acid sequences	96
Figure 4.34	Multiple alignments of <i>OsMADS_UMS2</i> overall amino acid sequences to <i>AGL20</i> , <i>OsMADS50</i> , <i>OsMADS56</i> and <i>RAGL20</i> overall amino acid sequences	96

## LIST OF TABLES

		Page
Table 2.1	The MIKC-type MADS box genes in <i>Japonica</i> rice subspecies	14
Table 3.1	Descriptions of the 3' RACE primers	35
Table 3.2	Descriptions of the 5' RACE primers	42
Table 3.3	Descriptions of the full-length target cDNA primers	46
Table 3.4	Descriptions of the real-time PCR primers	56
Table 3.5	Sequences used in the phylogenetic tree construction	58
Table 4.1	The ORF, 5'-UTR and 3'-UTR lengths of <i>OsMADS_UMS1a</i> and <i>OsMADS_UMS1b</i> cDNA	68
Table 4.2	Ct values and rescaled normalized expression of emerging, expanding and mature leaf tissues relative to the normalized expression of emerging coleoptile	90
Table 4.3	Percent sequence identity of <i>OsMADS_UMS1</i> and <i>OsMADS_UMS2</i> with <i>AGL19</i> , <i>AGL20</i> , <i>OsMADS50</i> , <i>OsMADS56</i> and <i>FDRMAD8</i>	95





## LIST OF APPENDICES

	Page	
Appendix A	Reagents and Media Recipes	133
Appendix B	Common Stock Solutions	136
Appendix C	pGEM-T Easy Cloning Vector Map	139
Appendix D	Summary of RAGL20 Report	140
Appendix E	Summary of OSMADS50 Report	142
Appendix F	Summary of FDRMADS8 Report	144
Appendix G	Summary of OSMADS56 Report	146
Appendix H	Summary of AGL19 Report	148
Appendix I	Summary of AGL20 Report	150
Appendix J	Summary of OsMADS_UMS1 Report	152
Appendix K	Summary of OsMADS_UMS2 Report	154
Appendix L	Genetic Code Symbols	155
Appendix M	Methodology General Flow Chart	157
Appendix N	<i>OsMADS_UMS1</i> Sequence Chromatogram	158
Appendix O	<i>OsMADS_UMS2</i> Sequence Chromatogram	159



## LIST OF ABBREVIATIONS

aa	Amino acid
BLAST	Basic Local Alignment Search Tool
bp	Base pairs
CaCl <sub>2</sub>	Calcium chloride
cDNA	Complementary DNA
CIAP	Calf intestinal alkaline phosphatase
DNA	Deoxyribonucleic acid
DEPC	Diethylpyrocarbonate
DMSO	Dimethylsulfoxide
dNTP	Deoxyribonucleoside triphosphate
EDTA	Ethylenediaminetetra-acetate
EtBr	Ethidium bromide
EBI	European Bioinformatics Institute
g	Gram
HE	Highly expressed
H <sub>2</sub> O	Water
HCl	Hydrochloric acid
h	Hour
kb	Kilobase pairs
LE	Lowly expressed
LB	Luria-Bertani
M	Molar
mg	Miligram
MgCl <sub>2</sub>	Magnesium chloride
min	Minutes
ml	Mililitre
mM	Milimolar
mm	Milimeter
U	Unit
µg	Microgram
µl	Microlitre



μM	Micromolar
NCBI	National Center for Biotechnology Information
NaOAc	Sodium acetate
ng	Nanogram
NLS	Nuclear localization signal
nm	Nanometer
NO <sub>3</sub> <sup>-</sup>	Nitrate
nts	Nucleotides
OD	Optical density
ORF	Open reading frame
PCR	Polymerase chain reaction
RACE	Rapid Amplification of cDNA Ends
RNA	Ribonucleic acid
rpm	Revolutions per minute
RT	Reverse transcriptase
sdH <sub>2</sub> O	Sterile distilled water
SAM	Shoot apical meristem
SD	Standard deviation
SDS	Sodium dodecyl sulphate
sec	Seconds
TAP	Tobacco acid pyrophosphatase
TAE	Tris acetate EDTA
TBE	Tris borate EDTA
TE	Tris-EDTA
TSS	Transcription start site
UTR	Untranslated region
UV	Ultraviolet
V	Voltage



## LIST OF SYMBOLS

$\sim$	Approximately
$/$	<i>Per</i>
$<$	Less than
$>$	More than
$\leq$	Less than or equal to
$'$	Prime
$\%$	Percent
$\alpha$	Alpha
$\beta$	Beta
$\delta$	Delta
$^{\circ}\text{C}$	Degree celcius



## CHAPTER ONE

### INTRODUCTION

MADS-box transcription factors are named after the initials of the four originally identified members. The yeast MINICHROMOSOME MAINTAINANCE 1 (MCM1) protein regulates mating-type-specific gene expression, the *Arabidopsis* AGAMOUS (AG) and *Antirrhinum* DEFICIENS (DEF) proteins play regulatory roles in specifying the identity of floral organs, and the human serum response factor (SRF) is involved in the transcriptional regulation of the protooncogene, *c-fos*. After the isolation of the first plant MADS-box genes, *DEF* and *AG* from plants (Sommer *et al.*, 1990; Yanofsky *et al.*, 1990), a large number of MADS-box genes have been sequentially identified in different species of angiosperms, and the orthologues of floral homeotic MADS-box genes have also been found in gymnosperm (Becker *et al.*, 2000; Munster *et al.*, 1997). MIKC type MADS-box genes were also cloned from a number of species outside the seed plant lineage including ferns, mosses, lycophids and from the closest relatives of land plants, the fresh water green algae charophyceans (Tanabe *et al.*, 2005; Svensson and Engstrom, 2002; Henschel *et al.*, 2002; Hasebe *et al.*, 1998; Munster *et al.*, 1997).

MADS-box genes are characterized by the conserved MADS domain and found in a wide range of eukaryotes including metazoans, fungi, slime mold, and green plants. The gene family can be divided into two main lineages, referred to as type I and type II, both of which are present in plants, animals and fungi (Alvarez-Buylla *et al.*, 2000a). All members of the family possess the 180-nucleotide long (on average) MADS box. In plants, type II MADS-domain proteins, referred to as MIKC-type proteins, possess three additional functional domains: a well-conserved K (keratin) domain, a less-well-conserved I (intervening) domain, and a variable C-terminal region.



In contrast to type II genes, which have been the subject of extensive research, not much is known about the type I genes in plants. Except for the MADS box, the type I genes share no sequence similarity with type II genes. However, some type I genes share conserved C-terminal motifs with each other (De Bodt *et al.*, 2003b; Parenicova *et al.*, 2003). In addition, a third group of genes has recently been identified that possesses only half of the MADS box or are overall highly divergent. These are referred to as MADS-like genes (De Bodt *et al.*, 2003b).

Throughout plant evolution, MADS-box genes have been recruited as transcriptional regulators active in the development of diverse plant structures. Since the discovery of the first MADS-box genes more than a decade ago, biologists have made great progress in elucidating the roles of these genes in plant development. Expression studies and mutant analyses on MADS-box genes in diverse plant species revealed the crucial importance of MADS-box genes in the regulation of both reproductive (flower, seed, fruit) (Vrebalov *et al.*, 2002; Nesi *et al.*, 2002; Liljegren *et al.*, 2000; Fernandez *et al.*, 2000; Mao, 2000; Western and Haughn, 1999; Gu *et al.*, 1998) and vegetative (root, leaf) development (Rosin *et al.*, 2003; Prakash and Kumar, 2000; Garcia-Maroto *et al.*, 2000; Heuer *et al.*, 2001, 2000; Zhang and Forde, 1998). Furthermore, MADS-box genes used in the control of floral patterning form the ideal genetic toolkit to study the diversification of flower architecture (Pinyopich *et al.*, 2003; Goto *et al.*, 2001; Theissen, 2001; Theissen and Saedler, 2001; Pelaz *et al.*, 2001, 2000; Alvarez and Smyth, 1999; Colombo *et al.*, 1995; Coen and Meyerowitz, 1991).

In monocots, MADS-box genes have been identified in maize, rice, and orchids (Lee *et al.*, 2003a; Pelucchi *et al.*, 2002; Heuer *et al.*, 2001; Yu and Goh, 2000). The fact that MADS box genes also function in monocotyledons is of great interest because of the agronomic importance of the species that belong to this group, not least among which are the cereals. MADS-box genes may thus provide a tool for elucidating the complex genetic system controlling development in cereals.

However, the structural and functional characterization of new MADS-box genes in cereals is required to further our understanding of this complex family of regulatory genes and their involvement in controlling the developmental processes in

cereals. Rice was chosen because of its importance as a worldwide crop and, additionally, because of its advantages as a biological model system for cereals in general, its genetics is well developed, extremely detailed genetic linkage maps are available and it can be efficiently transformed (Greco *et al.*, 1997).

MADS-box gene family in rice has resulted in the discovery of non-redundant 35 MADS-box genes (Yamaguchi *et al.*, 2006; Lee *et al.*, 2003a). The MIKC-type MADS box genes in *Japonica* rice subspecies is shown in Table 2.1. Although rice MADS-box genes involved in flower and fruit development have been well characterized, rice MADS-box genes expressed in vegetative structures have yet to be explored. Much work is still needed to uncover the molecular mechanisms of vegetative development.

Therefore, the aim of this project is to isolate and analyze the cDNA sequences encoding for putative MADS-box transcription factors from MR84 leaves. MR84 is currently the popular local cultivar which produced good yield and quality and confers resistance to Karah and Merah diseases as well. Besides that, it also aimed to characterize leaf-expressed MADS-box genes and to examine their expression in developing leaves.

Objectives of the study:

- a. To isolate cDNA of MADS-box gene in rice leaves
- b. To characterize the cDNA sequence using bioinformatic applications (BLAST search, CLUSTALW multiple alignment, exon/intron and RNA motifs/elements prediction and phylogenetic analysis)
- c. To investigate the expression level of the MADS-box gene during rice leaves development

## CHAPTER TWO

### LITERATURE REVIEW

#### 2.1 THE MADS-BOX GENE FAMILY

The MADS box is a highly conserved sequence motif found in a family of transcription factors. The conserved domain was named after the first four members of the family, which were MCM1, AGAMOUS, DEFICIENS and SRF. The name MADS was constructed from the "initials" of these four "founders".

##### 2.1.1 The Modular Structure of Plant MADS-Box Proteins

Typical plant transcription factors consist of a DNA-binding region, an oligomerization site, a transcription regulation domain and a nuclear localization signal although some lack either a transcription regulation domain or a specific DNA-binding domain (Liu *et al.*, 1999). By mutational and functional analysis it has been demonstrated that MADS-box proteins consist of a DNA-binding region, a region which serves as an interface for dimerization and interactions with other proteins, and sometimes contain a transcriptional activation domain. There is a considerable overlap between these functional domains and the M, I, K and C structural domains, although none of the functions can exclusively be assigned to just one single domain. Such modular organization is common to many eukaryotic transcription factors.

##### a. DNA Binding

DNA binding by plant type II MADS-box proteins to *cis*-acting promoter elements, named the CARG-box, is mediated by the MADS-domain and the K-box is believed to contribute to the binding specificity. Studies to identify the minimal DNA-binding domain of the *Antirrhinum* MADS-box proteins SQUAMOSA (SQUA) and PLENA (PLE) demonstrated that the MADS- and I-domains are sufficient to permit sequence-specific DNA binding by the proteins (West *et al.*, 1998). Similar results were





## REFERENCES

- Acton, T. B., Mead, J., Steiner, A. M. and Vershon, A. K. 2000. Scanning mutagenesis of MCM1: residues required for DNA binding, DNA bending and transcriptional activation by a MADS-box protein. *Mol. Cell. Biol.* **20**: 1–11.
- Aeschbacher, R.A., Schrott, M., Potrykus, I. and Saul, M.W. 1991. Isolation and molecular characterization of *PosF21*, an *Arabidopsis thaliana* gene which shows characteristics of a b-ZIP class transcription factor. *Plant J.* **1**: 303-316.
- Alvarez, J. and Smyth, D.R. 1999. *CRABS CLAW* and *SPATULA*, two *Arabidopsis* genes that control carpel development in parallel with *AGAMOUS*. *Development.* **126**: 2377-2386.
- Alvarez-Buylla, E.R., Liljegren, S.J., Gold, S.E., Burgeff, C., Ditta, G.S., de Pouplana, L.R., Martinez-Castilla, L. and Yanofsky, M.F. 2000a. An ancestral MADS-box gene duplication occurred before the divergence of plants and animals. *Proc. Natl. Acad. Sci. USA.* **97**: 5328-5333.
- Alvarez-Buylla, E. R., Liljegren, S. J., Pelaz, S., Gold, S. E., Burgeff, C., Ditta, G. S., Vergara-Silva, F., and Yanofsky, M. F. 2000b. MADS-box gene evolution beyond flowers: expression in pollen, endosperm, guard cells, roots and trichomes. *Plant J.* **24**: 1-11.
- Ambrose, B.A., Lerner, D.R., Ciceri, P., Padilla, C.M., Yanofsky, M.F., Schmidt, R.J. 2000. Molecular and genetic analyses of the *silky1* gene reveal conservation in floral organ specification between eudicots and monocots. *Mol. Cell.* **5**: 569-579.
- Ando, S., Sarlis, N.J., Krishnan, J., Feng, X., Refetoff, S., Zhang, M.Q., Oldfield, E.H. and Yen, P.M. 2001. Aberrant alternative splicing of thyroid hormone receptor in a TSH-secreting pituitary tumor is a mechanism for hormone resistance. *Mol. Endocrinol.* **15**: 1529–1538.
- Angenent, G.C., Busscher, J., Franken J., Mol, J.N.M. and van Tunen, A.J. 1992. Differential expression of two MADS-box genes in wild-type and mutant petunia flowers. *Plant Cell.* **4**: 983-993.
- Angenent, G.C., Franken, J., Busscher, M., Colombo, L. and van Tunen, A.J. 1993. Petal and stamen formation in petunia is regulated by the homeotic gene *fbp1*. *Plant J.* **4**: 101-112.
- Angenent, G.C., Franken, J., Busscher, M., van Dijken, A., van Went, J.L., Dons, H.J.M. and van Tunen, A.J. 1995. A novel class of MADS box genes is involved in ovule development in petunia. *Plant Cell.* **7**: 1569-1582.



- Angenent, G.C., Franken, J., Busscher, M., Weiss, D. and van Tunen, A.J. 1994. Co-suppression of the petunia homeotic gene *fbp2* affects the identity of the generative meristem. *Plant J.* **5**: 33-44.
- Baralle, D. and Baralle, M. 2005. Splicing in action: assessing disease causing sequence changes. *J. Med. Genet.* **42**: 737-748.
- Becker A., Winter, K. U., Meyer B., Saedler H. and Theissen, G. 2000. MADS-box gene diversity in seed plants 300 million years ago. *Mol. Biol. Evol.* **17**: 1425-1434.
- Berbel, A., Navarro, C., Ferrándiz, C., Cañas, L.A., Madueño, F. and Beltrán, J. 2001. Analysis of *PEAM4*, the pea *AP1* functional homologue, supports a model for *AP1*-like genes controlling both floral meristem and floral organ identity in different plant species. *Plant J.* **25**: 441-451.
- Blazquez, M.A., Green, R., Nilsson, O., Sussman, M.R. and Weigel, D. 1998. Gibberellins promote flowering of *Arabidopsis* by activating the *LEAFY* promoter. *Plant Cell.* **10**: 791-800.
- Bowman, J. L., Alvarez, J., Weigel, D., Meyerowitz, E. M. and Smyth, D. R. 1993. Control of flower development in *Arabidopsis thaliana* by *APETALA1* and interacting genes. *Development.* **119**: 721-743.
- Bowman, J. L., Sakai, H., Jack, T., Weigel, D., Mayer, U. and Meyerowitz, E. M. 1992. *SUPERMAN*, a regulator of floral homeotic genes in *Arabidopsis*. *Development.* **114**: 599-615.
- Buratti, E. and Baralle, F.E. 2004. Influence of RNA secondary structure on the pre-mRNA splicing process. *Mol. Cell Biol.* **24**:10505-10514.
- Bustin, S. A., Benes, V., Nolan, T. and Pfaffl, M. W. 2005. Quantitative real-time RT-PCR- a perspective. *J. Mol. Endocrinol.* **34**: 597-601.
- Carlsbecker, A., Tandre, K., Johanson, U., Englund, M. and Engstrom, P. 2004. The MADS-box gene *DAL1* is a potential mediator of the juvenile-to-adult transition in Norway spruce *Picea abies*. *Plant J.* **40**: 546-557.
- Carmona, M.J., Ortega, N. and Garcia-Maroto, F. 1998. Isolation and molecular characterization of a new vegetative MADS-box gene from *Solanum tuberosum* L. *Planta.* **207**: 181-188.
- Chen, X. 2004. A microRNA as a translational repressor of *APETALA2* in *Arabidopsis* flower development. *Science.* **303**: 2022-2025.
- Chen, Z.X., Wu, J.G., Ding, W.N., Chen, H.M., Wu, P. and Shi, C.H. 2006. Morphogenesis and molecular basis on naked seed rice, a novel homeotic mutation of OsMADS1 regulating transcript level of AP3 homologue in rice. *Planta.* **223**: 882-890.

- Cheng, X.F., Wittich, P.E., Kieft, H., Angenent, G., Han, X.X. and van Lammeren, A.A.M. 2000. Temporal and spatial expression of MADS box genes, *FBP7* and *FBP11*, during initiation and early development of ovules in wild type and mutant *Petunia hybrida*. *Plant Biol.* **2**: 693–702.
- Cho, S., Jang, S., Chae, S., Chung, K.M., Moon, Y.H., An, G. and Jang, S.K. 1999. Analysis of the C-terminal region of *Arabidopsis thaliana* APETALA1 as a transcription activation domain. *Plant Mol. Biol.* **40**: 419–429.
- Coen E.S. and Meyerowitz, E. M. 1991. The war of the whorls: genetic interactions controlling flower development. *Nature.* **353**: 31-37.
- Colombo, L., Franken, J., Koetje, E., van Went, J., Dons, H.J.M., Angenent, G.C. and van Tunen, A.J. 1995. The petunia MADS box gene *FBP11* determines ovule identity. *Plant Cell.* **7**: 1859-1868.
- Conner, J. and Liu, Z. 2000. LEUNIG, a putative transcriptional corepressor that regulates AGAMOUS expression during flower development. *Proc. Natl. Acad. Sci. USA.* **97**: 12902–12907.
- Courey, A.J., Holtzman, D.A., Jackson, S.P. and Tjian, R. 1989. Synergistic activation by the glutamine-rich domains of human transcription factor Sp1. *Cell.* **59**: 827-836.
- Curry, J., McHale, C. and Smith, M.T. 2002. Low efficiency of the Moloney murine leukemia virus reverse transcriptase during reverse transcription of rare t8;21. fusion gene transcripts. *Biotechniques.* **32**: 768–770.
- Davies, B. & Schwarz-Sommer, Z. 1994. Control of Floral Organ Identity by Homeotic MADS-Box Transcription Factors. In Nover, L. ed... *Plant Promoters and Transcription Factors, Results and Problems in Cell Differentiation No. 20*, pp. 235–258. Berlin, Heidelberg:Springer.
- Davies, B., Egea-Cortines, M., de Andrade Silva, E., Saedler, H. and Sommer, H. 1996. Multiple interactions amongst floral homeotic MADS box proteins. *EMBO J.* **16**: 4330–4343.
- Davies, B., Motte, P., Keck, E., Saedler, H., Sommer, H. and Schwarz-Sommer, Z. 1999. PLENA and FARINELLI: redundancy and regulatory interactions between two *Antirrhinum* MADS-box factors controlling flower development. *EMBO J.* **18**: 4023-4034.
- Davis, L., Kuehl, M. & Battey, J. 1994. *Basic methods in molecular biology.* 2<sup>nd</sup> edition.. Connecticut: Appleton and Lange.
- De Bodt, S., Raes, J., de Peer, Y.V. and Theissen, G. 2003a. And then there were many: MADS goes genomic. *Trends Plant Sci.* **8**: 475-483.

- De Bodt, S., Raes, J., Florquin, K., Rombauts, S., Rouze, P., Theissen, G. and de Peer, Y. V. 2003b. Genomewide structural annotation and evolutionary analysis of the type I MADS-box genes in plants. *J. Mol. Evol.* **56**: 573-586.
- Decroocq, V., Zhu, X., Kauffman, M., Kyojuka, J., Peacock, W.J., Dennis, E.S. and Lewellyn, D.J. 1999. A *TM3*-like MADS-box gene from *Eucalyptus* expressed in both vegetative and reproductive tissues. *Gene.* **228**: 155-160.
- Ditta, G., Pinyopich, A., Robles, P., Pelaz, S. and Yanofsky, M.F. 2004. The *SEP4* gene of *Arabidopsis thaliana* functions in floral organ and meristem identity. *Curr. Biol.* **14**: 1935-1940.
- Egea-Cortines, M., Saedler, H. and Sommer, H. 1999. Ternary complex formation between the MADS-box proteins SQUAMOSA, DEFICIENS, and GLOBOSA is involved in the control of floral architecture in *Antirrhinum majus*. *EMBO J.* **18**: 5370-5379.
- Eisen, J.A. 1998. Phylogenomics: Improving functional predictions for uncharacterized genes by evolutionary analysis. *Genome Res.* **8**: 163-167.
- Fan, H.Y., Hu, Y., Tudor, M. and Ma, H. 1997. Specific interactions between the K domains of AG and AGLs, members of the MADS domain family of DNA binding proteins. *Plant J.* **12**: 999-1010.
- Favaro, R., Pinyopich, A., Battaglia, R., Kooiker, M., Borghi, L., Ditta, G., Yanofsky, M.F., Kater, M.M. and Colombo, L. 2003. MADS box protein complexes control carpel and ovule development in *Arabidopsis*. *Plant Cell.* **15**: 2603-2611.
- Felsenstein, J. 1996. Inferring phylogenies from protein sequences by parsimony, distance, and likelihood methods. *Meth. in Enzym.* **266**: 419-427.
- Fernandez, D.E., Heck, G. R., Perry, S.E., Patterson, S.E., Bleecker, A.B. and Fang, S. C. 2000. The embryo MADS domain factor AGL15 acts postembryonically: Inhibition of perianth senescence and abscission via constitutive expression. *Plant Cell.* **12**: 183-198.
- Ferrández, C., Liljegren, S. and Yanofsky, M. 2000. Negative regulation of the *SHATTERPROOF* genes by FRUITFULL during *Arabidopsis* fruit development. *Science.* **289**: 436-438.
- Ferrández, C., Pelaz, S., Yanofsky, M.F. 1999. Control of carpel and fruit development in *Arabidopsis*. *Ann. Rev. Biochem.* **68**: 321-354.
- Ferrario, S., Immink, R. G. H., Shchennikova, A., Busscher-Lange, J. and Angenent, G. C. 2003. The MADS box gene *FBP2* is required for SEPALLATA function in petunia. *Plant Cell.* **15**: 914-925.
- Fleming, A.J., Mandel, T., Roth, I. and Kuhlemeier, C. 1993. The patterns of gene expression in the tomato shoot apical meristem. *Plant Cell.* **5**: 297-309.

- Fornara, F., Parenicova, L., Falasca, G., Pelucchi, N., Masiero, S., Ciannamea, S., Lopez-Dee, Z., Altamura, M.M., Colombo, L. and Kater, M.M. 2004. Functional characterization of *OsMADS18*, a member of the *AP1/SQUA* subfamily of MADS box genes. *Plant Physiol.* **135**: 2207-2219
- Garcia-Maroto, F., Ortega, N., Lozano, R. and Carmona, M.J. 2000. Characterization of the potato MADS-box gene *STMADS16* and expression analysis in tobacco transgenic plants. *Plant Mol. Biol.* **42**: 499–513.
- Garvie, C. W. and Wolberger, C. 2001. Recognition of specific DNA sequences. *Mol. Cell.* **8**: 937-946.
- Goto, K., Kyojuka, J. and Bowman, J. L. 2001. Turning floral organs into leaves, leaves into floral organs. *Curr. Opin. Genet. Dev.* **11**: 449 -456.
- Greco, R., Stagi, L., Colombo, L., Angenent, G.C., Sari-Gorla, M., Pe, M.E. 1997. MADS box genes expressed in developing inflorescences of rice and sorghum. *Mol. Gen. Genet.* **253**: 615-623.
- Gu, Q., Ferrandiz, C., Yanofsky, M. and Martienssen, R. 1998. The FRUITFULL MADS-box gene mediates cell differentiation during *Arabidopsis* fruit development. *Development.* **125**: 1509–1517.
- Gustafson-Brown, C., Savidge, B. and Yanofsky, M.F. 1994. Regulation of the *Arabidopsis* floral homeotic gene *APETALA1*. *Cell.* **76**: 131-143.
- Gutierrez-Cortines, M.E. and Davies, B. 2000. Beyond the ABCs: ternary complex formation in the control of floral organ identity. *Trends Plant Sci.* **5**: 471-476.
- Hasebe, M., Wen, C.K., Kato, M. and Banks, J.A. 1998. Characterization of MADS homeotic genes in the fern *Ceratopteris richardii*. *Proc. Natl. Acad. Sci. USA.* **95**: 6222-6227.
- Hassler, M. and Richmond, T.J. 2001. The B-box dominates SAP-1–SRF interactions in the structure of the ternary complex. *EMBO J.* **20**: 3018–3028.
- Heard, J. and Dunn, K. 1995. Symbiotic induction of a MADS-box gene during development of alfalfa root nodules. *Proc. Natl. Acad. Sci. USA.* **92**: 5273-5277.
- Henschel, K., Kofuji, R., Hasebe, M., Saedler, H., Munster, T. and Theissen, G. 2002. Two ancient classes of MIKC-type MADS-box genes are present in the moss *Physcomitrella patens*. *Mol.Biol.Evol.* **19**: 801-814.
- Heuer, S., Hansen, S., Bantini, J., Brettschneider, R., Kranz, E., Lorz, H. and Dresselhaus, T. 2001. The Maize MADS box gene *ZmMADS3* affects node number and spikelet development and is co-expressed with *ZmMADS1* during flower development, in egg cells and early embryogenesis. *Plant Physiol.* **127**: 33-45.

- Heuer, S., Lorz, H. and Dresselhaus, T. 2000. The MADS box gene *ZmMADS2* is specifically expressed in maize pollen and during maize pollen tube growth. *Sex Plant Reprod.* **13**: 21-27.
- Hillis, D.M., Allard, M.W., and Miyamoto, M.M. 1993. Analysis of DNA sequence data: phylogenetic inference. *Meth. in Enzym.* **224**: 456-487.
- Honma, T. and Goto, K. 2001. Complexes of MADS-box proteins are sufficient to convert leaves into floral organs. *Nature.* **409**: 525–529.
- Hope, I.A. and Struhl, K. 1986. Functional dissection of an eukaryotic transcriptional activator protein, GCN4 of yeast. *Cell.* **46**: 885-894.
- Hoshikawa, K. 1989. *The growing rice plant. An anatomical monograph.* Tokyo, Japan: Nosan Gyoson Bunka, Kyokai Nobunkyo..
- Huang, H., Tudor, M., Su, T., Zhang, Y., Hu, Y. and Ma, H. 1996. DNA binding properties of two Arabidopsis MADS domain proteins: binding consensus and dimer formation. *Plant Cell.* **8**: 81-94.
- Huijser, P., Klien, J., Lönnig, W.E., Meijer, H., Saedler, H. and Sommer, H. 1992. Bracteomania, an inflorescence anomaly, is caused by the loss of function of the MADS-box gene *squamosa* in *Antirrhinum majus*. *EMBO J.* **11**: 1239-1249.
- Immink, R.G.H., Ferrario, S., Busscher-Lange, J., Kooiker, M., Busscher, M. and Angenent, G.C. 2003. Analysis of the petunia MADS-box transcription factor family. *Mol. Genet. Genomics.* **268**: 598–606.
- Immink, R.G.H., Gadella, T.W., Jr., Ferrario, S., Busscher, M. and Angenent, G.C. 2002. Analysis of MADS box protein-protein interactions in living plant cells. *Proc. Natl. Acad. Sci. USA.* **99**: 2416–2421.
- Immink, R.G.H., Hannapel, D. J., Ferrario, S., Busscher, M., Franken, J., Campagne, M.M.L. and Angenent, G.C. 1999. A petunia MADS box gene involved in the transition from vegetative to reproductive development. *Development.* **126**: 5117-5126.
- Irish, V.F. and Litt, A. 2005. Flower development and evolution: gene duplication, diversification and redeployment. *Curr. Opin. Genet. Dev.* **15**: 454-460.
- Jack, T., Fox, G.L. and Meyerowitz, E.M. 1994. Arabidopsis homeotic gene *APETALA3* ectopic expression: Transcriptional and posttranscriptional regulations determine floral organ identity. *Cell.* **76**: 703–716.
- Jeon, J.S., Jang, S., Lee, S., Nam, J., Kim, C., Lee, S.H., Chung, Y.Y., Kim, S.R., Lee, Y.H. and Cho, Y.G. 2000. *leafy hull sterile1* is a homeotic mutation in a rice MADS box gene affecting rice flower development. *Plant Cell.* **12**: 871–84.

- Joshi C.P. 1987. An inspection of the domain between putative TATA box and translation start site in 79 plant genes. *Nucleic Acids Res.* **15**: 6643-6652.
- Jelesko, J.G, Carter, K., Thomson, W., Kinoshita, Y. and Gruissem, W. 2004. Meiotic recombination between paralogous RBCSB genes on sister chromatids of *Arabidopsis thaliana*. *Genetics.* **166**: 947-957.
- Kang, H.G. and An, G.H. 1997a. Isolation and characterization of a rice MADS box gene belonging to the *AGL2* gene family. *Mol. Cells.* **7**: 45-51.
- Kang, H.G., Jang, S., Chung, J.E., Cho, Y.G. and An, G. 1997b. Characterization of two rice MADS box genes that control flowering time. *Mol. Cell.* **7**: 559-566.
- Kang, H.G., Jeon, J.S., Lee, S. and An, G. 1998. Identification of class B and class C floral organ identity genes from rice plants. *Plant Mol. Biol.* **38**: 1021-1029.
- Kang, H.G., Noh, Y.S., Chung, Y.Y., Costa, M.A., An, K. and An, G. 1995. Phenotypic alterations of petal and sepal by ectopic expression of a rice MADS box gene in tobacco. *Plant Mol. Biol.* **29**: 1-10.
- Kochetov, A.V. 2005. AUG codons at the beginning of protein coding sequences are frequent in eukaryotic mRNAs with a suboptimal start codon context. *Bioinformatics.* **21**: 837-840.
- Kochetov, A.V., Ischenko, I.V., Vorobiev, D.G., Kel, A.E., Babenko, V.N., Kisselev, L.L. and Kolchanov, N.A. 1998a. Eukaryotic mRNAs encoding abundant and scarce proteins are statistically dissimilar in many structural features. *FEBS Lett.* **440**: 351-355.
- Kochetov, A.V., Pilugin, M.V., Kolpakov, F.V., Babenko, V.N., Kolchanov, N.A. and Shumny, V.K. 1998b. Sequence analysis of 5'untranslated regions of plant mRNAs. *Adv. Curr. Biol.* **118**: 754-770.
- Kofuji, R., Sumikawa, N., Yamasaki, M., Kondo, K., Ueda, K., Ito, M. and Hasebe, M. 2003. Evolution and divergence of MADS-box gene family based on genome wide expression analyses. *Mol. Biol. Evol.* **20**: 1963-1977.
- Kohler, C., Hennig, L., Spillane, C., Pien, S., Gruissem, W. and Grossniklaus, U. 2003. The *polycomb*-group protein MEDEA regulates seed development by controlling expression of the MADS-box gene *PHERES1*. *Genes Dev.* **17**: 1540-1553.
- Kozak, M. 1989. The scanning model for translation: an update. *J. Cell Biol.* **108**: 229-241.
- Kozak, M. 2002. Pushing the limits of the scanning mechanism for initiation of translation. *Gene.* **299**: 1-34.

- Krizek, B.A. and Meyerowitz, E.M. 1996. Mapping the protein regions responsible for the functional specificities of the *Arabidopsis* MADS domain organ-identity proteins. *Proc. Natl. Acad. Sci. USA*. **93**: 4063-4070.
- Kyozuka, J., Kobayashi, T., Morita, M. and Shimamoto, K. 2000. Spatially and temporally regulated expression of rice MADS box genes with similarity to *Arabidopsis* class A, B and C genes. *Plant Cell Physiol*. **41**: 710–718.
- Kyozuka, J., Konishi, S., Nemoto, K., Izawa, T. and Shimamoto, K. 1998. Down-regulation of *RFL* the *FLO/IFY* homolog of rice accompanied with panicle branch initiation. *Proc. Natl. Acad. Sci. USA*. **95**: 1979–1982.
- Lee, H., Suh, S.S., Park, E., Cho, E., Ahn, J.H., Kim, S.G., Lee, J.S., Kwon, Y.M. and Lee, I. 2000. The AGAMOUS-LIKE 20 MADS domain protein integrates floral inductive pathways in *Arabidopsis*. *Genes & Development*. **14**: 2366–2376.
- Lee, S., Kim, J., Han, J.J., Han, M.J. and An, G. 2004. Functional analyses of the flowering time gene *OsMADS50*, the putative *SUPPRESSOR OF OVEREXPRESSION OF CO 1/AGAMOUS-LIKE 20 SOC1/AGL20*. ortholog in rice. *Plant J*. **38**: 754–764.
- Lee, S., Kim, J., Son, J.S., Nam, J., Jeong, D.H., Lee, K., Jang, S., Yoo, J., Lee, J., Lee, D.Y., Kang, H.G. and An, G. 2003a. Systematic reverse genetic screening of T-DNA tagged genes in rice for functional genomic analysis: MADS-box genes as a test case. *Plant Cell Physiol*. **44**: 1403-1411.
- Lee, S., Jeon, J.S., An, K., Moon, Y.H., Lee, S., Chung, Y.Y. and An, G. 2003b. Alteration of floral organ identity in rice through ectopic expression of *OsMADS16*. *Planta*. **217**: 904-11.
- Levy, S., Avni, D., Hariharan, N., Perry, R.P. and Meyuhas O. 1991. Oligopyrimidine tract at the 5' end of mammalian ribosomal protein mRNAs is required for their translational control. *Proc. Natl. Acad. Sci. USA*. **88**: 3319-23.
- Liljegren, S.J., Ditta, G.S., Eshed, Y., Savidge, B., Bowman, J.L. and Yanofsky, M.F. 2000. SHATTERPROOF MADS-box genes control seed dispersal in *Arabidopsis*. *Nature*. **404**: 766-770.
- Lim, F.L, Hayes, A., West, A.G., Pic-Taylor, A., Darieva, Z., Morgan, B.A., Oliver, S.G. and Sharrocks, A.D. 2003. Mcm1p-induced DNA bending regulates the formation ternary transcription factor complexes. *Mol. Cell. Biol*. **23**: 450-461.
- Lim, J., Moon, Y.H., An, G. and Jang, S.K. 2000. Two rice MADS domain proteins interact with *OsMADS1*. *Plant Mol. Biol*. **44**: 513-527.
- Liu, L, White, M.J. and MacRae T.H. 1999. Transcription factors and their genes in higher plants. *Eur. J. Biochem*. **262**: 247-257.
- Liu, W., Xu, Z.H., Luo, D. and Xue, H.W. 2003. Roles of *Oskn1*, a rice casein kinase 1, in root development and plant hormone sensitivity. *Plant J*. **36**: 189-202.



- Liu, Z. and Meyerowitz, E.M. 1995. *LEUNIG* regulates *AGAMOUS* expression in *Arabidopsis* flowers. *Development*. **121**: 975–991.
- Loke, J.C., Stahlberg, E.A., Strenski, D.G., Haas, B.J., Wood, P.C. and Li, Q.Q. 2005. Compilation of mRNA polyadenylation signals in *Arabidopsis* revealed a new signal element and potential secondary structures. *Plant Physiol*. **138**: 1457–1468.
- Lopez-Dee, Z. P., Wittich, P., Pe, M.E., Rigola, D. and Del Buono I. 1999. *OsmADS13*, a novel rice MADS-box gene expressed during ovule development. *Dev. Genet*. **25**: 237-244.
- Lukas, J., Gao, D.Q., Keshmeshian, M., Wen, W.H., Wei, D.T., Rosenberg, D. and Press, M.F. 2001. Alternative and aberrant messenger RNA splicing of the *mdm2* oncogene in invasive breast cancer. *Cancer Res*. **61**: 3212–3219.
- Ma, H., Yanofsky, M. F. and Meyerowitz, E. M. 1991. AGL1-AGL6, an *Arabidopsis* gene family with similarity to floral homeotic and transcription factor genes. *Genes Dev*. **5**: 484–495.
- Ma, J. and Ptashne, M. 1987. A new class of yeast transcriptional activators. *Cell*. **51**:113-119.
- Mandel, M.A. and Yanofsky, M. F. 1995. The *Arabidopsis* AGL8 MADS box gene is expressed in inflorescence meristems and is negatively regulated by APETALA1. *Plant Cell*. **7**: 1763-1771.
- Mandel, M.A., Lutziger, I. and Kuhlemeier, C. 1994. A ubiquitously expressed MADS-box gene from *Nicotiana tabacum*. *Plant Mol. Biol*. **25**: 319-321.
- Mao, L., Begum, D., Chuang, H. W., Budiman, M. A., Szymkowwiak, E. J., Irish, E. E. and Wing, R. A. 2000. *JOINTLESS* is a MADS-box gene controlling tomato flower abscission zone development. *Nature*. **406**: 910-913.
- May, A.C. 2004. Percent sequence identity; the need to be explicit. *Structure Camb*. **12**: 737-738.
- McGonigle, B., Bouhidel, K. and Irish, V.F. 1996. Nuclear localization of the *Arabidopsis* APETALA3 and PISTILLATA homeotic gene products depends on their simultaneous expression. *Genes Dev*. **10**: 1812–1821.
- Mena, M., Ambrose, B.A., Meeley, R.B., Briggs, S.P., Yanofsky, M.F. and Schmidt, R.J. 1996. Diversification of C-function activity in maize flower development. *Science*. **274**: 1537–1540.
- Mermod, N., O'Neill, E.A., Kelly, T.J. and Tjian, R. 1989. The proline-rich transcriptional activator of CTF/NF-1 is distinct from the replication and DNA binding domain. *Cell*. **58**: 741-753.

- Meyer, I.M. and Miklos, I. 2005. Statistical evidence for conserved, local secondary structure in the coding regions of eukaryotic mRNAs and pre-mRNAs. *Nucleic Acids Res.* **33**: 6338-6348.
- Meyuhas, O. 2000. Synthesis of the translational apparatus is regulated at the translational level. *Eur. J. Biochem.* **267**: 6321-30.
- Michaels, S. D. and Amasino, R. M. 1999. *FLOWERING LOCUS C* encodes a novel MADS domain protein that acts as a repressor of flowering. *Plant Cell.* **11**: 949-956.
- Michaels, S.D. and Amasino, R.M. 2001. Loss of *FLOWERING LOCUS C* activity eliminates the late-flowering phenotype of *FRIGIDA* and autonomous pathway mutations but not responsiveness to vernalization. *Plant Cell.* **13**: 935-941.
- Mizukami, Y., Huang, H., Tudor, M., Hu, Y. and Ma, H. 1996. Functional domains of the floral regulator *AGAMOUS*: characterization of the DNA binding domain and analysis of dominant negative mutations. *Plant Cell.* **8**: 831-845.
- Montag, K., Salamini, F. and Thompson, R.D. 1995. *ZEMa*, a member of a novel group of MADS box genes, is alternatively spliced in maize endosperm. *Nucleic Acids Res.* **23**: 2168-2177.
- Moon, Y. H., Chen, L., Pan, R.L., Chang, H.S., Zhu, T., Maffeo, D. M. and Sung, Z. R. 2003. *EMF* genes maintain vegetative development by repressing the flower program in *Arabidopsis*. *Plant Cell.* **15**: 681-693.
- Moon, Y.H., Jung, J.Y., Kang, H.G. and An, G. 1999a. Identification of a rice *APETALA3* homolog by yeast two-hybrid screening. *Plant Mol. Biol.* **40**: 167-177.
- Moon, Y.H., Kang, H.G., Jung, J.Y., Jeon, J.S., Sung, S.K. and An, G. 1999b. Determination of the motif responsible for interaction between the rice *APETALA1/AGAMOUS-LIKE9* family proteins using a yeast two-hybrid system. *Plant Physiol.* **120**: 1193-1204.
- Motte, P., Saedler, H. and Schwarz-Sommer, Z. 1998. *STYLOSA* and *FISTULATA*: regulatory components of the homeotic control of *Antirrhinum* floral organogenesis. *Development.* **125**: 71-84.
- Mouradov, A., Glassick, T.V., Hamdorf, B.A., Murphy, L.C., Marla, S.S., Yang, Y. and Teasdale, R.D. 1998. Family of MADS-box genes expressed early in male and female reproductive structures of Monterey pine. *Plant Physiol.* **117**: 55-62.
- Muller, B.M., Saedler, H. and Zachgo, S. 2001. The MADS-box gene *DEFH28* from *Antirrhinum* is involved in the regulation of floral meristem identity and fruit development. *Plant J.* **28**: 169-179.
- Munster, T., Pahnke, J., Di Rosa, A., Kim, J.T., Martin, W., Saedler, H. and Theissen, G. 1997. Floral homeotic genes were recruited from homologous MADS-box

- genes preexisting in the common ancestor of ferns and seed plants. *Proc. Natl. Acad. Sci. USA*. **94**: 2415-2420.
- Nam, J., dePhamphilis, C.W., Ma, H. and Nei, M. 2003. Antiquity and evolution of the MADS-box gene family controlling flower development in plants. *Mol. Biol. Evol.* **20**:1435-1447.
- Nam, J., Kim, J., Lee, S., An, G., Ma, H. and Nei, M. 2004. Type I MADS-box genes have experienced faster birth-and-death evolution than type II MADS-box genes in angiosperms. *Proc. Nat. Acad. Sci. USA*. **101**:1910-1915.
- Nesi, N., Debeaujon, I., Jond, C., Stewart, A.J., Jenkins, G.I., Caboche, M. and Lepiniec, L. 2002. The *TRANSPARENT TESTA16* locus encodes the *ARABIDOPSIS* B sister MADS domain protein and is required for proper development and pigmentation of the seed coat. *Plant Cell*. **14**: 2463–2479.
- Onouchi, H., Igeño, M.I., Périlleux, C., Graves, K. and Coupland, G. 2000. Mutagenesis of plants overexpressing *CONSTANS* demonstrates novel interactions among *Arabidopsis* flowering-time genes. *Plant Cell*. **12**: 885–900.
- Parenicova, L., de Folter, S., Kieffer, M., Horner, D.S., Favalli, C., Busscher, J., Cook, H.E., Ingram, R.M., Kater, M.M., Davies, B., Angenent, G.C. and Colombo, L. 2003. Molecular and phylogenetic analyses of the complete MADS-box transcription factor family in *Arabidopsis*: New openings to the MADS world. *Plant Cell*. **15**: 1538-1551.
- Pelaz S., Ditta, G. S., Baumann, E., Wisman, E. and Yanofsky, M.F. 2000. B and C floral organ identity functions require *SEPALLATA* MADS-box gene. *Nature*. **405**: 200-202.
- Pelaz, S., Tapia-Lopez, R., Alvarez-Buylla, E. R. and Yanofsky, M. F. 2001. Conversion of leaves into petals in *Arabidopsis*. *Curr. Biol.* **11**: 182-184.
- Pelucchi, N., Fornara, F., Favalli, C., Masiero, S., Lago, C., Pè, M.E., Colombo, L. and Kater, M.M. 2002. Comparative analysis of rice MADS box genes expressed during flower development. *Sex Plant Reprod.* **15**: 113–122.
- Pinyopich, A., Ditta, G. S., Savidge, B., Liljegren, S. J., Baumann, E., Wisman, E. and Yanofsky, M.F. 2003. Assessing the redundancy of MADS-box genes during carpel and ovule development. *Nature*. **42**: 85-88.
- Pnueli, L., Abu-Abeid, M., Zamir, D., Nacken, W., Schwarz-Sommer, Z. and Lifschitz, E. 1991. The MADS box gene family in tomato: Temporal expression during floral development, conserved secondary structures and homology with homeotic genes from *Antirrhinum* and *Arabidopsis*. *Plant Cell*. **1**: 255–266.
- Pnueli, L., Hareven, D., Broday, L., Hurwitz, C. and Lifschitz, E. 1994. The *TM5* MADS box gene mediates organ differentiation in the three inner whorls of tomato flowers. *Plant Cell*. **6**: 175–186.

- Prakash, A. P. and Kumar, P. P. 2002. *PkMADS1* is a novel MADS box gene regulating adventitious shoot induction and vegetative shoot development in *Paulownia kawakamii*. *Plant J.* **29**: 141-151.
- Prasad, K. and Vijayraghavan, U. 2003. Double-stranded RNA interference of a rice *PI/GLO* paralog, *OsMADS2*, uncovers its second-whorl-specific function in floral organ patterning. *Genet.* **165**: 301-2305.
- Purugganan, M. D. 1998. The molecular evolution of development. *BioEssays.* **20**: 700-711.
- Purugganan, M.D., Rounsley, S. D., Schmidt, R.J. and Yanofsky, M.F. 1995. Molecular evolution of flower development: Diversification of the plant MADS-box regulatory gene family. *Genet.* **140**: 345-356.
- Raghava, G.P.S. and Barton, G.J. 2006. Quantification of the variation in percentage identity for protein sequence alignments. *BMC Bioinformatics.* **7**: 415.
- Ratcliffe, O.J., Nadzan, G.C., Reuber, T.L. and Riechmann, J.L. 2001. Regulation of flowering in *Arabidopsis* by an *FLC* homologue. *Plant Physiol.* **126**: 122–132.
- Riechmann, J.L., and Meyerowitz, E.M. 1998. The AP2/EREBP family of plant transcription factors. *Biol. Chem.* **379**: 633-646.
- Riechmann, J.L., Krizek, B.A. and Meyerowitz, E.M. 1996a. Dimerization specificity of *Arabidopsis* MADS domain homeotic proteins APETALA1, APETALA3, PISTILLATA, and AGAMOUS. *Proc. Natl Acad. Sci. USA.* **93**: 4793–4798.
- Riechmann, J.L., Wang, M. and Meyerowitz, E.M. 1996b. DNA binding properties of *Arabidopsis* MADS domain homeotic proteins APETALA1, APETALA3, PISTILLATA and AGAMOUS. *Nucl. Acids Res.* **24**: 3134–3141.
- Rogozin, I.B., Kochetov, I.V., Kondrashov, F.A., Koonin, E.V. and Milanese, L. 2001. Presence of ATG triplets in 5' untranslated regions of eukaryotic cDNAs correlates with a 'weak' context of the start codon. *Bioinformatics.* **17**: 890-900.
- Rosin, F.M., Hart, J. K., Van Onckelen, H. and Hannapel, D.J. 2003. Suppression of a vegetative MADS box gene of potato activates axillary meristem development. *Plant Physiol.* **131**: 1613-1622.
- Roskopf, D., Manthey, I., Habich, C., Kielbik, M., Eisenhardt, A., Nikula, C., Melanie, U., Stefanie, K., Eva, G., Ursula, R. and Winfried, S. 2003. Identification and characterization of Gβ3s2, a novel splice variant of the G-protein β3 subunit. *Biochem. J.* **371**: 223-232.
- Rothnie, H.M., Reid, J. and Hohn, T. 1994. The contribution of AAUAAA and the upstream element UUUGUA to the efficiency of mRNA 3'-end formation in plants. *EMBO J.* **13**: 2200-2210.

- Rounsley, S.D., Ditta, G. S. and Yanofsky M.F. 1995. Diverse roles for MADS box genes in *Arabidopsis* development. *Plant Cell*. **7**: 1259-1269.
- Sakai, H., Medrano, L.J. and Meyerowitz, E.M. 1995. Role of *SUPERMAN* in maintaining *Arabidopsis* floral whorl boundaries. *Nature*. **378**: 199-203.
- Samach, A., Onouchi, H., Gold, S.E., Ditta, G.S., Schwarz-Sommer, Z., Yanofsky, M.F. and Coupland, G. 2000. Distinct roles of *CONSTANS* target genes in reproductive development of *Arabidopsis*. *Science*. **288**: 1613–1616.
- Sambrook, J. & Russell, D.W. 2001. *Molecular cloning. A lab manual* 3<sup>rd</sup> edition. New York: Cold Spring Harbor Lab Press.
- Santelli, E. and Richmond, T.J. 2000. Crystal structure of MEF2A core bound to DNA at 1.5 Å resolution. *J. Mol. Biol.* **297**: 437–449.
- Schindler, U., Terzaghi, W., Beckmann, H., Kadash, T. and Cashmore, A.R. 1992. DNA binding site preferences and transcriptional properties of the *Arabidopsis* factor GBF1. *EMBO J.* **11**: 1275-1289.
- Schmidt, R. J., Veit, B., Mandel, M. A., Mena, M., Hake, S. and Yanofsky, M. F. 1993. Identification and molecular characterization of *ZAG1*, the maize homolog of the *Arabidopsis* floral homeotic gene *AGAMOUS*. *Plant Cell*. **5**: 729-737.
- Schwarz-Sommer, Z., Hue, I., Huijser, P., Flor, P.J., Hansen, R., Tetens, F., Lonig, W.E., Saedler, H. and Sommer, H. 1992. Characterization of the *Antirrhinum* floral homeotic MADS-box gene *Deficiens*: evidence for DNA binding and autoregulation of its persistent expression throughout flower development. *EMBO J.* **11**: 251-263.
- Scortecci, K. C., Michaels, S. D. and Amasino, R. M. 2001. Identification of a MADS-box gene, *FLOWERING LOCUS M* that represses flowering. *Plant J.* **26**: 229-236.
- Shabalina, S.A., Ogurtsov, A.Y. and Spiridonov, N.A. 2006. A periodic pattern of mRNA secondary structure created by the genetic code. *Nucl. Acids Res.* **34**: 2428–2437.
- Sheldon, C.C., Burn, J.E., Perez, P.P., Metzger, J., Edwards, J.A., Peacock, W.J. and Dennis, E.S. 1999. The *FLF* MADS box gene: a repressor of flowering in *Arabidopsis* regulated by vernalization and methylation. *Plant Cell*. **11**: 445-458.
- Shinozuka, Y., Kojima, S., Shomura, A., Ichimura, H., Yano, M., Yamamoto, K. and Sasaki, T. 1999. Isolation and characterization of rice MADS box gene homologues and their RFLP mapping. *DNA Res.* **6**: 123-9.
- Shore, P. and Sharrocks, A.D. 1995. The MADS-box family of transcription factors. *Eur. J. Biochem.* **229**: 1–13.

- Solanas, M., Moral, R., and Escrich, E. 2001. Unsuitability of using ribosomal RNA as loading control for Northern blot analyses related to the imbalance between messenger and ribosomal RNA content in rat mammary tumors. *Anal. Biochem.* **288**: 99–102.
- Soltis, D.E., Soltis, P.S., Albert, V.A., Oppenheimer, D.G., dePamphilis, C.W., Ma, H., Frohlich, M.W. and Theissen, G. 2002. Missing links: the genetic architecture of flowers and floral diversification. *Trends Plant Sci.* **7**: 22–31.
- Sommer, H., Beltran, J.P., Huijser, P., Pape, H., Lonng, W.E., Saedler, H. and Schwarz-Sommer, Z. 1990. *Deficiens*, a homeotic gene involved in the control of flower morphogenesis in *Antirrhinum majus*: the protein shows homology to transcription factors. *EMBO J.* **9**: 605–613.
- Stahlberg, A., Hakansson, J., Xian, X., Semb, H. and Kubista, M. 2004. Properties of the reverse transcription reaction in mRNA quantification. *Clinical Chem.* **50**: 509–515.
- Sterky, F., Regan, S., Karlsson, J., Hertzberg, M., Rohde, A., Holmberg, A., Amini, B., Bhalerao, R., Larsson, M., Villarroel, R., Montagu, M.V., Sandberg, G., Olsson, O., Teeri, T.T., Boerjan, W., Gustafsson, P., Uhlen, M., Sundberg, B. and Lunderberg, J. 1998. Gene discovery in the wood-forming tissues of poplar: analysis of 5,692 expressed sequence tags. *Proc. Natl. Acad. Sci. USA.* **95**: 13330–13335.
- Sung, S.K., Yu, G.H., Nam, J., Jeong, D.H. and An, G.H. 2000. Developmentally regulated expression of two MADS-box genes, *MdMADS3* and *MdMADS4*, in the morphogenesis of flower buds and fruits in apple. *Planta.* **210**: 519–528.
- Svensson, M. E. and Engstrom, P. 2002. Closely related MADS-box genes in club moss *Lycopodium*. show broad expression patterns and are structurally similar to, but phylogenetically distinct from, typical seed plant MADS-box genes. *New Phytol.* **154**: 439-450.
- Svensson, M. E., Johannesson, H. and Engström, P. 2000. The *LAMB1* gene from the clubmoss, *Lycopodium annotinum*, is a divergent MADS-box gene, expressed specifically in sporogenic structures. *Gene.* **253**: 31-43.
- Tanabe, Y., Hasebe, M., Sekimoto, H., Nishiyama, T., Kitani, M., Henschel, K., Munster, T., Theissen, G., Nozaki, H. and Ito, M. 2005. Characterization of MADS-box genes in charophycean green algae and its implication for the evolution of MADS-box genes. *Proc. Nat. Acad. Sci. USA.* **102**: 2436-2441.
- Tandre, K., Svenson, M., Svensson, M.E. and Engstrom, P. 1998. Conservation of gene structure and activity in the regulation of reproductive organ development of conifers and angiosperms. *Plant J.* **15**: 615-623.
- Taylor, S.A., Hofer, J.M. I., Murfet, I.C., Sollinger, J.D., Singer, S.R., Knox, M.R. and Ellis, T.H.N. 2002. *PROLIFERATING INFLORESCENCE MERISTEM*, a MADS-box

- gene that regulates floral meristem identity in pea. *Plant Physiol.* **129**: 1150-1159.
- Theissen, G. 2001. Development of floral organ identity: stories from the MADS house. *Curr. Opin. Biol.* **4**: 75–85.
- Theissen, G. and Saedler, H. 2001. Floral quartets. *Nature* **409**: 469–471.
- Theissen, G., Becker, A., Di Rosa, A., Kanno, A., Kim, J.T., Münster, T., Winter, K.U. and Saedler, H. 2000. A short history of MADS-box genes in plants. *Plant Mol. Biol.* **42**: 115–149.
- Theissen, G., Kim, J. T. and Saedler, H. 1996. Classification and phylogeny of the MADS-box multigene family suggest defined roles of MADS-box gene subfamilies in the morphological evolution of eucaryotes. *J. Mol. Evol.* **43**: 484-516.
- Tonaco, I.A., Borst, J.W., de Vries, S.C., Angenent, G.C. and Immink, R.G.H. 2006. *In vivo* imaging of MADS-box transcription factor interactions. *J. Exp. Bot.* **57**: 33-42.
- Tran, H.T., Gordenin, D.A. and Resnick, M.A. 1996. The prevention of repeat-associated deletions in *Saccharomyces cerevisiae* by mismatch repair depends on size and origin of deletions. *Genet.* **143**: 1579-1587.
- van der Krol, A.I. and Chua, N.H. 1993. Flower development in petunia. *Plant Cell.* **5**: 1195-1203.
- Vandesompele, J., Katleen, D.P., Filip, P., Poppe, B., Van Roy, N., De Paepe, A. and Speleman, F. 2002. Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biol.* **3**: 34.1-34.11.
- Venables, J.P. 2004. Aberrant and alternative splicing in cancer. *Cancer Res.* **64**: 7647–7654.
- Viguera, E., Canceill, D. and Erlich, S.D. 2001. Replication slippage involves DNA polymerase pausing and dissociation. *EMBO J.* **20**: 2587-2595.
- Vrebalov, J., Ruezinsky, D., Padmanabhan, V., White, R., Medrano, D., Drake, R., Schuch, W. and Giovannoni, J. 2002. A MADS-box gene necessary for fruit ripening at the tomato *Ripening-inhibitor Rin*. locus. *Science.* **296**: 343-346.
- Walden, A.R., Wang, D.Y., Walter, C. and Gardner, R.C. 1998. A large family of *TM3* MADS-box cDNAs in *Pinus radiata* includes two members with deletions of the conserved K domain. *Plant Sci.* **138**: 167-176.
- Wang, X.Q. and Rothnagel, J.A. 2004. 5'-Untranslated regions with multiple upstream AUG codons can support low-level translation via leaky scanning and reinitiation. *Nucl. Acids Res.* **32**: 1382 - 1391.

- Weigel, D. and Meyerowitz, E.M. 1994. The ABCs of floral homeotic genes. *Cell*. **78**: 203-209.
- West, A. G. and Sharrocks A. D. 1999. MADS-box transcription factors adopt alternative mechanisms for bending DNA. *J. Mol. Biol.* **286**: 1311-23.
- West, A.G., Causier, B.G., Davies, G. and Sharrocks, A.D. 1998. DNA binding and dimerization determinants of *Antirrhinum majus* MADS-box transcription factors. *Nucl. Acids Res.* **26**: 5277-5287.
- Western, T.L. and Haughn, G.W. 1999. *BELL1* and *AGAMOUS* genes promote ovule identity in *Arabidopsis thaliana*. *Plant J.* **18**: 329-336.
- Wilkinson, M., de Andrade, S.E., Zachgo, S., Saedler, H. and Schwarz-Sommer, Z. 2000. *CHORIPETALA* and *DESPENTEADO*: general regulators during plant development and potential floral targets of *FIMBRIATA*-mediated degradation. *Development.* **127**: 3725-34.
- Witte, C.P., Tiller, S., Isidore, E., Davies, H.V. and Taylor, M.A. 2005. Analysis of two alleles of the *urease* gene from potato: polymorphisms, expression, and extensive alternative splicing of the corresponding mRNA. *J. Exp. Bot.* **56**: 91-99.
- Wray, G.A., Hahn, M.W., Abouheif, E., Balhoff, J.P., Pizer, M., Rockman, M.V. and Romano, L.A. 2003. The evolution of transcriptional regulation in eukaryotes. *Mol. Biol. Evol.* **20**:1377-1419.
- Xiao, H., Wang, Y., Liu, D., Wang, W., Li, X., Zhao, X., Xu, J., Zhai, W. and Zhu, L. 2003. Functional analysis of the rice *AP3* homologue *OsMADS16* by RNA interference. *Plant Mol. Biol.* **52**: 957-966.
- Yamaguchi, T., Lee, D.Y., Miyao, A., Hirochika, H., An, G. and Hirano, H.Y. 2006. Functional diversification of the two C-class MADS box genes *OSMADS3* and *OSMADS58* in *Oryza sativa*. *Plant Cell*.**18**: 15-28.
- Yang, Y., Fanning, L. and Jack, T. 2003. The K domain mediates heterodimerization of the *Arabidopsis* floral organ identity proteins, APETALA3 and PISTILLATA. *Plant J.* **33**: 47-59.
- Yanofsky, M.F., Ma, H., Bowman, J.L., Drews, J.L., Feldmann, K.A. and Meyerowitz, E.M. 1990. The protein encoded by the *Arabidopsis* homeotic gene *agamous* resembles transcription factors. *Nature.* **346**: 35-39.
- Yu, H. and Goh, C. J. 2000. Identification and characterization of three orchid MADS-Box genes of the *API/AGL9* subfamily during floral transition. *Plant Physiol.* **123**: 1325-1336.
- Zachgo, S., Ede, S.A., Motte, P., Trobner, W., Saedler, H. and Schwarz-Sommer, Z. 1995. Functional analysis of the *Antirrhinum* floral homeotic *DEFICIENS* gene



*in vivo* and *in vitro* by using a temperature-sensitive mutant. *Development*. **121**: 2861-75.

- Zachgo, S., Saedler, H. and Schwarz-Sommer, Z. 1997. Pollen-specific expression of *DEFH125*, a MADS-box transcription factor in *Antirrhinum* with unusual features. *Plant J.* **11**: 1043-50.
- Zahna, L.M., Leebens-Macka, J.H., Arringtona, J.M., Hua, Y., Landherra, L.L., dePamphilisa, C.W., Becker, A., Theissen, G. and Ma, H. 2006. Conservation and divergence in the *AGAMOUS* subfamily of MADS-box genes; evidence of independent sub- and neofunctionalization events. *Evol. Dev.* **8**: 30-45.
- Zhang H. and Forde. B.G. 1998. An *Arabidopsis* MADS box gene that controls nutrient-induced changes in root architecture. *Science*. **279**: 407-409.
- Zhang, P., Tan, H.T., Pwee, K.H. and Kumar, P.P. 2004. Conservation of class C function of floral organ development during 300 million years of evolution from gymnosperms to angiosperms. *Plant J.* **37**: 566–577.
- Zhao, J., Hyman, L., and Moore, C. 1999. Formation of mRNA 3' ends in eucaryotes: mechanism, regulation and interrelationships with other steps in mRNA synthesis. *Microbiol. Mol. Biol. Rev.* **63**: 405-445.
- Zuker, M. 2003. Mfold web server for nucleic acid folding and hybridization prediction. *Nucl. Acids Res.* **31**: 3406–3415.