

**PHYSICOCHEMICAL CHARACTERISTICS OF DIFFERENT RICE
VARIETIES FOUND IN SABAH**

NEOH WEI TING

**DISSERTATION SUBMITTED IN PARTIAL FULFILMENT OF THE
REQUIREMENTS FOR THE DEGREE OF BACHELOR OF
AGRICULTURE SCIENCE WITH HONOURS**

**CROP PRODUCTION PROGRAM
SCHOOL OF SUSTAINABLE AGRICULTURE
UNIVERSITI MALAYSIA SABAH
2011**



UMS
UNIVERSITI MALAYSIA SABAH

DECLARATION

I hereby declare that this dissertation is based on my original work except for citations and quotations which have been duly acknowledged. I also declare that no part of this dissertation has been previously or concurrently submitted for a degree at this or any other university.



NEOH WEI TING
BR07110014
15th APRIL 2011

VERIFICATION


VERIFIED BY

1. Mr. Lum Mok Sam
SUPERVISOR



LUM MOK SAM
KETUA PROGRAM HG 35 HORTIKULTUR & LANSKAP
SEKOLAH PERTANIAN LESTARI
UNIVERSITI MALAYSIA SABAH

2. Mr. Clament Chin Fui Seung
EXAMINER 1




3. Mr. Januarius Gobilik
EXAMINER 2



JANUARIUS GOBILIK
Lecturer / Academic Supervisor
School of Sustainable Agriculture
Universiti Malaysia Sabah

4. Assoc. Prof. Dr. Mahmud Haji Sudin
DEAN



ASSOCIATE PROF. DR. MAHMUD-SUDIN
DEAN
SCHOOL OF SUSTAINABLE AGRICULTURE
UNIVERSITI MALAYSIA SABAH

ACKNOWLEDGEMENT

I would like to express the deepest appreciation to my supervisor, Mr. Lum Mok Sam for guiding me in my final year project and providing invaluable help and encouragement throughout this study. Without his guidance and persistent help this dissertation would not have been possible.

In addition, a thank you to Mdn. Christina Kungin, Mdn. Ahjia bt Osman, Mr. Malcolm Boxey Jilimin, Mr. Ahmad bin Manik, for their kind assistance and patience in helping me go through my project.

Last but not least, I would like to dedicate my special thanks to all my friends for their wonderful and endless support during my hard time in the whole progress of this dissertation writing. Once again, it was my pleasure and honor to have the support from all of you.



ACKNOWLEDGEMENT

I would like to express the deepest appreciation to my supervisor, Mr. Lum Mok Sam for guiding me in my final year project and providing invaluable help and encouragement throughout this study. Without his guidance and persistent help this dissertation would not have been possible.

In addition, a thank you to Mdn. Christina Kungin, Mdn. Ahjia bt Osman, Mr. Malcolm Boxey Jilimin, Mr. Ahmad bin Manik, for their kind assistance and patience in helping me go through my project.

Last but not least, I would like to dedicate my special thanks to all my friends for their wonderful and endless support during my hard time in the whole progress of this dissertation writing. Once again, it was my pleasure and honor to have the support from all of you.

ABSTRACT

Five different rice varieties found in Sabah (White rice, Red rice, Black rice, Brown rice and Aroma rice) were evaluated for physicochemical characteristics. The relationships between different characteristics were determined using Pearson correlation. Different varieties showed significantly variations in their physicochemical characteristics. Thousand grains weight, length–breadth (L/B) ratio, and moisture content varied between 15.30 g to 20.90 g, 2.18 to 3.33, and 12.30% to 16.40%, respectively among the various varieties. Minimum cooking time, water uptake ratio, gruel solid loss, and elongation ratio were found within 21.97 min to 47.20 min, 2.30 g g⁻¹ to 3.51 g g⁻¹, 6.02% to 13.62% and 1.25 to 1.65 respectively. Brown rice showed a lower elongation ratio and water uptake ratio, and a higher cooking time than the other rice varieties. The water uptake and elongation ratio of rice kernels showed negative correlation with cooking time, with a correlation coefficient of -0.960 and -0.900 respectively ($P \leq 0.05$). Starch content, starch swelling power (SSP), and amylose content (AC) of studied rice varieties varied between 33.73% to 64.98%, 9.31 g g⁻¹ to 13.01 g g⁻¹ and 10.83% to 14.93% respectively. Protein, phosphorus (P), sodium (Na), potassium (K), magnesium (mg) and copper (Cu) contents ranged from 13.13% to 24.60%, 19.95 mg/100 g to 27.60 mg/100 g, 74.98 mg/100 g to 627.17 mg/100 g, 68.40 mg/100 g to 126.73 mg/100 g and 0.57 mg/100 g to 1.63 mg/100 g, respectively among different rice varieties. The highest contents of these elements were observed in Red rice, Black rice and Brown rice. The result obtained showed that the traditional rice (Red rice, Black rice and Brown rice) found in Sabah had the better physicochemical characteristics and higher nutritional value as compare to cultivated White rice and Aroma rice.

CIRI-CIRI FIZIKOKIMIA BAGI VARIETI BERAS BERLAINAN YANG DITEMUI DI SABAH

ABSTRAK

Lima jenis varieti bera berlainan yang ditemui di Sabah (Beras putih, Beras Merah, Beras Hitam, Beras Coklat Dan Beras Wangi) telah dinilai dari segi ciri-ciri fizikokimia. Hubungan antara ciri-ciri yang berbeza ditentukan dengan menggunakan korelasi Pearson. Varieti yang berbeza menunjukkan variasi yang signifikan dalam fizikokimia mereka. Berat seribu bijih, nisbah panjang lebar (P/L), dan kelembapan di antara pelbagai jenis beras adalah berbeza antara 15.30 g hingga 20.90 g, 2.18 hingga 3.33 dan 12.30% hingga 16.40% masing-masing. Minima masa memasak, nisbah serapan air, kehilangan bubur padat, dan nisbah elongasi berbeza antara 21.97 minit hingga 47.20 minit, 2.30 g g⁻¹ hingga 3.51 g g⁻¹, 6.02% hingga 13.62% dan 1.25 hingga 1.65 masing-masing. Beras Coklat menunjukkan nisbah pemanjangan dan nisbah serapan air yang lebih rendah, dan minima masa memasak yang lebih tinggi berbanding dengan jenis beras yang lain. Nisbah serapan air dan nisbah pemanjangan beras berkorelasi negatif dengan masa memasak, dengan pekali korelasi -0.900 dan -0.960 masing-masing (P≤0.05). Kandungan kanji, daya pembengkakan kanji, dan kandungan amilosa berbeza antara 33.73% hingga 64.98%, 9.31 hingga 13.01 g g⁻¹ dan 10.83% hingga 14.93% masing-masing. Kandungan protin, fosforus, natrium, kalium, magnesium dan kuprum adalah 13.13% hingga 24.60%, 19.95 mg/100 g hingga 27.60 mg/100 g, 74.98 mg/100 g hingga 627.17 mg/100 g, 68.40 mg/100 g hingga 126.73 mg/100 g dan 0.57 mg/100 g hingga 1.63 mg/100 g masing-masing. Beras Merah, Beras Hitam dan Beras Coklat mempunyai kandungan mineral yang lebih tinggi berbanding dengan beras yang lain. Keputusan yang diperolehi menunjukkan bahawa beras tradisional (Beras Merah, Beras Hitam dan Beras Coklat) yang ditemui di Sabah memiliki sifat fizikokimia yang lebih baik dan nilai gizi yang lebih tinggi berbanding dengan Beras Putih dan Beras Aroma.

TABLE OF CONTENTS

CONTENT	PAGE
DECLARATION	ii
VERIFICATION	iii
ACKNOWLEDGEMENT	iv
ABSTRACT	v
<i>ABSTRAK</i>	vi
TABLE OF CONTENT	vii
LIST OF TABLES	ix
LIST OF FIGURES	x
LIST OF FORMULA	xi
LIST OF SYMBOLS, UNITS AND ABBREVIATIONS	xii
CHAPTER 1 INTRODUCTION	1
1.1 Foreword	1
1.2 Justifications	3
1.3 Objective	4
1.4 Hypothesis	4
CHAPTER 2 LITERATURE REVIEW	5
2.1 Rice	5
2.2 Types of Rice	5
2.2.1 Indica and Japonica	6
2.3 Rice in Malaysia	7
2.4 Nutritional Value	8
2.5 Rice Quality	8
2.6 Physico-chemical characteristics of Rice	10
2.6.1 Appearance	10
2.6.1.1 Size and Shape	11
2.6.1.2 Colour	12
2.6.2 Thousand Grains Weight (TGW)	13
2.6.3 Moisture Content (%)	14
2.6.4 Gelatinization Temperature (GT)	14
2.6.5 Starch Content	15
2.6.6 Starch Swelling Power (SSP)	16
2.6.7 Amylose Content (AC)	18
2.6.8 Cooking Time	19
2.6.9 Water Uptake Ratio	20
2.6.10 Elongation Ratio	21
2.6.11 Gruel Solid Loss	23
2.6.12 Protein Content	24
CHAPTER 3 METHODOLOGY	26
3.1 Materials	26
3.2 Physical Characteristics of Rice	26
3.2.1 Grain Appearance (Length, Shape and Colour)	26
3.2.2 A Thousand Grain Weight (g)	27
3.3 Chemical Characteristics of Rice	27
3.3.1 Moisture Content (%)	27
3.3.2 Gelatinization Temperature (GT)	27
3.3.3 Starch Content	28

3.3.4	Starch Swelling Power (SSP)	28
3.3.5	Amylose Content (AC)	29
3.3.6	Protein Content	30
3.4	Cooking Characteristics of Rice	30
3.4.1	Cooking Time	30
3.4.2	Water Uptake Ratio	30
3.4.3	Elongation Ratio	31
3.4.4	Gruel Solid Loss	31
3.5	Mineral Content of Rice	31
3.5.1	Phosphorus Content	31
3.5.2	Mineral Extraction	32
3.6	Data Analysis	33
CHAPTER 4 RESULTS		34
4.1	Morphological and Physical Characteristics	34
4.2	Chemical Characteristics	36
4.3	Cooking Characteristics	42
4.4	Mineral Contents	45
CHAPTER 5 DISCUSSION		47
5.1	Rice Appearance	47
5.1.1	Size and Shape of Rice	47
5.1.2	Colour of Rice	48
5.2	Thousand Grains Weight (TGW)	48
5.3	Moisture Content (%)	49
5.4	Gelatinization Temperature (GT)	50
5.5	Starch Content	51
5.6	Starch Swelling Power (SSP)	52
5.7	Amylose Content (AC)	52
5.8	Cooking Time	53
5.9	Water Uptake Ratio	54
5.10	Elongation Ratio	55
5.11	Gruel Solid Loss	56
5.12	Protein Content	56
5.13	Mineral Contents	57
CHAPTER 6 CONCLUSION		60
6.1	Conclusion	60
6.2	Recommendations	62
6.2.1	Nutrition Facts Labeling on the Rice Packaging	62
6.2.2	Served as Gene Sources in the Rice Breeding Program for Grain Quality	62
REFERENCES		63
APPENDICES		72

LIST OF TABLES

Table		Page
3.1	Size classification of rice	26
3.2	Shape classification of rice	26
3.3	Numerical scale for scoring gelatinous temperature	27
4.1	Morphological and physical characteristics of different rice varieties in Sabah	36
4.2	Chemical characteristics of different rice varieties in Sabah	37
4.3	Numerical scale for scoring gelatinization temperature of rice	37
4.4	Cooking characteristics of different rice varieties in Sabah	42
4.5	Mineral contents of different rice varieties in Sabah	46

LIST OF FIGURES

Figure		Page
4.1	The appearance of rice	35
4.2	The alkali digestion for rice	38
4.3	Correlation between cooking time and gelatinization temperature	39
4.4	Correlation between water uptake ratio and gelatinization temperature	40
4.5	Correlation between elongation ratio and gelatinization temperature	40
4.6	Correlation between starch content and SSP	41
4.7	Correlation between cooking time and water uptake ratio	43
4.8	Correlation between cooking time and elongation ratio	43
4.9	Correlation between L/B ratio and water uptake ratio of cooked rice	44
4.10	Correlation between L/B ratio and elongation ratio	44

LIST OF FORMULA

Formula	Page
3.1 % moisture content = $\frac{\text{fresh weight} - \text{dry weight}}{\text{fresh weight}} \times 100\%$	27
3.2 $\text{SSP} = \frac{\text{Sediment weight}}{\text{Mass of dry starch}} \times (100\% - \% \text{ of total mass of dried supernatant})$	28
3.3 % P = $\frac{1}{\text{Weight of dried sample (g)}} \times \frac{5 \text{ mL}}{1 \text{ mL}} \times \frac{100}{1} \times R$	32

LIST OF SYMBOLS, UNITS AND ABBREVIATIONS

AC	Amylose content
ANOVA	Analysis of variance
DOM	Degree of milling
FAO	Food and Agriculture Organization
GT	Gelatinous temperature
HRY	Head rice yield
IRRI	International Rice Research Institute
L/B	Length/Breadth
MC	Moisture content
SSP	Starch swelling power
TGW	Thousand grain weight
USDA	United State Department of Agriculture

CHAPTER 1

INTRODUCTION

1.1 Foreword

Rice (*Oryza sativa L.*) has been the major staple food in the world, especially Asian countries since ancient time. Generally, rice is unique due to the whole grain is eaten exclusively by human in contrast with other cereal grains such as wheat, barley and sorghum. It feeds more than one third of the world's population by supplying high value of carbohydrate and consisting of more than 50% of the human's daily calories intake (Anjum *et al.*, 2007).

Rice is very rich in carbohydrate. Although rice has a relatively low in protein content but still sufficient to sustain the daily needs for an adult. Moreover, rice can help to overcome malnutrition problem with small amounts of minerals. Minerals present in rice like potassium, manganese, iron and zinc play an important role in body regulatory functions (FAO, 2004). Besides just being as food, rice has the potential to alleviate poverty and hunger twice as compare to other agricultural products.

Among the cereals, rice has a great diversity with more than 120,000 different varieties (IRRI, 2008). All the varieties can be grouped into two sub-species: Japonica and Indica (Khush *et al.*, 2000). Even though there are many rice varieties to be found in the market, the preferences varied from one country to another depending on the consumer preference or the preference in the market. Some of them will prefer milled-white rice while other likes brown rice or Aroma rice. Some country will have great demand on long-grain rice while other prefers short-grain rice.

Like other countries, rice is also important for Malaysian. Rice plays a crucial cultural role in Malaysia beyond providing nourishment. For thousands of years, it has shaped the cultures, economies and diets of many regions in Malaysia. It plays an important role as the essence of Malaysian life. In Malaysia, each ethnic cuisine has developed individual use of rice. For example, rice is cooked as desserts which known as "kuih-muih" for Malayan. Other than that, rice is ground into flour for making "kuih bakul" by Chinese during celebration of Chinese New Year. Rice wine is produced by Kadazan and Dusun ethnics through fermentation. These unique cultures of rice usage in Malaysia have been kept until today from one generation to another generation.

Government has put huge effort in increasing the rice production to fulfill the local demand. Even so, it is unable to meet the huge demand and has to import from neighboring countries such as Thailand, Vietnam and Philippine. But, rice production in Malaysia is indeed gradually increasing from 2,169,000 tonnes in year 2000 to 2,354,000 tonnes in year 2008 (USDA, 2009). Rice produced in Malaysia varies significantly especially traditional rice in Sabah. Traditional rices in Sabah market show variations in colour such as white, brown, red and black. Traditional rices in Sabah have largely contributed in the rice market but they have not yet been documented. Therefore, it is important to have a proper study and research on these rice varieties.

Generally, rice in Sabah is grown by farmers in small farm size of one hectare or less. Certified rice seeds approved by the government normally are a luxury item that is beyond the farmer's financial ability and what they can afford is the traditional seeds that sold in the market or seeds they collected during harvesting. Due to small scale of cultivation, these rice seeds have not yet been proven on their eating and cooking quality. However, we cannot deny their quality since they still have a foothold in this competitive market. This proved that the quality of those traditional varieties are accepted and preferred by the local consumers. Therefore, the traditional rice varieties should own a chance to prove their quality.

With the rapid economic growth in Malaysia, consumers are becoming wiser in their choice of quality of rice. The quality of rice becomes the choice determinant factor. Many countries become self-sufficiency in rice production during this few ten years and starting involve in global market. The competitiveness of rice export among these countries becomes more drastic and again verifies the importance of rice quality. Therefore, it is necessary to understand the factors contributing to the quality of rice in order to meet the requirement of consumer.

There are many varieties of rice and each type has its own characteristics. Unlike other cereals, rice is consumed as a whole grain. Therefore, the physical appearance of rice such as size, shape and colour are of utmost importance. However, quality of rice does not only involve the physical appearance, rather it encompasses both the chemical and cooking properties. In conclusion, physicochemical characteristics of rice are the determinant factor. This fact has been proved from previous studies and researches (Singh *et al.*, 2003; Singh *et al.*, 2005; Yadav *et al.*, 2007). Present study is designed to access the physicochemical characteristics of Sabah varieties to better understanding for the consumers and world exporters.

1.2 Justification

Determination for rice quality is one of the common problems faced by the consumer (end users) especially during the purchasing of rice. Consumers basically purchase their rice based on basic visual observation which show low awareness on rice quality. Lacking of understanding regarding the essential factors which determined the rice quality cause the consumers to make the wrong judgment and end up buying low quality of rice. Therefore, this study may lend a hand to consumer by providing them some rice quality indicator's recommendation. With comprehend on those quality indicators, consumer welfare can be satisfied easily by consuming high quality of rice.

Nowadays, low quality rice still can be found easily in local market. This phenomenon is caused by the lacking of rice quality measurement and standard to monitor the quality assurance and food security of rice production. Variation in rice quality might be due to the lacking of inspection and supervision under the standard quality assurance system. Therefore, a locally suited quality control system should be established based on the results in this study. Consequently, all the rice produced which undergoes the strict regulation can be an assurance of the quality.

Traditional rice varieties which place a foothold in the local market of Sabah and serve the capability in accomplishing consumer satisfaction for rice quality but they are still not been documented so far by government. Therefore, this study help to discover the physicochemical characteristics of those rice varieties in term of verifying their quality and set foot on the way to be certified by government. As certified rice, the rice will earn more attention from government that they will strive to ensure the success of the rice production in Malaysia.

Traditional rice in Sabah is usually grown by farmer in small scale basis. Those rice varieties only been recognized by local people in the particular rural local market. Traditional rice production will remains unchanged due to the similar demand in the market. This is the time to publish and recommend the traditional rice in Sabah to other places as well as other countries. Once the rice varieties have been certified for their quality, then it can be market nationally or globally where the consumers had higher awareness on rice quality. The increasing demand on traditional Sabah's rice will increase farmers' income and profits as the rice production and yield is getting higher. On the other hand, Malaysia will become famous in international trade by supplying high quality of rice.

1.3 Objective

The objective of this study is to study the physicochemical characteristics of rice varieties found in Sabah.

1.4 Hypothesis

- H₀: There are no differences in physicochemical characteristics among the rice varieties.
- H₁: There are differences in physicochemical characteristics among the rice varieties.

CHAPTER 2

LITERATURE REVIEW

2.1 Rice

Rice plant (*Oryza sativa L.*) belongs to the tribe Oryzae under the sub-family Pooideae in the grass family Gramineae (Poaceae) (Khush *et al.*, 2000; Sparks, 2004). Rice is one of the most essential cereals and commercially more than two thousand varieties of rice are grown throughout the world (Deepa *et al.*, 2008). Rice is well-off genetic diversity by thousands of varieties grown throughout the world. In its natural un-milled state rice comes in many different colours, including brown, red, purple and even black. These colourful rice varieties are often prized for their health assets (FAO, 2004).

Rice is one of the leading food crops of the world and is a staple food of around one-half of the world population (Singh *et al.*, 2005; Yu *et al.*, 2010). Rice is the major staple food for 17 countries in Asia and the Pacific, nine countries in North and South America and eight countries in Africa. Rice supplies 20% of the world's dietary energy and 30% of the dietary energy for the Asian countries such as Malaysia, Indonesia, Philippines and China (FAO, 2004).

2.2 Types of Rice

The genus *Oryza* has two cultivated and twenty-one wild species. The cultivated rice, *Oryza Sativa* is grown all over the world. While for the African cultivated rice, *Oryza glaberrima* is grown in West Africa on a small scale. From its subtropical origins rice is now available between 55°N in China and 36°S in Chile. It is able to grown under diverse environmental conditions such as irrigated, rain-fed lowland, rain-fed upland and flood prone ecosystems. Besides that, human selection and rice adaptation to diverse environmental conditions has resulted in numerous varieties. Moreover, cultivation and farmer selection for centuries under varied growing conditions has



then resulted in a myriad of rice varieties. It is estimated that about 120,000 distinct rice varieties exist in the world after removing duplicates. About 80,000 of them are preserved in Genetic Resources Center of International Rice Research Institute (IRRI). China has about 40,000 varieties and India about 25,000 varieties in the gene banks. And other countries have the rest collections (Khush, 1997).

Oryza sativa varieties can be separated into two subspecies, the indica and japonica. Their origin showed as the result of selection by human in the process of domestication and selection of the wild rice under different environmental conditions. From the time 40 years ago, indica and japonica rice were considered by most rice scientists to be the most ecogeographic races. As hybridization between the groups continues, perhaps even that distinction will disappear, because of any of the identifying japonica and indica characteristics can be relocated in either direction through crossing and selection (Robert, 1979; Virmani, 1994).

2.2.1 Indica and Japonica

The indica rice varieties grown throughout the tropics, exhibit little tolerance to cold temperatures and respond in grain yield only to low applications of fertilizer. It is considerable drought tolerance and resistance to insect and disease attack. In general, the grains of indicas are medium-long to long, and medium to high content of amylose, causing the cooked rice to be dry and fluffy and to show little disintegration during cooked (Robert, 1979; Khush *et al.*, 2000).

Japonica rice varieties grow throughout the temperate countries such as Japan, Italy, Portugal and Spain. They are generally resistant to water-lodging and are more nitrogen responsive in their yield. Japonica has lower disease and insect resistance as compared to indica type. Generally, the grain is shorter and wider in its size and shape. The amylose content of the Japonica is lower, and this will result the rice stickier and glossier during cooking. The grains tend to disintegrate if boiled in too long period of time (Robert, 1979; Khush *et al.*, 2000).

The differences between indica and japonica were studied by Toriyama *et al.* (2005). Cooked indica rice is generally hard but not sticky. However, Japanese people are more favor on japonica cooked rice because of its moderate elasticity and stickiness. It is normally said that hardness and stickiness of cooked rice are influenced by the amylase content in rice. Japonica rice has lower amylose content and moderate elasticity and stickiness.

2.3 Rice in Malaysia

Rice (*Oryza sativa L.*) or nasi is the basic food of Malaysia and the most important source of employment and income of the rural population. As Asian rice demand is projected to increase by 30% in 2010, a sustainable approach of rice production has become important. In order to fulfill the rice demand for the growing population and with the target rice self-sufficiency of 65%, the government has to increase its rice productivity (Najim *et al.*, 2007). Currently, the self-sufficiency level of rice in Malaysia is about 75% (Vaghefi *et al.*, 2011).

Grain elongation, amylose content and aroma are key determinants of high quality rice varieties (Golam *et al.*, 2004). There is an increasing demand for quality rice in domestic markets. Almost all aromatic rice is imported, of which the majority are from Thailand and India. Many of our traditional rice varieties have aroma but there was only little effort to utilize the trait from such varieties. Malaysian rice breeders have already started to breed for high quality rice, where grain elongation, amylose content and aroma are major components in the special high quality breeding programme (Mohamad *et al.*, 2008).

Worldwide, there are more than 120,000 different rice varieties (IRRI, 2008), although only small number offers quality acceptable to be grown commercially in Malaysia. The rice germplasm collection at the Rice Genebank (Malaysia) stands at 11,470 registered accessions (Mustapha, 2007).

2.4 Nutritional Value

Rice has been considered as the best staple food among all cereal groups. It is a starchy staple food, which gives a large portion of dietary energy for around 90% of Asia population. White and brown rice contains about 90% and 75 to 85% carbohydrates, respectively. It has higher digestibility, biological value and protein efficiency ratio owing to presence of higher concentration (~ 4%) of lysine. Minerals like magnesium, phosphorus and calcium are present along with some traces of iron, zinc, copper and manganese. Mineral components were significantly higher in brown rice as compared to white rice (Shabbir *et al.*, 2008).

Many minerals are vital for body normal metabolic functions and are necessary components in a balanced diet. Calcium, phosphorus, sodium, potassium, and magnesium are required in large quantities and iron, copper, manganese, and zinc in smaller amounts. However, the ability of plants to assimilate minerals may vary with regard to variety and environmental conditions (Swain *et al.*, 1977; Fink *et al.*, 2008).

In addition to being a rich source of dietary energy, rice is a good source of thiamine, riboflavin and niacin. Unmilled rice contains a significant amount of dietary fiber. The amino acid profile of rice shows that it is high in glutamic and aspartic acid, while lysine is the limiting amino acid (FAO, 2004).

2.5 Rice Quality

Grain quality includes not only the traditional physical and visual properties of the rice grain but also cooked rice texture as indexed by apparent amylase content, alkali spreading value (index of gelatinization temperature), gel consistency, actual texture measurement of cooked grain, and protein content (nutritional value). Texture measurement is proceed primarily to verify the classification of amylose, but instrument methods are not sensitive enough to differentiate cooked rice within the same amylose type, as do sensory evaluation and processing (IRRI, 1991).

In many regions of the world, rice is the most important part of the human diet, so the consumer's daily bowl of rice needs to be safe and of good quality. In 1995, the Codex Alimentarius Commission agreed to adopt safety and quality criteria for the rice that is produced for human consumption (Codex Standard for Rice). These standards for rice are accepted by the World Trade Organization (WTO), so the Food and Agriculture Organization of the United Nations (FAO) and the World Health Organization (WHO) take care to ensure that they have a sound scientific basis (FAO, 2004).

Grain quality in rice is very difficult to define with precision as preferences for quality vary from country to country. Few people realize its complexity and the various quality components involved. The concept of quality varies according to the preparations for which grains are to be used. Although some of the quality characteristics desired by growers, millers, and consumers may be the same, each may place different emphasis on various quality characteristics. For instance, the miller's basis of quality is dependent upon total recovery and the proportion of head rice and brokens on milling. Consumers base their concept of quality on grain appearance, grain size and shape, behavior upon cooking, and the taste, tenderness, and flavor of cooked rice (Graham, 2002).

Rice quality is considered to have two general aspects. First are the milling, cooking, and processing qualities, which refer to suitability of the grain for a particular end-use. The second aspect is the physical quality, which means cleanliness, soundness, and being free from foreign materials. Usually, physical quality and milling, cooking, and processing qualities are interrelated in that certain market grades of rice are more suitable than others for consumer products (Toriyama *et al.*, 2005).

Rice forms an integral part of the culinary tradition for many cultures. Different cultures have different preferences regarding the taste, texture, colour and stickiness of the rice varieties that they consume. For example, dry flaky rice is eaten in South Asia and the Middle East; moist sticky rice in Japan, Taiwan Province of China, the Republic of Korea, Egypt and northern China; and red rice in parts of southern India. Many countries have signature rice recipes, such as sushi, fried rice, curry, paella, risotto, pancit, and beans with rice. There are also many sweets and candies made from rice (FAO, 2004).

Rice has a great diversity in its genetic background, amylose content, grain shape, and cooking quality. Varietal properties such as grain size, shape, thousand-kernel weight, hardness and bulk density affect the grain quality (Yadav *et al.*, 2007). The cooking quality of rice is determined on the basis of the variety and its physicochemical properties, but mainly the amylose content (Sujatha *et al.*, 2004). Shabbir *et al.* (2008) also had same conclusion on rice quality. The consumers are interested in to get rice of good cooking and eating quality. These characteristics are largely depending on the physico-chemical properties of starch, which make up 90% of milled rice. Other important characteristics like apparent amylose content (AC), water absorption ratio, volume expansion ratio and final starch gelatinization temperature (GT) collectively determine cooking and eating qualities of rice.

2.6 Physicochemical Characteristics of Rice

2.6.1 Appearance

Rice grain colour, size and shape are important criteria to understand the physical properties of rice (Deepa *et al.*, 2008). As rice is consumed and processed in whole kernel form, the physical attributes of the intact endosperm are the foremost importance for its market price all the time. The uniformity in physical dimensions of rice, such as length, breadth, shape and weight, also affect its market value (Virmani, 1994).

The appearance of rice is the focus point of consumers. Rice grain appearance depends on the size and shape of the grain and translucency and chalkiness of the grain. The physical dimensions of rice grains are very take seriously to those engaged in the many facets of the rice industry. Rice varieties objectively classified into grain-type categories based upon two physical parameters: length and shape. Length is a measure of the rice grain in its greatest dimension. Grain size and shape can be visually classified, and more exact measurements are needed for a more critical comparison of varieties (Graham, 2002).

2.6.1.1 Size and Shape

Grain size and shape are the first criteria that breeders consider in for the rice quality to develop new varieties for release for commercial production (Graham, 2002). Rice grain size and shape are important to consumers. This is because they determine the physical appearance and affect the cooking quality of the rice grain. In addition, rice size is important in the evolution of cereal crops because people tended to select for large seed size during the early domestication (Li *et al.*, 2004).

Rice varieties may be objectively classified into grain-type categories based on two physical parameters, the length and shape. Length is a measure of the rice kernel in its greatest dimension. While shape is determined by the length: width ratio (Khush *et al.*, 2000). Methods for measuring rice grains include of using photographic enlargers to magnify grains or simply measuring with a ruler the thickness, width and length of several rice grains placed in adjacent positions for particular measurements. Even though it will consumer longer period of time, probably it is the simplest methods for thickness, width and length determination (Luh, 1991).

The rice varieties can be classified as short grain varieties (L/B ratio less than 2.0), medium grain varieties (L/B ration less than 3.0) and long grain varieties (L/B more than 3.0) (Graham, 2002). While according to USDA (2005), rice is primarily classified according to its grain shape. The category known as long grain contains milled rice that is around three times longer than it is wide. The rice with 2.1 to 2.9 times longer than it is wide is medium grain rice. The short grain is less than two times longer than it is wide (USDA, 2005).

Preferences for grain size and shape differ from one group of consumers to another. Some of them favor short bold grains, some prefer medium-long grains, and others highly prize long slender grains. Generally, long grains are preferred in the Indian subcontinent, but, in Southeast Asia, they are more preferred on medium to medium-long rice. In temperate areas, short-grain varieties are their choice. However, there is a strong demand for long-grain rice on the international market (Graham, 2002).

REFERENCES

- Adu-Kwarteng, E., Ellis, W.O., Odurob, I. and Manful, J. T. 2003. Rice grain quality: a comparison of local varieties with new varieties under study in Ghana. *Food Control* **14(7)**: 507-514
- Ahrohmit, A., Ledward, D. A. and Niranjana, K. 2006. High pressure induced water uptake characteristics of Thai glutinous rice. *Journal of Food Engineering* **72(3)**: 225-233
- Akubugwo, I. E., Obasi, N. A., Chinyere, G. C. and Ugbo, A. E. 2007. Nutritional and chemical value of *Amaranthus hybridus* L. leaves from Afikpo, Nigeria. *African Journal of Biotechnology* **6(24)**: 2833-2839
- Amissah, J. G. N., Ellis, W. O., Oduroa, I. and Manful, J. T. 2003. Nutrient composition of bran from new rice varieties under study in Ghana. *Food Control* **14(1)**: 21-24
- Anjum, F. M., Pasha, I., Bugti, M. A. and Butt, M. S. 2007. Mineral composition of different rice varieties and their milling fractions. *Pakistan Journal of Agricultural Science* **44(2)**: 332-336
- AOAC. 1984. *Standard Official Methods of Analysis of the Association of Analytical Chemists* (14th edition). Washington, DC: Association of Official Analytical Chemists
- AOAC. 2000. *Official Methods of Analysis* (17th edition). Washington, DC: Association of Official Analytical Chemists
- Bai, X. F., Luo, L. J., Yan, W. H., Kovi, M. R., Zhan, W. and Xing, Y. Z. 2010. Genetic dissection of rice grain shape using a recombinant inbred line population derived from two contrasting parents and fine mapping a pleiotropic quantitative trait locus qGL7. *BMC Genetics* **11**: 16
- Balch, P. A. 2003. *Prescription for Dietary Wellness*. New York: Penguin Group (USA) Inc.
- Balch, P. A. 2006. *Prescription for Nutritional Healing*. England: Penguin Group
- Batres-Marquez, S. P. and Jensen, H. H. 2005. Rice Consumption in The United States: New Evidence From Food Consumption Surveys. Center for Agricultural and Rural Developments, Iowa State University, Ames, Iowa.
- Bett-Garber, K. L., Champagne, E. T., Ingram, D. A. and McClung, A. M. 2007. Influence of water-to-rice ratio on cooked rice flavor and texture. *Cereal Chemistry* **84(6)**: 614-619
- Bhattacharya, K. R., and Sowbhagya, C. M. 1971. Water uptake by rice during cooking. *Cereal Science Today* **16**: 420-424

- Brink, M. and Belay, G. 2006. *Cereals and Pulses*. Netherlands: PORTA Foundation / Backhuys Publishers / CTA
- Brown, T. and Brown K. 2011. *Biomolecular Archaeology: An Introduction*. USA: Wiley-Blackwell
- Cagampang, C. B., Perez, C. M. and Juliano, B. O. 1973. A gel consistency for eating quality of rice. *Journal of Science of Food and Agriculture* **24**: 88-94
- Cameron, D. K. and Wang, Y. J. 2005. A better understanding of factors that affect the hardness and stickiness of long-grain rice. *Cereal Chemistry* **82(2)**: 113-119
- Chanput, W., Theerakilkait, C. and Nakai, S. 2009. Antioxidative properties of partially purified barley hordein, rice bran protein fractions and their hydrolysates. *Journal of Cereal Science* **49**: 422-428
- Chaudhary, R. C. 2003. Specialty rices of the world: effect of WTO and IRP on its production trend and marketing. *Journal of Food, Agriculture and Environment* **1(2)**: 34-41
- Chen, N., Luo, Y. K., Xie, L. H., Zhu, Z. W., Duan, B. W. and Zhang, L. P. 2006. Protein content and its correlation with other quality parameters of rice in China. *Acta Agronomic Sinica* **32(8)**: 1193-1196
- Cichoke, A. J. 1999. *The Complete Cook of Enzyme Therapy*. New York: Penguin Putnam Inc.
- Cuevas, R.P. and Fitzgerald, M. 2007. Linking Starch Structure to Rice Cooking Quality. *IREC Farmers' Newsletter*. No. 177, Summer 2007-2008. 16-17
- Danbaba, N., Anounye, J. C., Gana, A. S., Abo, M. E. and Ukwungwu, M. N. 2011. Grain quality characteristics of Ofada rice (*Oryza sativa L.*): cooking and eating quality. *International Food Research Journal* **18**: 619-624
- Debet, M. R. and Gidley, M. J. 2006. Three classes of starch granule swelling: Influence of surface proteins and lipids. *Carbohydrate Polymers* **64**: 452-465
- Deepa, G., Singh, V. and Naidu, K. A. 2008. Nutrient composition and physicochemical properties of Indian medicinal rice – Njavara. *Food Chemistry* **106(1)**: 165-171
- Delseny, M., Salsesa, J., Cookea, R., Sallaudb, C., Regadb, F., Lagodab, P., Guiderdonib, E., Ventelonc, M., Brugidouc, C. and Ghesquièrec, A. 2001. Rice genomic: present and future. *Plant physiology and Biochemistry* **39(3-4)**: 323-334
- Dipti, S. S., Bari, M. N. and Kabir, K. A. 2003. Grain quality characteristics of some Beruin rice varieties of Bangladesh. *Pakistan Journal of Nutrition* **2(4)**: 242-245

- El-Qudah, J., Dabadneh, B. F., Al-Bakheit, A. A., Al-Qudah, M. M., Al-Rawashdeh, A., Khataibeh, M., Abu-Alruz, K. and Ereifej, K. I. 2008. Nutrient contents per serving of twelve varieties of cooked rice marketed in Jordan. *American Journal of Agricultural and Biological Sciences* **3(3)**: 617-622
- Fageria, N. K., Baligar, V. C. and Clark, R. B. 2006. *Physiology of Crop Production*. New York: The Haworth press, Inc.
- FAO. 2004. *Rice is life*. International year of rice 2004 and its implementation. Rome, Italy: FAO
- Faruq, G., Hadjim, M. O. K. and Meisner, C. A. 2003. Kernel ageing: an analysis in four Malaysian rice varieties. *International Journal of Agriculture and Biology* **5(3)**: 230-232
- Fink, H. H., Burgoon, L. A. and Mikesky, A. E. 2008. *Practical Applications in Sports Nutrition*. 2nd edition. Sudbury, Canada: Jones and Barlett Publishers
- FND. 2002. Food and nutrition board, Institute of medicine. National Academy of Sciences. Dietary reference Intake for Energy, Carbohydrate, Fibre, Fat, Fatty Acids, Cholesterol, protein and Amino acid (micro-nutrients)
- Gariboldi, F. 1988. *Rice Milling Equipment Operation and Maintenance*. Philippines: FAO
- Gayin, J., Manful, J. T. and Johnson, P-N. T. 2009. Rheological and sensory properties of rice varieties from improvement programmes in Ghana. *International Food Research Journal* **16**: 167-174
- Golam, F., Mohamad, O., Hadzim, K. and Meisner, C. A. 2004. Inheritance of kernel elongation in rice. *International Journal of Agricultural Biology* **6(5)**: 813-815
- Golob, P., Hodges, R. and Farrell, G. 2004. *Crop Post-Harvest Science and Technology*. Volume II. UK: Balckwell Science Ltd.
- Graham, R. 2002. *A Proposal for IRRI to Establish a Grain Quality and Nutrition Research Center*. Philippines: IRRI
- Gujral, H. S. and Kumar, V. 2003. Effect of accelerated aging on the physicochemical and textural properties of brown and milled rice. *Journal of Food Engineering* **59**: 117-121
- Harold, McG. 2004. *On Food and Cooking: The Science and Lore of the Kitchen*. New York: Scribner
- Hassan, L. G. and Umar, K. J. 2006. Nutritional value of Balsam Apple (*Momordica balsamina L.*) leaves. *Pakistan Journal of Nutrition* **5(6)**: 522-529
- Hoeger, W. W. K. and Hoeger, S. A. 2010. *Lifetime Physical Fitness and Wellness: A Personalized Program*. 11th edition. USA: Wadsworth, Cengage Learning

- Hsieh, S. C. 1991. Studies on physicochemical aspect of eating quality and grading of rice. *Taichung DAIS* **31**: 1-11
- Hui, Y. H. and Corke, H. 2006. *Bakery products: Science and Technology*. USA: Blackwell Publishing
- Insel, P., Ross, D., McMahon, K. and Bernstein, M. 2010. *Nutrition*. 4th edition. Sudbury: Jones and Barlett Publishers
- Insel, P., Ross, D., McMahon, K. and Bernstein, M. 2010. *Nutrition*. Sudbury: Jones and Barlett Publishers
- International Rice Research Institute. 1991. *Rice Grain Marketing and Quality Issues*. Selected papers from the International Rice Research Conference Seoul, Korea. Philippines: IRRI
- Itani, T., Tamaki, M., Arai, E. and Horino, T. 2002. Distribution of amylose, nitrogen, and minerals in rice kernels with various characters. *Journal of Agriculture and Food Chemistry* **50**: 5326-5332
- Jane, J., Chen, Y. Y., Lee, L. F., McPherson, A., Wong, K. S. and Radosavljevic, M. 1999. Effects of amylopectin branch chain length and amylose content on the gelatinization and pasting properties of starch. *Cereal Chemistry* **76**: 629-637
- Jane, J., Kasemsuwan, T. and Chen, J. F. 1996. Phosphorus in rice and other starches. *Cereal Food World* **41**: 827-832
- Jeng, T. L., Wang, C. S., Chen, C. L. and Sung, J. M. 2003. Effects of grain position on the panicle on starch biosynthetic enzyme activity in developing grains of rice cultivar Tainung 67 and its NaN³-induced mutant. *Journal of Agricultural Science* **141**: 303-311
- Juliano, B. O. 1971. A simplified assay for milled rice amylose. *Cereal Science* **16**: 334-360
- Juliano, B. O. and Villareal, C. P. 1993. *Grain Quality Evaluation of World Rices*. Manila: IRRI
- Juliano, B. O. and Bechtel, D. B. 1985. *The Rice Grain and Its Gross Composition. Rice: Chemistry and Technology*. Minnesota: The American Association of cereal Chemists, Inc.
- Keeratipibul, S., Luangsakul, N. and Lertsatchayarn, T. 2008. The effect of Thai glutinous rice cultivars, grain length and cultivating locations on the quality of rice cracker (arare). *Food Science and Technology* **41**: 1934-1943
- Khan, A. G., Shabbir, G., Sadiq, M. and Hassan, G. 2003. Grain quality traits of candidate rice variety PB-95. *Asian Journal of Plant Sciences* **2(6)**: 483-484
- Khan, M. S. and Ali, C. A. 1985. Cooking quality of some rice varieties. *Journal of Agricultural Research Pakistan* **23(3)**: 231-233

- Khatoon, S. and Gopalakrishna, A. G. 2004. Fat-Soluble Nutraceuticals and Fatty Acid Composition of Selected Indian Rice Varieties. *Journal of American Oil Chemists' Society* **81(10)**: 939-943
- Khush, G. S. 1997. Origin, dispersal, cultivation and variation of rice. *Plant Molecular Biology* **35**: 25-34
- Khush, G. S., Singh, U. S. and Singh, R. K. 2000. *Aromatic Rices*. New Delhi: Oxford and IBH Publishing Co. Pvt. Ltd.
- Kibanda, J. M. N. and Luzi-Kihupi, A. 2007. Influence of genetic and genotype x environment interaction on quality of rice grain. *African Crop Science Journal* **15(4)**: 173-182
- King, T. L. and Brucker, M. C. 2009. *Pharmacology for Women's Health*. Sudbury: Jones and Barlett Publishers
- Lamberts, L., Bie, E. D., Vandeputte, G. E., Veraverbeke, W. S., Derycke, V., De Man, W. and Delcour, J. A. 2007. Effect of milling on colour and nutritional properties of rice. *Food Chemistry* **100(2)**: 1496-1503
- Lee, Y. E. and Osman, E. M. 1991. Correlation of morphological changes of rice starches granules with rheological properties during heating in excess water. *Journal of Korean Agricultural Chemical Society* **34**: 379-385
- Li, J. M., Thomson, M. and McCouch, S. R. 2004. Fine mapping of a grain-weight quantitative trait locus in the pericentromeric region of rice chromosome. *Genetics* **168**: 2187-2195
- Li, Y., Shoemaker, C. F., Ma, J. G., Moon, K. J. and Zhong, F. 2008. Structure-viscosity relationships for starches from different rice varieties during heating. *Food Chemistry* **106**: 1105-1112
- Little, R. R., Hilder, G. B. and Dawson, E. H. 1958. Differential effect of dilute alkali on 25 varieties of milled white rice. *Cereal Chemistry* **35**: 111-126
- Luh, B. S. 1991. *Rice Utilization*. 2nd edition. New York: Van Nostrand Reinhold
- Mohamad, O., Hadzim, K., Azlan, S., Abdullah, M. Z., Amiran, N., Nur Samahah, M. Z., Siti Noorfahana, M. I., Noraida, M. R., Cheng, A. and Salwa, A. S. 2008. Molecular screening for high quality rice. Plenary paper presented at the 6th National Genetics Congress, 12-14 May 2005, Kuala Lumpur
- Mohapatra, D. and Bal, S. 2006. Cooking quality and instrumental textural attributes of cooked rice for different milling fractions. *Journal of Food Engineering* **73**: 253-259
- Mustapha, N. A. 2007. Physicochemical and rheological characterization of Malaysian rice flours and starches. Master of Science Dissertation. Universiti Putra Malaysia

- Nag, A. 2006. *Analytical Techniques in Agriculture, Biotechnology and Environmental Engineering*. New Delhi: Prentice Hall of India Private Limited
- Nahar, K., Hasanuzzaman, M. and Majumder, R. R. 2009. Effect of low temperature stress in transplanted Aman Rice. *Academic Journal of Plant Sciences* **2(3)**: 132-138
- Najim, M. M. M., Lee, T. S., Haque, M. A. and Esham, M. 2007. Sustainability of rice production: a Malaysian perspective. *The Journal of Agricultural Sciences* **3(1)**: 1-12
- NIIR Board of Consultants and Engineers. 2006. *Wheat, Rice, Corn, Oat, Barley and Sorghum Processing Handbook (Cereal Food Technology)*. New Delhi: Asia Pacific Business Press Inc.
- Ntanos, D. A. and Koutroubas, S. D. 2002. Dry matter and N accumulation and translocation for Indica and Japonica rice under Mediterranean conditions. *Field Crops Research* **74**: 93-101
- Oko, A. O. and Ugwu, S. I. 2011. The proximate and mineral compositions of five major rice varieties in Abakaliki, South-Eastern Nigeria. *International Journal of Plant Physiology and Biochemistry* **3(2)**: 25-27
- Organisation for Economic Co-operation and Development (OECD). 2006. *Safety Assessment of Transgenic Organisms: OECD Consensus documents*. Vol 1. USA: OECD Publishing
- Otegbayo, B. O., Osamuel, F. and Fashakin, J. B. 2001. Effect of parboiling on physico-chemical qualities of two local rice varieties in Nigeria. *The Journal of Food Technology in Africa* **6(4)**: 130-132
- Owen, S. 2003. *The Rice Book*. London : Frances Lincoln Limited
- Ramesh, M., Ali, S. R. and Bhattacharya, K. R. 1999. Structure of rice starch and its relation to cooked-rice texture. *Carbohydrate Polymers* **38**: 337-347
- Randall, G., Mutters, G. and Thompson, J. F. 2009. *Rice Quality Handbook*. California: University of California
- Resurrection, A. P., Juliano, B. O. and Tanaka Y. 2006. Nutrient content and distribution in milling fractions of the rice grain. *Journal of the Science of Food and Agriculture* **30(5)**: 475-481
- Robert, A. R. 2003. *The Encyclopedia of Nutrition and Good Health*. New York: Facts on File, Inc.
- Robert, F. C. Jr. 1979. *Rice in The Tropics: A Guide to the Development of National Programs*. Colorado, USA: Westview Press
- Saleh, M. and Meullenet, J. F. 2007. Effect of long-grain rice degree of milling, moisture uptake, and solids leach during cooking on rice textural properties. *B.R. Wells Rice Research Studies*. 251-260

- Sareen, S., Gropper, S., Smith, J. L. and Groff, J. L. 2009. *Advanced Nutrition and Human Metabolism*. Canada: Wadsworth, Cengage Learning
- Shabbir, M. A., Anjum, F. M., Zahoor, T. and Nawaz, H. 2008. Mineral and pasting characterization of Indica rice varieties with different milling fractions. *International Journal of Agriculture Biology* **10**: 556-560
- Shils, M. E. and Shike, M. 2006. *Modern Nutrition in Health and Disease*. New York: Lippincott Williams and Wilkins
- Shu, X. L., Shen, S. Q., Bao, J. S., Wu, D. X., Nakamura, Y. and Shu, Q. Y. 2006. Molecular and biochemical analysis of the gelatinization temperature characteristics of rice (*Oryza sativa* L.) starch granules. *Journal of Cereal Science* **44**: 40-48
- Singh, N. and Sodhi, N. S. 2003. Morphological, thermal and rheological properties of starches separated from rice cultivars grown in India. *Food Chemistry* **80**: 99-108
- Singh, N., Kaur, L., Sandhu, K. S., Kaur, J. and Nishinari, K. 2006. Relationships between physicochemical, morphological, thermal, rheological properties of rice starches. *Food Hydrocolloids* **20**: 532-542
- Singh, N., Kaur, L., Sodhi, N. S. and Sekhon, K. S. 2005. Physicochemical, cooking and textural properties of milled rice from different Indian rice cultivars. *Food Chemistry* **89(2)**: 253-259
- Singh, N., Sodhi, N. S., Kaur, M. and Saxena, S. K. 2003. Physicochemical, morphological, thermal, cooking and textural properties of chalky and translucent rice kernels. *Food Chemistry* **82(3)**: 433-439
- Sivak, M. N. 1998. *Starch: Basic Science to Biotechnology*. USA: Elsevier Inc.
- Smith, C. W. and Dilday, R. H. 2003. *Rice: Origin, History, Technology, and Production*. New Jersey: John Wiley & Sons, Inc.
- Sompong, R., Siebenhandl-Ehn, S., Linsberger-Martin, G. and Berghofer, E. 2011. Physicochemical and antioxidative properties of red and black rice varieties from Thailand, China and Sri Lanka. *Food Chemistry* **124**: 132-140
- Soponronnarit, S., Chiawwet, M., Prachayawarakorn, S., Tungtrakul, P. and Taechapiroj, C. 2007. Comparative study of physicochemical properties of accelerated and naturally aged rice. *Journal of Food Engineering* **85**: 268-276
- Sotelo, A., Sousa, V., Montalvo, I., Hernandez, M. and Hernandez-Aragon, L. 1990. Chemical composition of different fractions of 12 Mexican varieties of rice obtained during milling. *Cereal Chemistry* **67(2)**: 209-212
- Sparks, D. L. 2004. *Advances in Agronomy*. Volume 82. California, USA: An Imrpint of Elsevier

- Srichuwong, S., Sunarti, T. C., Mishima, T., Isonoa, N. and Hisamatsua, M. 2005. Starches from different botanical sources II: Contribution of starch structure to swelling and pasting properties. *Carbohydrate Polymers* **62**: 25-34
- Sthapit, B. R., Upadhyay, M. P., Shrestha, P. K. and Jarvis, D. I. (eds.). 2004. On-Farm Conservation Agricultural Biodiversity in Nepal. Volume II. *Managing Diversity and Promoting its Benefits*. Rome, Italy: International Plant Genetic Resources Institute
- Sujatha, S. J., Ahmad, R. and Bhat, P. R. 2004. Physicochemical properties and cooking qualities of two varieties of raw and parboiled rice cultivated in the coastal region of Dakshina Kannada, India. *Food Chemistry* **86(2)**: 211-216
- Swain, E. W., Wang, H. L. and Hesseltine, C. W. 1977. Note on vitamins and minerals of wild rice. *Cereal Chemistry* **55(3)**: 412-414
- Tahir, H. A., Ahmad, M., Inayat, A., Muhammad, A. and Zaheen, M. 2007. Contribution of tillers within a rice plant to yield and yield components. *Journal of Agriculture Research* **45(3)**: 237-243
- Tang, H., Mitsunaga, T., and Kawamura, Y. 2005. Functionality of starch granules in milling fractions of normal wheat grain. *Carbohydrate Polymers* **59**: 11-17
- Tester, R. F. 1997. Starch: The polysaccharide fractions. In Frazier, R. J., Donald, A. M. and Richmond, P. (Eds.). *Starch Structure and Functionality*. Cambridge: Royal Society of Chemistry
- Tester, R. F. and Morrison, W. R. 1990. Swelling and gelatinization of cereal starches. I. Effect of amylopectin, amylose and lipids. *Cereal Chemistry* **67**: 551-557
- Toriyama, K., Heong, K. L. and Hardy, B. 2005. *Rice is Life: scientific perspectives for the 21st century*. Philippines: IRRI
- USDA. 2005. Rice quality categories. <http://www.ars.usda.gov/Research/docs.htm?docid=7060>. Access on 11 July 2005. Verified on 12 October 2010.
- USDA. 2009. Rough rice production by country and geographical region. beta.irri.org/solutions/.../wrs_jun09_2009_table01_usda_prod.xls. Access on 10 June 2009. Verified on 12 October 2010.
- Vaghefi, N., Shamsudin, M. N., Makmom, A. and Bagheri, M. 2011. The economic impacts of climate change on the rice production in Malaysia. *International Journal of Agricultural Research* **6**: 67-74
- Vanaja, T. and Babu, L. C. 2003. Variability in grain quality attributes of high yielding rice varieties (*Oryza sativa* L.) of diverse origin. *Journal of Tropical Agriculture* **44(1-2)**: 61-63
- Virmani, S. S. 1994. *Heterosis and Hybrid Rice Breeding*. Monographs on theoretical and applied genetics. Volume XXII. Berlin: Springer

- Wang, L. and Wang, Y. J. 2001. Comparison of protease digestion at neutral pH with alkaline steeping method for rice starch isolation. *Cereal Chemistry* **78**: 690-692
- Wang, L. and Wang, Y. J. 2004. Rice starch isolation by neutral protease and high intensity ultrasound. *Journal of Cereal Science* **39**: 291-296
- Wang, L., Xie, B. J., Shi, J., Xue, S., Deng, Q. C., Wei, Y. W. and Tian, B. Q. 2010. Physicochemical properties and structure of starches from Chinese rice cultivars. *Food Hydrocolloids* **24**: 208-216
- Watson, R. and Thelander, T. 2007. *The MediterrAsian Way*. New Jersey: John Wiley & Sons, Inc.
- Webb, F. S., Whitney, E., Sizer, F. S. and Whitney, E. N. 2008. *Nutrition: Concepts and Controversies*. Vol. 10. USA: Thomson Higher Education
- Williams, S. R. and Schlenker, E. D. 2003. *Essentials of Nutrition and Diet Therapy*. Vol 1. USA: AN Affiliate of Elsevier Science
- Yadav, B. K. and Jindal, V. K. 2007. Water uptake and solid loss during cooking of milled rice (*Oryza sativa L.*) in relation to its physicochemical properties. *Journal of Food Engineering* **80(1)**: 46-54
- Yadav, R. B., Khatkar, B. S. and Yadav, B. S. 2007. Morphological, physicochemical and cooking properties of some Indian rice (*Oryza sativa L.*) cultivars. *Journal of Agricultural Technology* **3(2)**: 203-210
- Yoroshimura, M., Takahashi, H. and Nakanishi, T. 1991. Role of sodium, potassium , calcium, magnesium on blood pressure regulation and antihypertensive dietary therapy. *Japanese Journal of Nutrition* **49**: 53-62
- Yu, S. F., Ma, Y., Menager, L. and Sun, D. W. 2010. Physicochemical properties of starch and flour from different rice cultivars. *Food Bioprocessing Technology*. Springer
- Zhong, F., Lia, Y., Ibanez, M. A., Oh, M. H., McKenzie, K. S. and Shoemaker, C. 2006. The effect of rice variety and starch isolation method on the pasting and rheological properties of rice starch pastes. *Food Hydrocolloids* **23**: 406-414