

Growth and biochemical composition of *Kappaphycus* (Rhodophyta) in customized tank culture system

Abstract

The study was conducted to determine the growth and biochemical composition of *Kappaphycus* cultivated in a customized tank culture system. Two red seaweed species (*Kappaphycus alvarezii* and *Kappaphycus striatum*) were selected and cultivated using suspension culture method in the tank. Three cycles of 40-day culture trials were performed during September to December 2014, and both *K. alvarezii* and *K. striatum* were successfully grown in the tank. This is the first report on the success of seaweed culture in Malaysia involving land-based facility. Interestingly, *K. striatum* was found to grow better than *K. alvarezii* in the tank. The daily growth rate (DGR) and daily weight productivity (DWP) of *K. alvarezii* ranged from 1.96 ± 0.08 to 2.29 ± 0.11 % day⁻¹ and 3.70 ± 0.20 to 4.55 ± 0.34 g DW m⁻² day⁻¹, and those of *K. striatum* ranged from 2.25 ± 0.06 to 2.96 ± 0.02 % day⁻¹ and 4.48 ± 0.19 to 6.17 ± 0.18 g DW m⁻² day⁻¹, respectively. These values were influenced by the changes in the water quality variables during the culture period. On the other hand, the biochemical composition of *K. alvarezii* and *K. striatum* was not significantly different ($p > 0.05$) from each other. Both growth and biochemical composition of *K. alvarezii* and *K. striatum* in the present study were comparable with those cultured in the open sea. In conclusion, the findings indicate the ability of *Kappaphycus* to grow well in land-based cultivation system which can be further explored to support the development of local seaweed farming industry especially for the high-quality seed production.