

**PHYTOREMOVAL AND PHYTOTOXICITY OF
TYPHA ANGUSTIFOLIA L. EXPOSED TO
NORETHINDRONE UNDER HYDROPONIC
CULTURE**

LEE JIE YINN

**THESIS SUBMITTED IN FULFILMENT FOR
THE DEGREE OF MASTER OF SCIENCE**

**PERPUSTAKAAN
UNIVERSITI MALAYSIA SABAH**

**INSTITUTE FOR TROPICAL BIOLOGY AND
CONSERVATION**

UNIVERSITY OF MALAYSIA SABAH

2019



UMS
UNIVERSITI MALAYSIA SABAH

UNIVERSITI MALAYSIA SABAH
BORANG PENGESAHAN STATUS TESIS

JUDUL: **PHYTOREMOVAL AND PHYTOTOXICITY OF TYPHA ANGUSTIFOLIA L. EXPOSED TO NORETHINDRONE UNDER HYDROPONIC CULTURE**
IJAZAH: **SARJANA SAINS (PROSES EKOLOGI)**

Saya **LEE JIE YINN**, Sesi **2016-2019**, mengaku membenarkan tesis Sarjana ini disimpan di Perpustakaan Universiti Malaysia Sabah dengan syarat-syarat kegunaan seperti berikut:-

1. Tesis ini adalah hak milik Universiti Malaysia Sabah.
2. Perpustakaan Universiti Malaysia Sabah dibenarkan membuat salinan untuk tujuan pengajian sahaja.
3. Perpustakaan dibenarkan membuat salinan tesis ini sebagai bahan pertukaran antara institusi pengajian tinggi.
4. Sila tandakan (/)

SULIT

(Mengandungi maklumat yang berdarjah keselamatan atau kepentingan Malaysia seperti yang termaktub di dalam AKTA RAHSIA 1972)

TERHAD

(Mengandungi maklumat TERHAD yang telah ditentukan oleh organisasi/badan di mana penyelidikan dijalankan)

TIDAK TERHAD



LEE JIE YINN
MX1611014T

PERPUSTAKAAN
UNIVERSITI MALAYSIA SABAH



Disahkan Oleh,

NORAZLYNNE MOHD. JOHAN @ JAC'LYNE
PUSTAKAWAN
UNIVERSITI MALAYSIA SABAH

(Tandatangan Pustakawan)


PROF. MADYA DR. KARTINI SAIBEH
PENSYARAH D554
FAKULTI PERTANIAN LESTARI
UMS SANDAKAN



Tarikh: 19 Februari 2019

Penyelia utama
(PROF MADYA DR KARTINI SAIBEH)

Penyelia bersama
(PROF DR CHARLES S. VAIRAPPAN)
Institute for Tropical Agriculture and Forestry
Universiti Malaysia Sabah, Kota Kinabalu,
Sabah, 88400, Malaysia




UMS
UNIVERSITI MALAYSIA SABAH

DECLARATION

I hereby declare that the material in this thesis is my own except for the quotations, excerpts, equations, summaries and references, which have been duly acknowledged.

25 January 2019



LEE JIE YINN
MX1611014T

CERTIFICATION

NAME : LEE JIE YINN
MATRICE NO : MX1611014T
TITLE : PHYTOREMOVAL AND PHYTOTOXICITY OF *TYPHA ANGUSTIFOLIA* L. EXPOSED TO NORETHINDRONE UNDER HYDROPONIC CULTURE
DEGREE : MASTER OF SCIENCE (ECOLOGICAL PROCESSES)
VIVA DATE : 25 JANUARY 2019

CERTIFIED BY;

SIGNATURE

1. MAIN SUPERVISOR

Assoc. Prof. Dr. Kartini Binti Saibeh

PROF. MADYA DR. KARTINI SAIBEH
PENYARAH DSS4
FAKULTI PERTANIAN LESTARI
UMS SANDAKAN



2. CO-SUPERVISOR

Pror. Dr. Charles S. Vairappan



ACKNOWLEDGEMENT

It is gratefully acknowledged that this research was financially supported by the grant from Universiti Malaysia Sabah, UMSGreat- GUG0054-2/2016. I also would like to thank Universiti Malaysia Sabah true and for the financial support by Teaching Scheme Scholarship so that I can finish my study without any worries.

Besides, I express my deepest gratitude and appreciation to my supervisors, Associate Professor Dr Kartini Saibeh and Professor Dr Charles S. Vairappan for their advice, guidance and supports during the entire research study. They have helped me from every perspective, in field work, in identification, in writing, in presenting myself, and thank you for giving me the confidence. Besides, I also appreciate for all field and laboratory assistance from Institute for Tropical Biology and Conservation, Faculty of Sustainable Agriculture and Faculty of Science and Natural Resources. I also want to send my thanks to my friends in the institute too for the accompaniment through the hard times.

Finally, I would like to express my profound gratitude to my family for providing me with unfailing support and continuous encouragement throughout my years of study and through the process of researching and writing this thesis. This accomplishment would not have been possible without them. Thank you.

LEE JIE YINN
25 January 2019



ABSTRACT

This study was conducted to assess the potential of *Typha angustifolia* L. to tolerate and remove norethindrone. The research objectives were (i) to investigate the phytoremoval efficiency of *T. angustifolia* when exposed to norethindrone; and (ii) to assess the phytotoxicity of norethindrone by *T. angustifolia* through phytoremediation. *Typha angustifolia* were collected and acclimatised under hydroponic culture. Young seedlings were harvest and transferred to 0 – 2.0 mg/L of norethindrone for a maximum period of 28 days. Norethindrone in water was analysed using dispersive liquid-liquid microextraction with the solidification of floating organic (DLLME-SFO) and followed with the chromatography analysis by HPLC-UV/Vis. Meanwhile, the plant samples were separated into three parts (root, stem and leaves), air-dried in room temperature and then followed with sea sand matrix disrupting extraction (SSDM), cleaned up with preparative thin layer chromatography (PTLC) and finally analysed by HPLC-UV/Vis. Calculations like BCF, Ti, TF, and RGR were applied, followed with statistical analysis with ANOVA and correlation test. An evaluation showed that norethindrone removal efficiency by plants reached a value of 90% from the initial contents at 28 days. The rate of removal kinetic followed the first order reaction rate with rate constant of 0.5mg/L treatment (-0.131 h^{-1}) > 1.0 mg/L (-0.02 h^{-1}) > 2.0 mg/L (-0.003 h^{-1}). The relative growth rate of plant weight and height showed that decreasing order from 0.5 mg/L > 1.0 mg/L > 2.0 mg/L norethindrone exposure treatment. The plants exposed to norethindrone did not experience chlorosis effect on root and necrosis effect on leaves. Norethindrone accumulation was higher in the root (0.012-0.07 mg/g) than stem (0.018-0.049 mg/g) and leaves (0.006-0.041 mg/g). The plants in (i) 1.0 mg/L norethindrone treatment and (ii) 2.0 mg/L norethindrone treatment have similar increasing BCF order: $\text{BCF}_{\text{leaves}}$ (0.016; 0.017) < BCF_{stem} (0.019;0.02) < BCF_{root} (0.023;0.022), except for the plants in 0.5 mg/L norethindrone treatment with the increasing order BCF_{root} (0.006) < BCF_{stem} (0.026) < $\text{BCF}_{\text{leaves}}$ (0.027). The root concentration factor was higher in 0.5 mg/L treatment (25.829) than 1.0 mg/L (21.005) and 2.0 mg/L (21.495). The translocation factor (TF) values and tolerance index (Ti) values were >1 but the values were decreasing when exposure of norethindrone concentration increasing. Although the removal performance was greater in 0.5 mg/L norethindrone treatment, *T. angustifolia* still has performed its ability in the removal of norethindrone from aqueous medium. It also proves that *T. angustifolia* able to tolerate the toxicity of norethindrone even they are exposed to high concentration.



ABSTRAK

Fito-penyngkiran dan Fito-ketoksikan *Typha angustifolia* L. Terdedah kepada Norethindrone secara Kultur Hidroponik

*Kajian ini dijalankan untuk menilai potensi *Typha angustifolia* L. untuk menyingkir dan bertoleransi norethindrone. Objectif penyelidikan ini adalah (i) mengkaji kecekapan *T. angustifolia* untuk menyingkir norethindrone; (ii) mengkaji penyesuaian *T. angustifolia* terhadap norethindrone melalui fito-remediasi. *T. angustifolia* diambil dan ditumbuh secara hidroponik. Anak tumbuhan telah dituai dan dipindahkan ke 0-2.0 mg/L rawatan norethindrone untuk tempoh maksimum 28 hari. Norethindrone dalam air dianalisa dengan menggunakan dispersive liquid-liquid microextraction with solid floating organic (DLLME-SFO) dan diikuti dengan HPLC-UV/VIS. Sementara itu, sampel tumbuhan dibahagi kepada tiga bahagian (akar, batang dan daun), kering dalam suhu bilik dan kemudian diikuti dengan sea sand matrix disrupting extraction (SSDM), dibersihkan dengan kromatografi lapisan nipis (PTLC) dan akhirnya dianalisis oleh HPLC-UV/VIS. Pengiraan seperti BCF, Ti, TF, dan RGR digunakan, diikuti dengan analisis statistik dengan ujian ANOVA dan korelasi. Penilaian menunjukkan bahawa kecekapan penyingkiran norethindrone oleh tumbuhan telah mencapai nilai 90% dari kandungan pada 28 hari. Kadar kinetik penyingkiran mengikut kadar tindak balas pesanan pertama dengan kadar tetap rawatan 0.5mg/L (-0.131 h^{-1}) > 1.0 mg/L (-0.02 h^{-1}) > 2.0 mg/L (-0.003 h^{-1}). Kadar pertumbuhan relatif berat dan ketinggian tumbuhan menurun daripada rawatan norethindrone 0.5 mg/L > 1.0 mg/L > 2.0 mg/L. Tanaman yang terdedah kepada norethindrone tidak mengalami kesan klorosis pada kesan akar dan nekrosis pada daun. Kepekatan norethindrone didapati lebih tinggi pada akar (0.012-0.07 mg/g), batang (0.018-0.049 mg/g) dan daun (0.0061-0.041 mg/g). Tumbuhan dalam rawatan 1.0mg/L norethindrone dan rawatan 2,0 mg/L norethindrone mempunyai sama susunan BCF iaitu $BCF_{\text{daun}} (0.016; 0.017) < BCF_{\text{batang}} (0.019; 0.02) < BCF_{\text{akar}} (0.023; 0.022)$ tetapi tumbuhan dalam rawatan 0.5 mg/L norethindrone pula dengan peningkatan $BCF_{\text{akar}} (0.006) < BCF_{\text{batang}} (0.026) < BCF_{\text{daun}} (0.027)$. Manakala, faktor kepekatan akar didapati lebih tinggi dalam rawatan 0.5 mg/L(25.8289) daripada 1.0 mg/L(21.0054) dan 2.0 mg/L(21.4953). Faktor tumpuan akar (RCF) lebih tinggi dalam rawatan 0.5 mg/L (25.829) daripada 1.0 mg/L (21.005) dan 2.0 mg/L (21.495). Nilai translocation factor (TF) dan indeks toleransi (Ti) adalah > 1 tetapi nilai-nilai itu berkurangan apabila pendedahan kepekatan norethindrone meningkat. Sebagai kesimpulan, Walaupun prestasi penyingkiran lebih besar dalam rawatan 0,5 mg/L norethindrone, *T. angustifolia* berpotensi untuk menyingkirkan norethindrone dalam air. Ia juga membuktikan bahawa *T. angustifolia* dapat bertoleransi dengan kesan ketoksikan norethindrone walaupun mereka terdedah kepada kepekatan yang tinggi.*

LIST OF CONTENT

	Pages
TITLE	i
DECLARATION	ii
CERTIFICATION	iii
ACKNOWLEDGEMENT	iv
ABSTRACT	v
<i>ABSTRAK</i>	vi
LIST OF CONTENT	vii
LIST OF FIGURES	ix
LIST OF TABLES	xii
LIST OF EQUATIONS	xiii
LIST OF ABBREVIATIONS	xiv
LIST OF APPENDICES	xv
CHAPTER 1 INTRODUCTION	
1.1 Objectives	4
1.2 Thesis contribution	4
1.3 Scopes of study	5
CHAPTER 2 LITERATURE REVIEWS	
2.1 Pharmaceutical pollution	6
2.2 Synthetic steroid and hormonally pharmaceutical residues act as emerging micro-pollutants in the environment.	7
2.3 Norethindrone	9
2.3.1 Chemical characteristic of norethindrone	9
2.3.2 Quantification of norethindrone in various matrix	12
2.3.3 Distribution of norethindrone in environment	13
2.3.4 Toxicological effects of norethindrone on microbials and animals	17
2.4 Phytoremediation	20
2.5 <i>Typha angustifolia</i>	22



CHAPTER 3	METHODOLOGY	
3.1	Chemical reagents and materials	27
3.2	Hydroponic culture vessels set up	28
3.3	Plants collection and acclimatisation	29
3.4	Phytoremoval experiment set up	32
3.5	Phytotoxicity experiment set up	33
3.6	Quantification of norethindrone in nutrient solution	34
3.7	Quantification of norethindrone in plants	35
3.8	Quality assurance and quality control	36
3.9	Statistical analysis on norethindrone uptake evaluation	39
CHAPTER 4	RESULTS AND DISCUSSION	
4.1	Phytoremoval efficiency of <i>Typha angustifolia</i>	42
	4.1.1 Comparison: plants and without plants	42
	4.1.2 Comparison: among different initial concentration	44
4.2	Phytotoxicity investigation of plants on norethindrone	48
	4.2.1 Physical and morphological changes of plant in height, weight, leaves and roots	48
	4.2.2 Accumulation of norethindrone in plants tissues	58
CHAPTER 5	CONCLUSION	74
	REFERENCES	76
	APPENDICES	96

LIST OF FIGURES

		Pages
Figure 2.1	Fate of Pharmaceutical Residues from Human to Environment	8
Figure 2.2	Norethindrone is Structurally Related to the Natural Hormonal Steroids: Progesterone and Testosterone.	9
Figure 2.3	Structural relationships of the 19-nortestosterone derivatives. All compounds were derived from testosterone. The compounds with similar structures relate to the dotted arrows and were differentiated by functional groups	10
Figure 2.4	The Inner Box Shows that Synthesis of Norethindrone from Testosterone. Meanwhile, Metabolism Pathway of Norethindrone Also Schemes with Its Metabolites. However, the Substitutions At 7 α - And 11- Position May Inhibit the Metabolism To 5 α -Reduced Forms. It Demonstrates that 5 α -Reduction is the Rate-Limiting Step in Formation of 3 α -OH and 5 α -H-Reduced Metabolites	12
Figure 2.5	Residual Norethindrone Over Time (Days) in Surface Water and Wastewater Measured At 4 °C and 21.5 °C with an Initial Spiked Concentration 400 μ g/L.	14
Figure 2.6	The World Map Illustrates the Summary of Studies on the Application of <i>Typha</i> Spp. in Various Countries.	25
Figure 3.1	A Plastic Vessel with Dimension Length 520.0 mm, Width 320 mm, Height 212.0 mm was Selected to be the Hydroponic Culture Vessel. A Polyester Broad was Cut and Designed as Cover. This Figure was Sketched with Coreldraw X6® Under the Scale 1:8.	27
Figure 3.2	Wild <i>T. Angustifolia</i> Grew in Rural Area, Located at Mile 10, Sandakan, Sabah, in which this Area is Far Away from the City of Sandakan.	28
Figure 3.3	Matured <i>T. angustifolia</i> Has Height 1.75-2.0 m.	29
Figure 3.4	Matured <i>T. Angustifolia</i> was Collected in Mile 10, Sandakan, Sabah. This Area is Located Nearly to Forest and Swamp, in which It is Almost Free from Source Point of Pollution Like Industry and Urban Area.	30
Figure 3.5	Phytoremediation Experiment in Closed Environment with 12 Hours Daylight (1500 Lux), Temperature Maintained at around 24-27 °C and Humidity Maintained at around 65-75%.	31
Figure 3.6	Flow Chart of the Methodology in Quantification of Norethindrone from Water	36

Figure 3.7	Quantification of Norethindrone in Plant Tissues by Modified Sea Sand Dispersive Extraction Method	37
Figure 4.1	(A) The Percentage of Remaining Norethindrone in the Water and (B) The Percentage of Norethindrone Removal from Water as Function of the Exposure Time, for Initial Concentration 1.0 Mg/L and the Control without Plants (N=9). Error Bars Represent the Standard Deviations of the Trend (Two-Way ANOVA, $F_{[6,112]} = 218.735$, $P < 0.01$).	42
Figure 4.2	The Percentage of Norethindrone Removal from Water as Function of the Exposure Time, for Initial Concentration 0.5 mg/L, 1.0 mg/L and 2.0 mg/L.	43
Figure 4.3	Kinetics of Norethindrone Removal from Hydroponic Solution by <i>T. angustifolia</i> from the Treatments with Initial Concentration of 0.5 mg/L, 1.0 mg/L, 2.0 mg/L within 28 Days (672 hours).	46
Figure 4.4	The Plant Heights in Respective Treatments to Exposure Time with Respectively Error Bars as Their Standard Deviation.	48
Figure 4.5	Relative Growth Rate-Weight (RGR-W) as Function with Natural Logarithm of Differences between Weights against Exposure Time. (n=9; ANOVA; $F_{[9, 128]} = 14.964$, $p < 0.01$).	49
Figure 4.6	The Root of Plants Has Less or No Influenced by Norethindrone Effect when the Plants Were Exposed to Different Norethindrone Content Treatments (0 mg/L Treatment).	51
Figure 4.7	Chlorosis Effect on the Plant Leaves Was Shown Less Significantly in 0.5 mg/L Norethindrone Treatment. The Tip Part of Leaves Shown Lack of Green Pigment. The Colour of Leaves and Some of Tip Parts Were Remained when Compared to Plants in 2.0 mg/L Norethindrone Treatment.	52
Figure 4.8	Chlorosis Effect on the Plant Leaves Was Shown Moderate Significantly in 1.0 mg/L Norethindrone Treatment. The Tip Part of Leaves Shown Lack of Green Pigment. The Colour of Leaves (Mid Part) Was Remained when Compared to Plants in 2.0 mg/L Norethindrone Treatment.	53
Figure 4.9	Chlorosis Effect on Plant Leaves Was Shown Significantly in 2.0 mg/L Norethindrone Treatment. The Tip Part of Leaves Shown Lack of Green Pigment. The Colour of Leaves also Shown Yellowish Compared to Other Treatment.	54
Figure 4.10	Mean Concentration of Norethindrone Found in Different Parts of Plant (Stem: Concentration in Stem; Root: Concentration in Root; Leaves: Concentration Found in Leaves) after Harvesting on 28 th Day.	58

Figure 3.7	Quantification of Norethindrone in Plant Tissues by Modified Sea Sand Dispersive Extraction Method	37
Figure 4.1	(A) The Percentage of Remaining Norethindrone in the Water and (B) The Percentage of Norethindrone Removal from Water as Function of the Exposure Time, for Initial Concentration 1.0 Mg/L and the Control without Plants (N=9). Error Bars Represent the Standard Deviations of the Trend (Two-Way ANOVA, $F_{[6,112]} = 218.735$, $P < 0.01$).	42
Figure 4.2	The Percentage of Norethindrone Removal from Water as Function of the Exposure Time, for Initial Concentration 0.5 mg/L, 1.0 mg/L and 2.0 mg/L.	43
Figure 4.3	Kinetics of Norethindrone Removal from Hydroponic Solution by <i>T. angustifolia</i> from the Treatments with Initial Concentration of 0.5 mg/L, 1.0 mg/L, 2.0 mg/L within 28 Days (672 hours).	46
Figure 4.4	The Plant Heights in Respective Treatments to Exposure Time with Respectively Error Bars as Their Standard Deviation.	48
Figure 4.5	Relative Growth Rate-Weight (RGR-W) as Function with Natural Logarithm of Differences between Weights against Exposure Time. ($n=9$; ANOVA; $F_{[9, 128]} = 14.964$, $p < 0.01$).	49
Figure 4.6	The Root of Plants Has Less or No Influenced by Norethindrone Effect when the Plants Were Exposed to Different Norethindrone Content Treatments (0 mg/L Treatment).	51
Figure 4.7	Chlorosis Effect on the Plant Leaves Was Shown Less Significantly in 0.5 mg/L Norethindrone Treatment. The Tip Part of Leaves Shown Lack of Green Pigment. The Colour of Leaves and Some of Tip Parts Were Remained when Compared to Plants in 2.0 mg/L Norethindrone Treatment.	52
Figure 4.8	Chlorosis Effect on the Plant Leaves Was Shown Moderate Significantly in 1.0 mg/L Norethindrone Treatment. The Tip Part of Leaves Shown Lack of Green Pigment. The Colour of Leaves (Mid Part) Was Remained when Compared to Plants in 2.0 mg/L Norethindrone Treatment.	53
Figure 4.9	Chlorosis Effect on Plant Leaves Was Shown Significantly in 2.0 mg/L Norethindrone Treatment. The Tip Part of Leaves Shown Lack of Green Pigment. The Colour of Leaves also Shown Yellowish Compared to Other Treatment.	54
Figure 4.10	Mean Concentration of Norethindrone Found in Different Parts of Plant (Stem: Concentration in Stem; Root: Concentration in Root; Leaves: Concentration Found in Leaves) after Harvesting on 28 th Day.	58

Figure 4.11	Correlation Matrix on Relationship between Different Concentration Treatments and Concentration Found in Different Parts of Plant Is Performed in Correlogram. The Significance is Displayed with Symbol "***" Represent p-Value with $p < 0.05$.	59
Figure 4.12	Estimated Bioconcentration Factor (BCF) (Fresh Weight, mg/g) of Norethindrone in Root Part and Shoot Part of <i>T. angustifolia</i> . Error Bar Represents Standard Deviation With $n=9$ per Treatment ($F_{[4,72]} = 171.50, p < 0.01$).	60
Figure 4.13	Translocation Factor (TF) in <i>T. angustifolia</i> exposed for 28 Days to Different Norethindrone Concentrations (Mean Data \pm S.D.).	64
Figure 4.14	Tolerance Index (TI) in <i>T. angustifolia</i> exposed for 28 Days to Different Norethindrone Concentrations (Mean Data \pm S.D.).	66
Figure 4.15	Correlation between $\text{Log BCF}_{\text{root}}$ and Log K_{ow} for Norethindrone Measured in <i>T. angustifolia</i> Grown in Hydroponic Culture with Concentration of 0.5, 1.0 and 2.0 mg/L	69
Figure 4.16	Correlation between $\text{Log BCF}_{\text{root}}$ and Log D_{ow} for Norethindrone Measured in <i>T. angustifolia</i> Grown in Hydroponic Culture with Concentration of 0.5, 1.0 and 2.0 mg/L	70
Figure 4.17	Positive Linear Relationship Showed in between Root Concentration Factor (Log RCF) and Octanol-Water Partition Coefficient (Log K_{ow}).	70
Figure 4.18	Positive Linear Relationship Showed in between Root Concentration Factor (Log RCF) and Octanol-Water Partition Coefficient (Log D_{ow}).	71
Figure 4.19	Negatively Linear Correlation Found in between Log TF and Log K_{ow} .	72

LIST OF TABLES

		Pages
Table 2.1	The data shows various water solubility of norethindrone	10
Table 2.2	Occurrence of norethindrone in different environment matrix.	15
Table 2.3	Application of phytoremediation with various hydrophytes and terrestrial macrophytes in eliminating various types of pollutants.	20
Table 2.4	Application of Typhaceae in phytoremediation that have been conducted to improve the knowledge of constructed wetland.	23
Table 4.1	The percentage of norethindrone removal from water was calculated by determining the remaining final concentration in water according to respectively treatments with concentration 0.5 mg/L, 1.0 mg/L, and 2.0 mg/L.	45
Table 4.2	The relative growth rate of plants height (RGR-H) within 28 days in the respective treatments.	48
Table 4.3	The concentrations of norethindrone (mg/g) found in the plants in different parts were analysed with one-way ANOVA to test the significance differences of the concentration found in the parts of plants according to respectively treatment.	59
Table 4.4	The estimated bioconcentration factor (BCF) (fresh weight, mg/g) of norethindrone in root part and shoot part of <i>T. angustifolia</i> .	62
Table 4.5	The estimated root concentration factor (RCF) of norethindrone in <i>Typha angustifolia</i> .	63
Table 4.6	Translocation factor (TF) of <i>Typha angustifolia</i> after exposure to norethindrone was analysed with one-way ANOVA	64
Table 4.7	The result of tolerance index (Ti) of <i>Typha angustifolia</i> after exposure to norethindrone were obtained and analysed with one-way ANOVA according to Tukey's test.	65

LIST OF EQUATIONS

	Pages
Equation 3.1 Relative growth rate- Weight (RGR-W)	38
Equation 3.2 Relative growth rate- Height (RGR-H)	38
Equation 3.3 Translocation factor (TF)	39
Equation 3.4 Tolerance index (TI)	39
Equation 3.5 Bioconcentration factor (BCF)	39
Equation 3.6 Root concentration factor (RCF)	40
Equation 4.1 The function of the norethindrone removal rate within 21 days	42
Equation 4.2 Reaction kinetics of norethindrone removal in the 0.5 mg/L treatment	45
Equation 4.3 Reaction kinetics of norethindrone removal in the 1.0 mg/L treatment	45
Equation 4.4 Reaction kinetics of norethindrone removal in the 2.0 mg/L treatment	46



LIST OF ABBREVIATIONS

%	- percent
ANOVA	- analysis of variance
CHCl ₃	- chloroform
<i>et al.</i>	- and others; latin phrase <i>exempli gratia</i>
g	- gram
GC	- gas chromatography
h	- hour(s)
HPLC-UV/VIS	- high performance liquid chromatography coupled with ultraviolet-visible spectroscopy detector
MeCN	- acetonitrile
MeOH	- methanol
mg	- milligram
mL	- millilitre
MS	- mass spectrometry
μL	- microlitre
n	- number of samples
ng	- nanogram
p	- significant different value
R	- regression
spp.	- species
<i>T. angustifolia</i>	- <i>Typha angustifolia</i>
TLC	- thin layer chromatography

LIST OF APPENDICES

	Pages
Appendices A The study of occurrence (is taken within 2006-2017) and ecotoxicological data on organisms are presented with corresponding references	96
Appendices B Analytical methodologies were developed by various researchers to quantify norethindrone in different matrices	100
Appendices C Quantification of norethindrone with optimised DLLME-SFO	106
Appendices D Quantification of norethindrone with optimised SSDM	111
Appendices E Raw data	112
Appendices F Correlation analysis	120
Appendices G R script	123
Appendices H Chromatography (HPLC and TLC)	126



CHAPTER 1

INTRODUCTION

Phytoremediation is a plant-based bioremediation technique approach in the detoxification or removing pollutants from soil and water environment. Since it is low cost, sustainable, and environmentally sound technology, application of this technique includes passive treatment using vegetation barriers, cleaning of soil and water through pollutant uptake and harvest, and pollutant degradation and volatilization provides a valuable component of integrated solutions for inorganic and organic pollution. Previous research on the plant uptake of pharmaceuticals and steroids, demonstrated their potential to accumulate pollutants (Boxall *et al.* 2006; Dordio *et al.* 2010; Seo *et al.* 2010; Dodgen *et al.* 2013) hydroponically or in nutrient solution has also been reported (Redshaw *et al.* 2008; Herklotz *et al.* 2010). Plants used in phytoremediation are usually evaluated with phytoremoval efficiency and phytotoxicity. Phytoremoval efficiency is evaluated by assessing the ability of plant in removing a compound from water, soil or soil-water in given exposure duration under controllable condition (Dordio *et al.* 2009b). Meanwhile, phytotoxicity or toxicity effect study of a compound to plant is used to determine the sensitivity of plant to xenobiotic action and occurrence of that compound in certain concentration and area (Carvalho *et al.* 2014; Bartikova *et al.* 2016). These two components are important to determine the potential and availability of plants used in phytoremediation.



Typhaceae such as *Typha agustifolia* L. is one of the frequent emergent plants and widely used in a constructed wetland. Various studies have reported Typhaceae ability to withstand and remove pharmaceutical and other trace emerging compounds such as clofibrac acid (Dordio *et al.*, 2009b), ibuprofen (Dordio *et al.* 2011a; Li *et al.* 2016), and carbamazepine (Dordio *et al.* 2011b) from water. However, it is interesting that there is lack of research in the application of *Typha* spp. for the accumulation of hormonal pharmaceutical compounds.

Steroids including estrogen, progestin and androgen pose ecotoxicity risk to environment. Norethindrone is a synthetic progestin derived from 19-nortestosterone that is used medicinally in the treatment of oral contraceptive with estrogen, hormone replacement therapy as well as menstrual disorders and endometriosis. However, compared to estrogens such as ethinylestradiol, estradiol and estrone, the presence of progesterone and progestin are not frequently reported (Fent 2015). Various studies believed that progestin like norethindrone can be converted into estrogenic compounds in animals' metabolism before discharged from body into sewage and wastewater (Kamyab *et al.* 1968; Mayberry *et al.* 1990; Kuhn *et al.* 1997). Therefore, many studies were focusing on the presence of the estrogenic steroid hormones in environment rather than progestogenic steroid hormones.

But numerous researchers have reported the presence of norethindrone in water even after the water treatments. The concentration have been found in surface water and wastewater from 0.2 ng/L up to 0.8 mg/L in several research such as in Labadie and Budzinski (2005), Vulliet *et al.* (2009), Vulliet and Cren-Olive (2011), and Al-Odaini *et al.* (2012, 2013). Presence of norethindrone in the aquatic environment has recognised norethindrone as an environmental pollutant due to poor efficiency of sewage and wastewater treatment plants. Cormier *et al.* (2015) reported that norethindrone is a non-natural degradable compound. It can persist in water for a long period at normal temperature. In another word, it can pose a risk to environment. Studies by Wibbels and Crews (1995), Maier and Herman (2001), Goto and Hiromi (2003), Gomez *et al.* (2010), Paulos *et al.* (2010), Zucchi *et al.* (2012), Safholm *et al.* (2014) and Wu *et al.* (2016b) have indicated that norethindrone has negative impacts likes decreasing fecundity, leading to population declines and affecting sex determination in aquatic organisms in/or extremely low concentration.

The current sewage treatment plants are not designed to effectively eliminate non-conventional toxic compounds like norethindrone in water. Generally, steroidal hormones have biodegradability, resulting in partial removal from the water phase and less deposited in the sedimentation and biological treatment in sewage and wastewater treatment plants (Ternes *et al.* 2004; Bendz *et al.* 2005; Heidler *et al.* 2006). A study showed that although the occurrence of pharmaceutical compounds in hospital wastewater is consistently higher than urban wastewater, the application of conventional treatments is still less efficiency to eliminate pharmaceutical compounds (Verlicchi *et al.* 2010).

The removal of norethindrone requires advanced technology such as advanced oxidation (Gultekin and Ince 2007; Broseus *et al.* 2009; Fayad *et al.* 2013), chemically removal (Zhang *et al.* 2015; Badejo *et al.* 2017), nano-/ultrafiltration (Verliefde *et al.* 2007; Yoon *et al.* 2007), microbial-based bioremediation (Passatore *et al.* 2014; Rana *et al.* 2014; Gavrilesco *et al.* 2015) and phytoremediation. However, most of the methods are highly cost and non-environmental friendly except for bioremediation and phytoremediation (Abd Manan *et al.* 2015). Most conventional wastewater treatment plants do not have treatment processes such as activated carbon, ozonation or membrane treatment. Therefore, these emerging micro-pollutants are not removed but are easily released to the receiving natural waters (Bolong *et al.* 2009). As a result, considerable amounts of pollutants remain in sewage treatment effluents and sludge. The discharge of the effluents will lead to the contamination of receiving waters (Batt *et al.* 2006).

The usage of phytoremediation technology is widely applied with success to remove some pollutants from wastewater. However, only very few studies have been carried out on the removal of pharmaceuticals by plants. Typhaceae is one of the frequent emergent plants and widely used in a constructed wetland. Various studies have assessed Typhaceae ability to withstand and remove pharmaceutical and other trace emerging compounds. Gray and Sedlak (2005) reported that the overall estradiol and ethinylestradiol removal from water ranged 36 to 41 % by a combination of *Typha* spp. and *Scirpus* spp. Ávila *et al.* (2015) also reported above 80% removal efficiencies for estrogenic compounds such as bisphenol A and ethinyl estradiol by *T. angustifolia*. Dai *et al.* (2017) reported that *T. angustifolia* able to remove bisphenol A and esterone about 50% in stacked constructed wetland at 0.5-

2.0 m/d flow rate. Thus, it is possible for *Typha angustifolia* absorbing norethindrone and tolerating to effects of norethindrone.

1.1 Objectives

- (i) To investigate the phyto-removal efficiency of *T. angustifolia* when exposed to norethindrone
- (ii) To access the phytotoxicity of norethindrone by *T. angustifolia* through phyto-remediation.

1.2 Thesis contribution

The increasing pharmaceutical residues such as norethindrone discharged from urban and agricultural areas into natural water resources has alarmed the public concerns. As results, there is a need to develop remediation technique to overcome the problem. Floating and emergent plants such as Typhaceae, mangrove and *Hydra* sp. have the advantage of absorbing these compounds and metabolising them from toxic to non-toxic characteristics. This can decrease the risk of the organisms from directly exposed to high concentration of these compounds. Furthermore, the occurrence of emerging or newly identified contaminants in our water resources is of continued concern for the health and safety of consuming public. This indicates the urgent need for finding cost-effective processes to retain and remove these pollutants before they reach water systems. This information is valuable for evaluating the ability of *Typha angustifolia* in tolerance and removal of norethindrone in the aquatic environment. It will later help in designing effectively constructed wetland to treat pharmaceutical contaminated water.

1.3 Scopes of study

In this context, the research was conducted in a closed environment to prevent the disturbance of weather on the studies of experiments. Besides, the phytoremoval efficiency was evaluated under hydroponic culture. Since the hydroponically research was conducted by using PVC containers, thus adsorption and other abiotic factors of norethindrone in water tank were accounted and performed as a controlled experiment of the research. Meanwhile, phytotoxicity experiment which studied on the toxicity effect of norethindrone in plants was only accounting on the physical characteristics of *T. angustifolia* and accumulation of norethindrone in the plant tissues.

CHAPTER 2

LITERATURE REVIEW

2.1 Pharmaceutical pollution

Pharmaceutical pollution due to uncontrolled agricultural and industrial revolution has resulted in a rapid increase of toxin in the aquatic environment (Ashraf *et al.* 2010). In past, Hignite and Azarnoff (1977) discovered the presence of chlorophenoxyisobutyrate and salicylic acid in the effluent of sewage disposal plant in Kansas City, Missouri. It was alarming to know that drugs, either in form of unchanged or biologically active metabolites, have high potential negative impacts to the ecosystem, especially for biota. Numerous researchers pointed out that the presence of these residues that discharged from urbanization (includes industrial) and agricultural activities are not totally eliminated with current conventional wastewater and sewage treatment plants, although the micro-pollutants remained in low concentration.

However, due to the characteristic of polymorphism and various water solubility in different pH matrix, this might lead to possible biological and toxicity effect to organisms (Caliman and Gavrilescu 2009; Fatta-Kassinos *et al.* 2011). Most of the predominantly water-soluble pharmaceuticals have limited volatility and biodegradability, resulting in partial removal from the water phase and less deposited in the sedimentation and biological treatment in sewage and wastewater treatment plants (Ternes *et al.* 2004; Bendz *et al.* 2005; Heidler *et al.* 2006). Despite their extremely low concentration, they are still capable of negative impacts on the aquatic environment, as well as interrupting the normal functioning of aquatic organisms and human health due to their unpredictable eco-toxicological effects (Stuart *et al.* 2012).



Therefore, even though these compounds usually are not commonly monitored, but they are nowadays a major cause of toxicity in plants and animals.

2.2 Synthetic steroid and hormonally pharmaceutical residues act as emerging micro-pollutants in the environment.

Steroids are originated from lipid compounds (terpenes) with containing or biologically derived from perhydro-1,2-cyclopentanophenanthrene ring system. These varieties are controlled by the type of side-chain substitution at C17, the degree of unsaturation, degree and nature of oxidation and the stereochemical characteristics of the ring junctions (Morgan and Moynihan 2000). Steroidal compounds exist in two categories: natural and synthetic, in which may not only influence in human health but also other organisms' health. Natural steroid such as sterols, bile acids, hormones (estrogen, phytoestrogen, and androgens), corticosteroids, and vitamin D can be found in biotas. On the other hand, synthetic steroid hormones like conjugated and esterified estrogen, and others (de Mes *et al.* 2005) are widely used in various pharmaceutical fields, product care fields, and industrial activities.

However, global demand for consumption has resulted the discharged of hormonal pharmaceutical residues in the overuse of hormones that has resulted with potentially negative impacts to the ecosystem. Severely damaging effects of endocrine disruption can be induced even in low concentration with those units like ng/L that are not commonly monitored (EEA 1996; Kavlock *et al.* 1996; Zoeller *et al.* 2012). The disruption process can be induced through: i) directly influence: mimic (involving agonist response, exhibit estrogenic, ant-androgenic, and androgenic activity); ii) indirectly process that involving hormone pathway disruption (Magnarelli and Fonovich 2013). These residues may not only influence gene expression process but also interrupts the normal biological process like cell proliferation and hormonal secretion. Numerous researches were conducted to study the concentration of these pharmaceutical residues in environment and the consequences of the occurrence to the environment (refer to Appendix A) Figure 2.1

REFERENCES

- Abd Manan, F., Chai T.-T. and Abd Samad A. 2015. Environmental pollution in Malaysia - Are medicinal plants potential phytoremediation agents? *Maejo Int. J. Sci. Technol.* 9(3): 288-300.
- Abdel-Ghani, N. T., Hegazy A. K., El-Chaghaby G. A. and Lima E. C. 2009. Factorial experimental design for biosorption of iron and zinc using *Typha domingensis* phytomass. *Desalination* 249(1): 343-347.
- Abdollahi, M. and Mostafalou S. 2014. Norethisterone (Norethindrone). 619-621.
- Abioye, P. 2011. Biological Remediation of Hydrocarbon and Heavy Metals Contaminated Soil. *Soil Contamination*. S. Pascucci: 127-142.
- Agostini, E., Talano M. A., Gonzalez P. S., Oller A. L. and Medina M. I. 2013. Application of hairy roots for phytoremediation: what makes them an interesting tool for this purpose? *Appl Microbiol Biotechnol* 97(3): 1017-1030.
- Ahmadi-Jouibari, T., Fattahi N. and Shamsipur M. 2014. Rapid extraction and determination of amphetamines in human urine samples using dispersive liquid-liquid microextraction and solidification of floating organic drop followed by high performance liquid chromatography. *J Pharm Biomed Anal* 94: 145-151.
- Ahn, C., Mitsch W. J. and Wolfe W. E. 2001. Effects of recycled FGD liner material on water quality and macrophytes of constructed wetlands: A mesocosm experiment. *Water Res* 35(3): 633-642.
- Al-Odaini, N. A., Zakaria M. P., Yaziz M. I., Surif S. and Abdulghani M. 2011. The occurrence of human pharmaceuticals in wastewater effluents and surface water of Langat River and its tributaries, Malaysia. *International Journal of Environmental Analytical Chemistry* 93(3): 245-264.
- Al-Odaini, N. A., Zakaria M. P., Yaziz M. I., Surif S. and Kannan N. 2012. Occurrence of synthetic hormones in sewage effluents and Langat River and its tributaries, Malaysia. *International Journal of Environmental Analytical Chemistry* 93(14): 1457-1469.
- Albuquerque, A., Oliveira J., Semitela S. and Amaral L. 2010. Evaluation of the effectiveness of horizontal subsurface flow constructed wetlands for different media. *Journal of Environmental Sciences* 22(6): 820-825.
- Almeida, C. and Nogueira J. M. 2006. Determination of steroid sex hormones in water and urine matrices by stir bar sorptive extraction and liquid chromatography with diode array detection. *J Pharm Biomed Anal* 41(4): 1303-1311.
- Amarante, L. d., Lima J. D. and Sodek L. 2006. Growth and stress conditions cause similar changes in xylem amino acids for different legume species. *Environmental and Experimental Botany* 58(1-3): 123-129.
- Amin, M. M., Bina B., Ebrahim K., Yavari Z. and Mohammadi F. 2018. Biodegradation of natural and synthetic estrogens in moving bed bioreactor. *Chinese Journal of Chemical Engineering* 26(2): 393-399.
- Asami, T., Oh K., Jikumaru Y., Shimada Y., Kaneko I., Nakano T., Takatsuto S., Fujioka S. and Yoshida S. 2004. A mammalian steroid action inhibitor spironolactone

- retards plant growth by inhibition of brassinosteroid action and induces light-induced gene expression in the dark. *J Steroid Biochem Mol Biol* 91(1-2): 41-47.
- Ashraf, M., Ozturk M. and Muhammad Sajid A. A. 2010. Plant Adaption and Phytoremediation. *Plant Adaption and Phytoremediation*. M. Ashraf, M. Ozturk and M. S. A. Ahmad, Springer: 1-34.
- Austin, H. B. 1991. The Effects of Norethindrone on the Mullerian of the American Alligator. *Gen Comp Endocrinol* 84: 300-307.
- Ávila, C., Bayona J. M., Martín I., Salas J. J. and García J. 2015. Emerging organic contaminant removal in a full-scale hybrid constructed wetland system for wastewater treatment and reuse. *Ecological Engineering* 80: 108-116.
- Ávila, C., Salas J. J., Martín I., Aragón C. and García J. 2013. Integrated treatment of combined sewer wastewater and stormwater in a hybrid constructed wetland system in southern Spain and its further reuse. *Ecological Engineering* 50: 13-20.
- Ayaz, S. Ç. 2008. Post-treatment and reuse of tertiary treated wastewater by constructed wetlands. *Desalination* 226(1-3): 249-255.
- Ayaz, S. C., Aktas O., Akca L. and Findik N. 2015. Effluent quality and reuse potential of domestic wastewater treated in a pilot-scale hybrid constructed wetland system. *J Environ Manage* 156: 115-120.
- Badejo, A. A., Omole D. O., Ndambuki J. M. and Kupolati W. K. 2017. Municipal wastewater treatment using sequential activated sludge reactor and vegetated submerged bed constructed wetland planted with *Vetiveria zizanioides*. *Ecological Engineering* 99: 525-529.
- Ballesteros, F., Vuong T. H., Secondes M. F. and Tuan P. D. 2016. Removal efficiencies of constructed wetland and efficacy of plant on treating benzene. *Sustainable Environment Research* 26(2): 93-96.
- Bartikova, H., Podlipna R. and Skalova L. 2016. Veterinary drugs in the environment and their toxicity to plants. *Chemosphere* 144: 2290-2301.
- Batt, A. L., Bruce I. B. and Aga D. S. 2006. Evaluating the vulnerability of surface waters to antibiotic contamination from varying wastewater treatment plant discharges. *Environ Pollut* 142(2): 295-302.
- Bendz, D., Paxeus N. A., Ginn T. R. and Loge F. J. 2005. Occurrence and fate of pharmaceutically active compounds in the environment, a case study: Hoje River in Sweden. *J Hazard Mater* 122(3): 195-204.
- Berlizo-Barbier, A., Vauchez A., Wiest L., Baudot R., Vulliet E. and Cren-Olive C. 2014. Multi-residue analysis of emerging pollutants in sediment using QuEChERS-based extraction followed by LC-MS/MS analysis. *Anal Bioanal Chem* 406(4): 1259-1266.
- Blakey, D. C. and White I. N. H. 1986. Destruction of cytochrome P-450 and formation of green pigments by contraceptive steroids in rat hepatocyte suspensions. *Biochem Pharmacol* 35(9): 1561-1567.

- Bolong, N., Ismail A. F., Salim M. R. and Matsuura T. 2009. A review of the effects of emerging contaminants in wastewater and options for their removal. *Desalination* 239(1-3): 229-246.
- Boxall, A. B., Johnson P., Smith E. J., Sinclair C. J., Stutt E. and Levy L. S. 2006. Uptake of veterinary medicines from soils into plants. *J Agric Food Chem* 54(6): 2288-2297.
- Braselton, W. E., Lin T. J., Mills T. M., Ellegood J. O. and Mahesh V. B. 1977. Identification and measurement by gas chromatography-mass spectrometry of norethindrone and metabolites in human urine and blood. *Journal of Steroid Biochemistry* 8(1): 9-18.
- Briggs, G. G., Bromilow R. H. and Evans A. A. 1982. Relationships between lipophilicity and root uptake and translocation of non-ionised chemicals by barley. *Pesticide Science* 13(5): 495-504.
- Broseus, R., Vincent S., Aboulfadl K., Daneshvar A., Sauve S., Barbeau B. and Prevost M. 2009. Ozone oxidation of pharmaceuticals, endocrine disruptors and pesticides during drinking water treatment. *Water Res* 43(18): 4707-4717.
- Budin, K. and Lee J.-Y. 2015. Occurrence of 17 α -Ethinylestradiol (EE2) in Inanam River, Sabah, Malaysia. *Adv. Environ. Biol.* 9(15): 28-33.
- Cakir, R., Gidirislioglu A. and Cebi U. 2015. A study on the effects of different hydraulic loading rates (HLR) on pollutant removal efficiency of subsurface horizontal-flow constructed wetlands used for treatment of domestic wastewaters. *J Environ Manage* 164: 121-128.
- Calheiros, C. S., Rangel A. O. and Castro P. M. 2007. Constructed wetland systems vegetated with different plants applied to the treatment of tannery wastewater. *Water Res* 41(8): 1790-1798.
- Calijuri, M. L., da Fonseca Santiago A., Moreira Neto R. F. and de Castro Carvalho I. 2010. Evaluation of the Ability of a Natural Wetland to Remove Heavy Metals Generated by Runways and Other Paved Areas from an Airport Complex in Brazil. *Water, Air, & Soil Pollution* 219(1-4): 319-327.
- Caliman, F. A. and Gavrilescu M. 2009. Pharmaceuticals, Personal Care Products and Endocrine Disrupting Agents in the Environment - A Review. *CLEAN - Soil, Air, Water* 37(4-5): 277-303.
- Cantarero, R., Richter P., Brown S., Ascar L. and Ahumada I. 2017. Effects of applying biosolids to soils on the adsorption and bioavailability of 17 α -ethinylestradiol and triclosan in wheat plants. *Environ Sci Pollut Res Int* 24(14): 12847-12859.
- Carballeira, T., Ruiz I. and Soto M. 2016. Effect of plants and surface loading rate on the treatment efficiency of shallow subsurface constructed wetlands. *Ecological Engineering* 90: 203-214.
- Carter, L. J., Harris E., Williams M., Ryan J. J., Kookana R. S. and Boxall A. B. 2014. Fate and uptake of pharmaceuticals in soil-plant systems. *J Agric Food Chem* 62(4): 816-825.

- Carvalho, P. N., Basto M. C. P., Almeida C. M. R. and Brix H. 2014. A review of plant–pharmaceutical interactions: from uptake and effects in crop plants to phytoremediation in constructed wetlands. *Environmental Science and Pollution Research* 21(20): 11729-11763.
- Chandra, R., Yadav S., Bharagava R. N. and Murthy R. C. 2008. Bacterial pretreatment enhances removal of heavy metals during treatment of post-methanated distillery effluent by *Typha angustata* L. *J Environ Manage* 88(4): 1016-1024.
- Chang, C.-C. and Huang S.-D. 2010a. Determination of the steroid hormone levels in water samples by dispersive liquid–liquid microextraction with solidification of a floating organic drop followed by high-performance liquid chromatography. *Analytica Chimica Acta* 662(1): 39-43.
- Chang, C. C. and Huang S. D. 2010b. Determination of the steroid hormone levels in water samples by dispersive liquid-liquid microextraction with solidification of a floating organic drop followed by high-performance liquid chromatography. *Anal Chim Acta* 662(1): 39-43.
- Chang, H., Wan Y. and Hu J. 2009. Determination and source apportionment of five classes of steroid hormones in urban rivers. *Environ Sci Technol* 43(20): 7691-7698.
- Chang, H., Wan Y., Wu S., Fan Z. and Hu J. 2011. Occurrence of androgens and progestogens in wastewater treatment plants and receiving river waters: comparison to estrogens. *Water Res* 45(2): 732-740.
- Chaoui, A. and El Ferjani E. 2013. β -Estradiol Protects Embryo Growth from Heavy-Metal Toxicity in Germinating Lentil Seeds. *Journal of Plant Growth Regulation* 32(3): 636-645.
- Chawla, S. 2010. Treatment of Endometriosis and Chronic Pelvic Pain with Letrozole and Norethindrone Acetate. *Medical Journal Armed Forces India* 66(3): 213-215.
- Chen, B., Li Y., Guo Y., Zhu L. and Schnoor J. L. 2008. Role of the Extractable Lipids and Polymeric Lipids in Sorption of Organic Contaminants onto Plant Cuticles. *Environmental Science & Technology* 42(5): 1517-1523.
- Chen, Y. L., Hong X. Q., He H., Luo H. W., Qian T. T., Li R. Z., Jiang H. and Yu H. Q. 2014. Biosorption of Cr (VI) by *Typha angustifolia*: mechanism and responses to heavy metal stress. *Bioresour Technol* 160: 89-92.
- Chiemchaisri, C., Chiemchaisri W., Junsod J., Threedeach S. and Wicranarachchi P. N. 2009. Leachate treatment and greenhouse gas emission in subsurface horizontal flow constructed wetland. *Bioresour Technol* 100(16): 3808-3814.
- Chuang, Y.-H. 2017. Uptake, accumulation and metabolism of chemicals of emerging concern in vegetables. Doctor of Philosophy PhD., Michigan State University.
- Clarke, E. and Baldwin A. H. 2002. Responses of wetland plants to ammonia and water level. *Ecological Engineering* 18(3): 257-264.
- Cormier, G., Barbeau B., Arp H. P. and Sauve S. 2015. The degradation behaviour of nine diverse contaminants in urban surface water and wastewater prior to water treatment. *Environ Sci Process Impacts* 17(12): 2051-2065.

- Crabbé, P. 1979. Some Aspects of Steroid Research Based on Natural Products from Plant Origin. *Bull. Sociétés Chim. Belg* 88(5): 345-358.
- Cui, H. and Schroder P. 2016. Uptake, translocation and possible biodegradation of the antidiabetic agent metformin by hydroponically grown *Typha latifolia*. *J Hazard Mater* 308: 355-361.
- D'Abrosca, B., Fiorentino A., Izzo A., Cefarelli G., Pascarella M. T., Uzzo P. and Monaco P. 2008. Phytotoxicity evaluation of five pharmaceutical pollutants detected in surface water on germination and growth of cultivated and spontaneous plants. *J Environ Sci Health A Tox Hazard Subst Environ Eng* 43(3): 285-294.
- Dai, Y.-n., Tao R., Tai Y.-p., Tam N. F.-y., A D. and Yang Y. 2017. Application of a full-scale newly developed stacked constructed wetland and an assembled bio-filter for reducing phenolic endocrine disrupting chemicals from secondary effluent. *Ecological Engineering* 99: 496-503.
- Davidson, S. E., Smith J. J., Helliwell C. A., Poole A. T. and Reid J. B. 2004. The pea gene LH encodes ent-kaurene oxidase. *Plant Physiol* 134(3): 1123-1134.
- Dawidowicz, A. L. and Czapczynska N. B. 2011. Sea sand disruption method (SSDM) as a valuable tool for isolating essential oil components from conifers. *Chem Biodivers* 8(11): 2045-2056.
- Dawidowicz, A. L. and Wianowska D. 2009. Application of the MSPD Technique for the HPLC Analysis of Rutin in *Sambucus nigra* L.: The Linear Correlation of the Matrix Solid-Phase Dispersion Process. *Journal of Chromatographic Science* 47(10): 914-918.
- Dawidowicz, A. L., Wianowska D. and Rado E. 2011. Matrix solid-phase dispersion with sand in chromatographic analysis of essential oils in herbs. *Phytochem Anal* 22(1): 51-58.
- de Mes, T., Zeeman G. and Lettinga G. 2005. Occurrence and fate of estrone, 17 β -estradiol and 17 α -ethynylestradiol in STPs for domestic wastewater. *Reviews in Environmental Science and Biotechnology* 2005(4): 275-311.
- Dean, M., Murphy B. T. and Burdette J. E. 2017. Phytosteroids beyond estrogens: Regulators of reproductive and endocrine function in natural products. *Mol Cell Endocrinol* 442: 98-105.
- Devillers, J., Marchand-Geneste N., Porcher J.-M. and Overview A. T. 2009. Mechanisms of Endocrine Disruptions. *Endocrine Disruption Modeling*. CRC Press: 11-46.
- Díaz-Cruz, M. S., López de Alda M. a. J. and Barceló D. 2003. Environmental behavior and analysis of veterinary and human drugs in soils, sediments and sludge. *TRAC Trends in Analytical Chemistry* 22(6): 340-351.
- Dodgen, L. K., Li J., Parker D. and Gan J. J. 2013. Uptake and accumulation of four PPCP/EDCs in two leafy vegetables. *Environ Pollut* 182: 150-156.
- Dolliver, H., Kumar K. and Gupta S. 2007. Sulfamethazine uptake by plants from manure-amended soil. *J Environ Qual* 36(4): 1224-1230.

- Dordio, A., Carvalho A. J., Teixeira D. M., Dias C. B. and Pinto A. P. 2010. Removal of pharmaceuticals in microcosm constructed wetlands using *Typha* spp. and LECA. *Bioresour Technol* 101(3): 886-892.
- Dordio, A., Ferro R., Teixeira D., Palace A. J., Pinto A. P. and Dias C. M. B. 2011a. Study on the use of *Typha* spp. for the phytotreatment of water contaminated with ibuprofen. *International Journal of Environmental Analytical Chemistry* 91(7-8): 654-667.
- Dordio, A., Pinto J., Barrocas Dias C., Pinto A. P., Palace Carvalho A. J. and Teixeira D. M. 2009a. Atenolol removal in microcosm constructed wetlands. *International Journal of Environmental Analytical Chemistry* 89(8-12): 835-848.
- Dordio, A. V., Belo M., Martins Teixeira D., Palace Carvalho A. J., Dias C. M., Pico Y. and Pinto A. P. 2011b. Evaluation of carbamazepine uptake and metabolization by *Typha* spp., a plant with potential use in phytotreatment. *Bioresour Technol* 102(17): 7827-7834.
- Dordio, A. V., Duarte C., Barreiros M., Carvalho A. J., Pinto A. P. and da Costa C. T. 2009b. Toxicity and removal efficiency of pharmaceutical metabolite clofibrac acid by *Typha* spp.--potential use for phytoremediation? *Bioresour Technol* 100(3): 1156-1161.
- Duman, F., Urey E. and Koca F. D. 2015. Temporal variation of heavy metal accumulation and translocation characteristics of narrow-leaved cattail (*Typha angustifolia* L.). *Environ Sci Pollut Res Int* 22(22): 17886-17896.
- EEA, E. C. A. 1996. European workshop on the impact of endocrine disruptors on human health and wildlife Weybridge, U. K., European Commission Agency.
- El Hamouri, B., Nazih J. and Lahjouj J. 2007. Subsurface-horizontal flow constructed wetland for sewage treatment under Moroccan climate conditions. *Desalination* 215(1-3): 153-158.
- Fan, Z., Wu S., Chang H. and Hu J. 2011. Behaviors of glucocorticoids, androgens and progestogens in a municipal sewage treatment plant: comparison to estrogens. *Environ Sci Technol* 45(7): 2725-2733.
- Fatta-Kassinos, D., Meric S. and Nikolaou A. 2011. Pharmaceutical residues in environmental waters and wastewater: current state of knowledge and future research. *Anal Bioanal Chem* 399(1): 251-275.
- Fawzy, M., Nasr M., Adel S., Nagy H. and Helmi S. 2016. Environmental approach and artificial intelligence for Ni(II) and Cd(II) biosorption from aqueous solution using *Typha domingensis* biomass. *Ecological Engineering* 95: 743-752.
- Fayad, P. B., Zamyadi A., Broseus R., Prevost M. and Sauve S. 2013. Degradation of progestagens by oxidation with potassium permanganate in wastewater effluents. *Chem Cent J* 7: 84.
- Fent, K. 2015. Progestins as endocrine disrupters in aquatic ecosystems: Concentrations, effects and risk assessment. *Environ Int* 84: 115-130.
- Franklin, A. M., Williams C. F., Andrews D. M., Woodward E. E. and Watson J. E. 2016. Uptake of Three Antibiotics and an Antiepileptic Drug by Wheat Crops Spray

- Irrigated with Wastewater Treatment Plant Effluent. *J Environ Qual* 45(2): 546-554.
- Garcia-Valcarcel, A. I., Loureiro I., Escorial C., Molero E. and Tadeo J. L. 2016. Uptake of azoles by lamb's lettuce (*Valerianella locusta* L.) grown in hydroponic conditions. *Ecotoxicol Environ Saf* 124: 138-146.
- Garvin, N., Doucette W. J. and White J. C. 2014. Investigating differences in the root to shoot transfer and xylem sap solubility of organic compounds between zucchini, squash and soybean using a pressure chamber method. *Chemosphere* 130: 98-102.
- Gavrilescu, M., Demnerova K., Aamand J., Agathos S. and Fava F. 2015. Emerging pollutants in the environment: present and future challenges in biomonitoring, ecological risks and bioremediation. *N Biotechnol* 32(1): 147-156.
- Ghosh, M., Paul J., Jana A., De A. and Mukherjee A. 2015. Use of the grass, *Vetiveria zizanioides* (L.) Nash for detoxification and phytoremediation of soils contaminated with fly ash from thermal power plants. *Ecological Engineering* 74: 258-265.
- Gillespie, W. B., Jr., Hawkins W. B., Rodgers J. H., Jr., Cano M. L. and Dorn P. B. 1999. Transfers and transformations of zinc in flow-through wetland microcosms. *Ecotoxicol Environ Saf* 43(2): 126-132.
- Gobas, F. A. P. C. 1993. A model for predicting the bioaccumulation of hydrophobic organic chemicals in aquatic food-webs: application to Lake Ontario. *Ecological Modelling* 69(1-2): 1-17.
- Gomez, C. F., Constantine L. and Huggett D. B. 2010. The influence of gill and liver metabolism on the predicted bioconcentration of three pharmaceuticals in fish. *Chemosphere* 81(10): 1189-1195.
- Gong, Z., Chandler K., Webster S., Kerley R., Buist S. and McCort-Tipton M. 2012. Simple and rapid determination of norethindrone in human plasma by supported liquid extraction and ultra performance liquid chromatography with tandem mass spectrometry. *Talanta* 91: 77-82.
- Goto, T. and Hiromi J. 2003. Toxicity of 17 α -ethynylestradiol and norethindrone, constituents of an oral contraceptive pill to the swimming and reproduction of cladoceran *Daphnia magna*, with special reference to their synergetic effect. *Marine Pollution Bulletin* 47(1-6): 139-142.
- Gray, J. L. and Sedlak D. L. 2005. The fate of estrogenic hormones in an engineered treatment wetland with dense macrophytes. *Water Environmental Research* 77(1): 24-31.
- Gray, S., Kinross J., Read P. and Marland A. 2000. The nutrient assimilative capacity of maerl as a substrate in constructed wetland systems for waste treatment. *Water Res* 34(8): 2183-2190.
- Gultekin, I. and Ince N. H. 2007. Synthetic endocrine disruptors in the environment and water remediation by advanced oxidation processes. *J Environ Manage* 85(4): 816-832.

- Guo, L., He K., Wu S., Sun H., Wang Y., Huang X. and Dong R. 2016. Optimization of high-rate TN removal in a novel constructed wetland integrated with microelectrolysis system treating high-strength digestate supernatant. *J Environ Manage* 178: 42-51.
- Haan, J. N. 1974. Effects of Diethylstilbestrol, Norethindrone, and Mestranol on Selected Microbes. Master of Science, Loyola University Chicago.
- Halder, S., Venu P. and Rao Y. V. 2014. The distinct *Typha angustifolia* (Typhaceae) ignored in Indian floras. *Rheedea* 24(1): 16-20.
- He, Y., Wang Y. and Song X. 2016. High-effective denitrification of low C/N wastewater by combined constructed wetland and biofilm-electrode reactor (CW-BER). *Bioresour Technol* 203: 245-251.
- Heidler, J., Sapkota A. and Halden R. U. 2006. Partitioning, Persistence, and Accumulation in Digested Sludge of the Topical Antiseptic Triclocarban during Wastewater Treatment. *Environmental Science & Technology* 40(11): 3634-3639.
- Hemond, H. F. and Fechner E. J. 2015. Chapter 2 - Surface Waters. Chemical Fate and Transport in the Environment (Third Edition). H. F. Hemond and E. J. Fechner. Boston, Academic Press: 75-218.
- Herklotz, P. A., Gurung P., Vanden Heuvel B. and Kinney C. A. 2010. Uptake of human pharmaceuticals by plants grown under hydroponic conditions. *Chemosphere* 78(11): 1416-1421.
- Hignite, C. and Azarnoff D. L. 1977. Drugs and drug metabolites as environmental contaminants: chlorophenoxyisobutyrate and salicylic acid in sewage water effluent. *Life Science* 20: 337-342.
- Hijosa-Valsero, M., Matamoros V., Martín-Villacorta J., Bécares E. and Bayona J. M. 2010. Assessment of full-scale natural systems for the removal of PPCPs from wastewater in small communities. *Water Res* 44(5): 1429-1439.
- Ho, Y. B., Zakaria M. P., Latif P. A. and Saari N. 2014. Occurrence of veterinary antibiotics and progesterone in broiler manure and agricultural soil in Malaysia. *Science of The Total Environment* 488-489: 261-267.
- Iino, M., Nomura T., Tamaki Y., Yamada Y., Yoneyama K., Takeuchi Y., Mori M., Asami T., Nakano T. and Yokota T. 2007. Progesterone: its occurrence in plants and involvement in plant growth. *Phytochemistry* 68(12): 1664-1673.
- Imai, S., Shiraishi A., Gamo K., Watanabe I., Okuhata H., Miyasaka H., Ikeda K., Bamba T. and Hirata K. 2007. Removal of phenolic endocrine disruptors by *Portulaca oleracea*. *J Biosci Bioeng* 103(5): 420-426.
- Inoue, J., Chamberlain K. and Bromilow R. H. 1998. Physicochemical factors affecting the uptake by roots and translocation to shoots of amine bases in barley. *Pesticide Science* 54(1): 8-21.
- Inthorn, D. 2004. Decolorization of basic, direct and reactive dyes by pre-treated narrow-leaved cattail (*Typha angustifolia* Linn.). *Bioresour Technol* 94(3): 299-306.



- Inthorn, D., Singhtho S., Thiravetyan P. and Khan E. 2004. Decolorization of basic, direct and reactive dyes by pre-treated narrow-leaved cattail (*Typha angustifolia* Linn.). *Bioresour Technol* 94(3): 299-306.
- Inui, H., Sawada M., Goto J., Yamazaki K., Kodama N., Tsuruta H. and Eun H. 2013. A major latex-like protein is a key factor in crop contamination by persistent organic pollutants. *Plant Physiol.* 161: 2128–2135.
- Inui, H., Wakai T., Gion K., Yamazaki K., Kim Y.-S. and Eun H. 2014. Congener Specificity in the Accumulation of Dioxins and Dioxin-Like Compounds in Zucchini Plants Grown Hydroponically. *Biosci Biotechnol Biochem* 75(4): 705-710.
- Iori, V., Zacchini M. and Pietrini F. 2013. Growth, physiological response and phytoremoval capability of two willow clones exposed to ibuprofen under hydroponic culture. *J Hazard Mater* 262: 796-804.
- Janezko, A. and Skoczowski A. 2005. Mammalian sex hormones in plants. *Folia Histochemica Et Cytobiologica* 43(2): 71-79.
- Jing, S.-R., Lin Y.-F., Lee D.-Y. and Wang T.-W. 2001. Nutrient removal from polluted river water by using constructed wetlands. *Bioresour Technol* 76(2): 131-135.
- Kamyab, S., Fotherby K. and Klopper A. I. 1968. Metabolism of [4-14c]Norethisterone in Women. *Journal of Endocrinology* 41(2): 263-272.
- Karnjanapiboonwong, A., Chase D. A., Canas J. E., Jackson W. A., Maul J. D., Morse A. N. and Anderson T. A. 2011. Uptake of 17alpha-ethynylestradiol and triclosan in pinto bean, *Phaseolus vulgaris*. *Ecotoxicol Environ Saf* 74(5): 1336-1342.
- Kaser, D. J., Missmer S. A., Berry K. F. and Laufer M. R. 2012. Use of norethindrone acetate alone for postoperative suppression of endometriosis symptoms. *J Pediatr Adolesc Gynecol* 25(2): 105-108.
- Kavlock, R. J., Daston G. P., DeRosa C., Fenner-Crisp P., Gray L. E., Kaattari S., Lucier G., Luster M., Mac M. J., Maczka C., Miller R., Moore J., Rolland R., Scott G., Sheehan D. M., Sinks T. and Tilson H. A. 1996. Research Needs for the Risk Assessment of Health and Environmental Effects of Endocrine Disruptors: A Report of the U.S. EPA-Sponsored Workshop. *Environmental Health Perspectives* 104: 715-740.
- Khandare, R. V., Rane N. R., Waghmode T. R. and Govindwar S. P. 2012. Bacterial assisted phytoremediation for enhanced degradation of highly sulfonated diazo reactive dye. *Environ Sci Pollut Res Int* 19(5): 1709-1718.
- Kinraide, T. B. 2006. Plasma Membrane Surface Potential (ψ_{pm}) as a Determinant of Ion Bioavailability: A Critical Analysis of New and Published Toxicological Studies and a Simplified Method for the Computation of Plant ψ_{pm} . *Environmental Toxicology and Chemistry* 25(12): 3188.
- Kissoon, L. T. T., Jacob D. L. and Otte M. L. 2011. Multiple elements in *Typha angustifolia* rhizosphere and plants: Wetland versus dryland. *Environmental and Experimental Botany* 72(2): 232-241.

- Kuhnz, W., Heuner A., Hümpel M., Seifert W. and Michaelis K. 1997. In vivo conversion of norethisterone and norethisterone acetate to ethinyl estradiol in postmenopausal women. *Contraception* 56(6): 379-385.
- Kurwadkar, S., Struckhoff G., Pugh K. and Singh O. 2017. Uptake and translocation of sulfamethazine by alfalfa grown under hydroponic conditions. *J Environ Sci (China)* 53: 217-223.
- Kusch, P., Wießner A., Kappelmeyer U., Weißbrodt E., Kästner M. and Stottmeister U. 2003. Annual cycle of nitrogen removal by a pilot-scale subsurface horizontal flow in a constructed wetland under moderate climate. *Water Res* 37(17): 4236-4242.
- Kuster, M., Azevedo D. A., Lopez de Alda M. J., Aquino Neto F. R. and Barcelo D. 2009. Analysis of phytoestrogens, progestogens and estrogens in environmental waters from Rio de Janeiro (Brazil). *Environ Int* 35(7): 997-1003.
- Kuster, M., López de Alda M. J., Hernando M. D., Petrovic M., Martín-Alonso J. and Barceló D. 2008. Analysis and occurrence of pharmaceuticals, estrogens, progestogens and polar pesticides in sewage treatment plant effluents, river water and drinking water in the Llobregat river basin (Barcelona, Spain). *Journal of Hydrology* 358(1-2): 112-123.
- Labadie, P. and Budzinski H. 2005. Development of an analytical procedure for determination of selected estrogens and progestagens in water samples. *Anal Bioanal Chem* 381(6): 1199-1205.
- Labella, A., Caniani D., Hughes-Riley T., Morris R. H., Newton M. I., Hawes P., Puigagut J., García J. and Uggetti E. 2015. Assessing the economic suitability of aeration and the influence of bed heating on constructed wetlands treatment efficiency and life-span. *Ecological Engineering* 83: 184-190.
- Leung, J. Y. S., Cai Q. and Tam N. F. Y. 2016. Comparing subsurface flow constructed wetlands with mangrove plants and freshwater wetland plants for removing nutrients and toxic pollutants. *Ecological Engineering* 95: 129-137.
- Li, G., Zhai J., He Q., Zhi Y., Xiao H. and Rong J. 2014. Phytoremediation of levonorgestrel in aquatic environment by hydrophytes. *J Environ Sci (China)* 26(9): 1869-1873.
- Li, W., Li Y. H., Li A. C., Zhou S. and Naidong W. 2005. Simultaneous determination of norethindrone and ethinyl estradiol in human plasma by high performance liquid chromatography with tandem mass spectrometry--experiences on developing a highly selective method using derivatization reagent for enhancing sensitivity. *J Chromatogr B Analyt Technol Biomed Life Sci* 825(2): 223-232.
- Li, W. C. 2014. Occurrence, sources, and fate of pharmaceuticals in aquatic environment and soil. *Environ Pollut* 187: 193-201.
- Li, X., Yu H., Xu S. and Hua R. 2013. Uptake of three sulfonamides from contaminated soil by pakchoi cabbage. *Ecotoxicol Environ Saf* 92: 297-302.
- Li, Y., Zhang J., Zhu G., Liu Y., Wu B., Ng W. J., Appan A. and Tan S. K. 2016. Phytoextraction, phytotransformation and rhizodegradation of ibuprofen associated with *Typha angustifolia* in a horizontal subsurface flow constructed wetland. *Water Res* 102: 294-304.

- Lim, P. E., Tay M. G., Mak K. Y. and Mohamed N. 2003. The effect of heavy metals on nitrogen and oxygen demand removal in constructed wetlands. *Science of The Total Environment* 301(1-3): 13-21.
- Lim, P. E., Wong T. F. and Lim D. V. 2001. Oxygen demand, nitrogen and copper removal by free-water-surface and subsurface-flow constructed wetlands under tropical conditions. *Environ Int* 26(5-6): 425-431.
- Limmer, M. A. and Burken J. G. 2014. Plant Translocation of Organic Compounds: Molecular and Physicochemical Predictors. *Environmental Science & Technology Letters* 1(2): 156-161.
- Lin, Y.-F., Jing S.-R. and Lee D.-Y. 2003. The potential use of constructed wetlands in a recirculating aquaculture system for shrimp culture. *Environ Pollut* 123(1): 107-113.
- Lin, Y.-F., Jing S.-R., Wang T.-W. and Lee D.-Y. 2002. Effects of macrophytes and external carbon sources on nitrate removal from groundwater in constructed wetlands. *Environ Pollut* 119(3): 413-420.
- Lin, Y. F., Jing S. R., Lee D. Y., Chang Y. F. and Shih K. C. 2008. Nitrate removal from groundwater using constructed wetlands under various hydraulic loading rates. *Bioresour Technol* 99(16): 7504-7513.
- Liu, L., Liu Y.-h., Liu C.-x., Wang Z., Dong J., Zhu G.-f. and Huang X. 2013. Potential effect and accumulation of veterinary antibiotics in *Phragmites australis* under hydroponic conditions. *Ecological Engineering* 53: 138-143.
- Liu, S., Ying G. G., Zhang R. Q., Zhou L. J., Lai H. J. and Chen Z. F. 2012a. Fate and occurrence of steroids in swine and dairy cattle farms with different farming scales and wastes disposal systems. *Environ Pollut* 170: 190-201.
- Liu, S., Ying G. G., Zhao J. L., Chen F., Yang B., Zhou L. J. and Lai H. J. 2011. Trace analysis of 28 steroids in surface water, wastewater and sludge samples by rapid resolution liquid chromatography-electrospray ionization tandem mass spectrometry. *J Chromatogr A* 1218(10): 1367-1378.
- Liu, S., Ying G. G., Zhou L. J., Zhang R. Q., Chen Z. F. and Lai H. J. 2012b. Steroids in a typical swine farm and their release into the environment. *Water Res* 46(12): 3754-3768.
- Lizama Allende, K., McCarthy D. T. and Fletcher T. D. 2014. The influence of media type on removal of arsenic, iron and boron from acidic wastewater in horizontal flow wetland microcosms planted with *Phragmites australis*. *Chemical Engineering Journal* 246: 217-228.
- Lo, V. Y. and Saibeh K. 2013. Phytoremediation using *Typha angustifolia* L. for mine water effluence treatment: case study of Ex-Mamut Copper Mine, Ranau, Sabah. *Borneo Science* 33: 16-22.
- Loh, S. H., Ong S. T., Ngu M. L. and Mohd Ariffin M. 2017. Rapid Extraction of Bisphenol A by Dispersive Liquid-liquid Microextraction based on Solidification of Floating Organic. *Sains Malaysia* 46(4): 615-621.
- López de Alda, M. J. and Barceló D. 2000. Determination of steroid sex hormones and related synthetic compounds considered as endocrine disrupters in water by

- liquid chromatography–diode array detection–mass spectrometry. *Journal of Chromatography A* 892(1-2): 391-406.
- López de Alda, M. J. and Barceló D. 2001. Use of solid-phase extraction in various of its modalities for sample preparation in the determination of estrogens and progestogens in sediment and water. *Journal of Chromatography A* 938(1-2): 145-153.
- Magnarelli, G. and Fonovich T. 2013. Protein phosphorylation pathways disruption by pesticides. *Advances in Biological Chemistry* 03(05): 460-474.
- Maier, W. E. and Herman J. R. 2001. Pharmacology and toxicology of ethinyl estradiol and norethindrone acetate in experimental animals. *Regul Toxicol Pharmacol* 34(1): 53-61.
- Manhita, A. C., Teixeira D. M. and da Costa C. T. 2006. Application of sample disruption methods in the extraction of anthocyanins from solid or semi-solid vegetable samples. *J Chromatogr A* 1129(1): 14-20.
- Mantovi, P., Marmiroli M., Maestri E., Tagliavini S., Piccinini S. and Marmiroli N. 2003. Application of a horizontal subsurface flow constructed wetland on treatment of dairy parlor wastewater. *Bioresour Technol* 88(2): 85-94.
- Martins, S., Aguilar C. N., Teixeira J. A. and Mussatto S. I. 2012. Bioactive compounds (phytoestrogens) recovery from *Larrea tridentata* leaves by solvents extraction. *Separation and Purification Technology* 88: 163-167.
- Mateus, D. M. R., Vaz M. M. N. and Pinho H. J. O. 2012. Fragmented limestone wastes as a constructed wetland substrate for phosphorus removal. *Ecological Engineering* 41: 65-69.
- Matsoukis, A., Kamoutsis A., Koutsouropoulou M. and Kanakis A. 2006. Study of meteorological parameters and the growth regulator ethephon in combination with growth of narcissus (*Narcissus cv. Ice Follies*). 8th Conference on Meteorology-Climatology-Atmospheric Physics. Athen. A.
- Mayberry, D. O., Kowblansky M., Lane P. A. and Wray P. E. 1990. Determination of norethindrone stability on red delicious apple. *Journal of Pharmaceutical Sciences* 79(8): 746-749.
- Miller, E. L., Nason S. L., Karthikeyan K. G. and Pedersen J. A. 2016. Root Uptake of Pharmaceuticals and Personal Care Product Ingredients. *Environ Sci Technol* 50(2): 525-541.
- Morari, F. and Giardini L. 2009. Municipal wastewater treatment with vertical flow constructed wetlands for irrigation reuse. *Ecological Engineering* 35(5): 643-653.
- Morgan, B. P. and Moynihan M. S. 2000. Steroids.
- Murano, H., Otani T. and Seike N. 2010. Dieldrin-dissolving abilities of the xylem saps of several plant families, particularly *Cucurbita pepo* L. *Environ. Toxicol. Chem.* 29: 2269-2277.
- Nilratnisakorn, S., Thiravetyan P. and Nakbanpote W. 2007. Synthetic reactive dye wastewater treatment by narrow-leaved cattails (*Typha angustifolia* Linn.): effects of dye, salinity and metals. *Sci Total Environ* 384(1-3): 67-76.

- Okuhata, H., Ikeda K., Miyasaka H., Takahashi S., Matsui T., Nakayama H., Kato K. and Hirata K. 2010. Floricultural *Salvia* plants have a high ability to eliminate bisphenol A. *J Biosci Bioeng* 110(1): 99-101.
- Olguin, E. J., Sanchez-Galvan G., Gonzalez-Portela R. E. and Lopez-Vela M. 2008. Constructed wetland mesocosms for the treatment of diluted sugarcane molasses stillage from ethanol production using *Pontederia sagittata*. *Water Res* 42(14): 3659-3666.
- Oliveira, J. P. V., Pereira M. P., Duarte V. P., Correa F. F., Castro E. M. and Pereira F. J. 2018. Cadmium tolerance of *Typha domingensis* Pers. (Typhaceae) as related to growth and leaf morphophysiology. *Braz J Biol* 78(3): 509-516.
- Osunmakinde, C. O. 2014. Dispersive Liquid-Liquid Micro-Extraction Of Steroidal Hormones And Determination In Wastewater Using High Pressure Liquid Chromatography-Charged Aerosol Detector, University of South Africa.
- Passatore, L., Rossetti S., Juwarkar A. A. and Massacci A. 2014. Phytoremediation and bioremediation of polychlorinated biphenyls (PCBs): state of knowledge and research perspectives. *J Hazard Mater* 278: 189-202.
- Paulos, P., Runnalls T. J., Nallani G., La Point T., Scott A. P., Sumpter J. P. and Huggett D. B. 2010. Reproductive responses in fathead minnow and Japanese medaka following exposure to a synthetic progestin, Norethindrone. *Aquat Toxicol* 99(2): 256-262.
- Payus, C., John C., Wan V. L., Hsiang T. W. and Kui W. N. 2016. Occurrence of Steroid Sex Hormone Progesterone in Influent and Effluent of Oxidation Pond and the River Outlet of Waste Water Treatment Case Study. *Journal of Environmental Science and Technology* 9(5): 399-406.
- Payus, C., Roger J. and Ping S. Y. 2017. Steroid Sex Hormone Ethinylestradiol in Recycled Wastewater at Recreational Park. *Journal of Pharmacology and Toxicology* 12(3): 148-153.
- Petersen, L. H., Hala D., Carty D., Cantu M., Martinovic D. and Huggett D. B. 2015. Effects of progesterone and norethindrone on female fathead minnow (*Pimephales promelas*) steroidogenesis. *Environ Toxicol Chem* 34(2): 379-390.
- Petrovic, M., Solé M., López De Alda M. J. and Barceló D. 2002a. Endocrine disruptors in sewage treatment plants, receiving river waters, and sediments: Integration of chemical analysis and biological effects on feral carp. *Environmental Toxicology and Chemistry* 21(10): 2146-2156.
- Petrovic, M., Tavazzi S. and Barcelo D. 2002b. Column-switching system with restricted access pre-column packing for an integrated sample cleanup and liquid chromatographic–mass spectrometric analysis of alkylphenolic compounds and steroid sex hormones in sediment. *Journal of Chromatography A* 971(1-2): 37-45.
- Pommier, F., Sioufi A. and Godbillon J. 1995. Simultaneous determination of norethisterone and six metabolites in human plasma by capillary gas chromatography with mass-selective detection. *Journal of Chromatography B: Biomedical Sciences and Applications* 674(2): 155-165.

- Rana, R. S., Singh P., Kandari V., Singh R., Dobhal R. and Gupta S. 2014. A review on characterization and bioremediation of pharmaceutical industries' wastewater: an Indian perspective. *Applied Water Science* 7(1): 1-12.
- Redder, A., Durr M., Daeschlein G., Baeder-Bederski O., Koch C., Muller R., Exner M. and Borneff-Lipp M. 2010. Constructed wetlands--Are they safe in reducing protozoan parasites? *Int J Hyg Environ Health* 213(1): 72-77.
- Redshaw, C. H., Wootton V. G. and Rowland S. J. 2008. Uptake of the pharmaceutical Fluoxetine Hydrochloride from growth medium by Brassicaceae. *Phytochemistry* 69(13): 2510-2516.
- Rezaee, M., Assadi Y., Milani Hosseini M. R., Aghaee E., Ahmadi F. and Berijani S. 2006. Determination of organic compounds in water using dispersive liquid-liquid microextraction. *J Chromatogr A* 1116(1-2): 1-9.
- Rodriguez-Hernandez, M. C., Garcia De la-Cruz R. F., Leyva E. and Navarro-Tovar G. 2017. *Typha latifolia* as potential phytoremediator of 2,4-dichlorophenol: Analysis of tolerance, uptake and possible transformation processes. *Chemosphere* 173: 190-198.
- Rogers, S. M. D. 2003. Tissue culture and wetland establishment of the freshwater monocots *Carex*, *Juncus*, *Scirpus*, and *Typha*. *In Vitro Cellular & Developmental Biology - Plant* 39(1): 1-5.
- Safholm, M., Ribbenstedt A., Fick J. and Berg C. 2014. Risks of hormonally active pharmaceuticals to amphibians: a growing concern regarding progestagens. *Philos Trans R Soc Lond B Biol Sci* 369(1656).
- Sahlberg, B.-L. 1987. The characterization of sulphated metabolites of norethindrone in human milk after oral administration of contraceptive steroids. *Journal of Steroid Biochemistry* 26(4): 481-485.
- Sahlberg, B.-L., Landgren B.-M. and Axelson M. 1987. Metabolic profiles of endogenous and ethynyl steroids in plasma and urine from women during administration of oral contraceptives. *Journal of Steroid Biochemistry* 26(5): 609-617.
- Sanagi, M. M., Abbas H. H., Ibrahim W. A. and Aboul-Enien H. Y. 2012. Dispersive liquid-liquid microextraction method based on solidification of floating organic droplet for the determination of triazine herbicides in water and sugarcane samples. *Food Chem* 133(2): 557-562.
- Santos, M., Hendry D., Sangi-Haghpeykar H. and Dietrich J. E. 2014. Retrospective review of norethindrone use in adolescents. *J Pediatr Adolesc Gynecol* 27(1): 41-44.
- Sasmaz, A., Obek E. and Hasar H. 2008. The accumulation of heavy metals in *Typha latifolia* L. grown in a stream carrying secondary effluent. *Ecological Engineering* 33(3-4): 278-284.
- Sawaittayothin, V. and Polprasert C. 2007. Nitrogen mass balance and microbial analysis of constructed wetlands treating municipal landfill leachate. *Bioresour Technol* 98(3): 565-570.

- Schoonen, W. G. E. J., Deckers G. H., de Gooijer M. E., de Ries R. and Kloosterboer H. J. 2000. Hormonal properties of norethisterone, 7 α -methyl-norethisterone and their derivatives. *J Steroid Biochem Mol Biol* 74(4): 213-222.
- Schulz, C., Gelbrecht J. and Rennert B. 2003. Treatment of rainbow trout farm effluents in constructed wetland with emergent plants and subsurface horizontal water flow. *Aquaculture* 217(1-4): 207-221.
- Senzia, M. A., Mashauri D. A. and Mayo A. W. 2003. Suitability of constructed wetlands and waste stabilisation ponds in wastewater treatment: nitrogen transformation and removal. *Physics and Chemistry of the Earth, Parts A/B/C* 28(20-27): 1117-1124.
- Seo, Y.-H., Cho B.-O., Kang A.-S., Jeong B.-C. and Jung Y.-S. 2010. Antibiotic Uptake by Plants from Soil Applied with Antibiotic-Treated Animal Manure. *Korean J. Soil Sci. Fert.* 43(4): 466-470.
- Shehzadi, M., Afzal M., Khan M. U., Islam E., Mobin A., Anwar S. and Khan Q. M. 2014. Enhanced degradation of textile effluent in constructed wetland system using *Typha domingensis* and textile effluent-degrading endophytic bacteria. *Water Res* 58: 152-159.
- Shu, X. P. 2010. *Typha* Linnaceus. *FL China* 23: 161-163.
- Simon, J. A., Liu J. H., Speroff L., Shumel B. S. and Symons J. P. 2003. Reduced vaginal bleeding in postmenopausal women who receive combined norethindrone acetate and low-dose ethinyl estradiol therapy versus combined conjugated equine estrogens and medroxyprogesterone acetate therapy. *Am J Obstet Gynecol* 188(1): 92-99.
- Solé, M., López de Alda M. J., Castillo M., Porte C., Ladegaard-Pedersen K. and Barceló D. 2000. Estrogenicity Determination in Sewage Treatment Plants and Surface Waters from the Catalanian Area (NE Spain). *Environmental Science & Technology* 34(24): 5076-5083.
- Stuart, M., Lapworth D., Crane E. and Hart A. 2012. Review of risk from potential emerging contaminants in UK groundwater. *Sci Total Environ* 416: 1-21.
- Stumm-Zollinger, E. and Fair G. M. 1965. Biodegradation of steroid hormones. *Journal (Water Pollution Control Federation)* 37(11): 1506-1510.
- Sui, Q., Cao X., Lu S., Zhao W., Qiu Z. and Yu G. 2015. Occurrence, sources and fate of pharmaceuticals and personal care products in the groundwater: A review. *Emerging Contaminants* 1(1): 14-24.
- Sun, L., Yong W., Chu X. and Lin J. M. 2009. Simultaneous determination of 15 steroidal oral contraceptives in water using solid-phase disk extraction followed by high performance liquid chromatography-tandem mass spectrometry. *J Chromatogr A* 1216(28): 5416-5423.
- Susarla, S., Medina V. F. and McCutcheon S. C. 2002. Phytoremediation: An ecological solution to organic chemical contamination. *Ecological Engineering* 18(5): 647-658.

- Tanoue, R., Sato Y., Motoyama M., Nakagawa S., Shinohara R. and Nomiya K. 2012. Plant uptake of pharmaceutical chemicals detected in recycled organic manure and reclaimed wastewater. *J Agric Food Chem* 60(41): 10203-10211.
- Tee, H. C., Seng C. E., Noor A. M. and Lim P. E. 2009. Performance comparison of constructed wetlands with gravel- and rice husk-based media for phenol and nitrogen removal. *Sci Total Environ* 407(11): 3563-3571.
- Teixeira, D. M., Canelas V. C., do Canto A. M., Teixeira J. M. G. and Dias C. B. 2009. HPLC-DAD Quantification of Phenolic Compounds Contributing to the Antioxidant Activity of *Maclura pomifera*, *Ficus carica* and *Ficus elastica* Extracts. *Analytical Letters* 42(18): 2986-3003.
- Teixeira, D. M. and Costa C. T. d. 2005. Novel methods to extract flavanones and xanthones from the root bark of *Maclura pomifera*. *Journal of Chromatography A* 1062(2): 175-181.
- Teixeira, D. M., Patao R. F., Coelho A. V. and da Costa C. T. 2006. Comparison between sample disruption methods and solid-liquid extraction (SLE) to extract phenolic compounds from *Ficus carica* leaves. *J Chromatogr A* 1103(1): 22-28.
- Ternes, T. A., Joss A. and Siegrist H. 2004. Peer Reviewed: Scrutinizing Pharmaceuticals and Personal Care Products in Wastewater Treatment. *Environmental Science & Technology* 38(20): 392A-399A.
- Trapp, S. 2000. Modelling uptake into roots and subsequent translocation of neutral and ionisable organic compounds. *Pest Management Science* 56(9): 767-778.
- Trapp, S. 2009. Bioaccumulation of polar and ionizable compounds in plants. *Ecotoxicology Modeling*. J. Devillers, Springer US: 299-353.
- Trapp, S. and Miglioranza K. S. B. 2001. Sorption of Lipophilic Organic Compounds to Wood and Implications for Their Environmental Fate. *Environmental Science & Technology* 35(8): 1561-1566.
- Turgeon, R. and Wolf S. 2009. Phloem transport: Cellular pathways and molecular trafficking. *Annu. Rev. Plant Biol.* 60: 207-221.
- Türker, O. C., Türe C., Böcük H., Çiçek A. and Yakar A. 2016. Role of plants and vegetation structure on boron (B) removal process in constructed wetlands. *Ecological Engineering* 88: 143-152.
- Uggetti, E., Hughes-Riley T., Morris R. H., Newton M. I., Trabi C. L., Hawes P., Puigagut J. and Garcia J. 2016. Intermittent aeration to improve wastewater treatment efficiency in pilot-scale constructed wetland. *Sci Total Environ* 559: 212-217.
- U. S. Environmental Protection Agency (USEPA), 2007. Method 1698: Steroid and Hormones in Water, Soil and Biosolids by HRGC/HRMS. USA, U. S. Environmental Protection Agency.
- Vafaei, F., Movafeghi A. and Khataee A. 2013. Evaluation of antioxidant enzymes activities and identification of intermediate products during phytoremediation of an anionic dye (C.I. Acid Blue 92) by pennywort (*Hydrocotyle vulgaris*). *Journal of Environmental Sciences* 25(11): 2214-2222.

- Verlicchi, P., Galletti A., Petrovic M. and Barceló D. 2010. Hospital effluents as a source of emerging pollutants: An overview of micropollutants and sustainable treatment options. *Journal of Hydrology* 389(3-4): 416-428.
- Verliefde, A., Cornelissen E., Amy G., Van der Bruggen B. and van Dijk H. 2007. Priority organic micropollutants in water sources in Flanders and the Netherlands and assessment of removal possibilities with nanofiltration. *Environ Pollut* 146(1): 281-289.
- Viñas, P., Campillo N. and Andruch V. 2015. Recent achievements in solidified floating organic drop microextraction. *TrAC Trends in Analytical Chemistry* 68: 48-77.
- Vulliet, E., Baugros J. B., Flament-Waton M. M. and Grenier-Loustalot M. F. 2007. Analytical methods for the determination of selected steroid sex hormones and corticosteroids in wastewater. *Anal Bioanal Chem* 387(6): 2143-2151.
- Vulliet, E., Cren-Olivé C. and Grenier-Loustalot M.-F. 2009. Occurrence of pharmaceuticals and hormones in drinking water treated from surface waters. *Environmental Chemistry Letters* 9(1): 103-114.
- Wallace, E. D. 1999. Some characteristics of the uterine progesterone receptor and the effects of norethindrone on conception in the rock hyrax (*Procavia capensis*), University of Pretoria.
- Wang, Q. and Kelly B. C. 2017. Occurrence, distribution and bioaccumulation behaviour of hydrophobic organic contaminants in a large-scale constructed wetland in Singapore. *Chemosphere* 183: 257-265.
- Wang, Y., Wang J., Zhao X., Song X. and Gong J. 2016. The inhibition and adaptability of four wetland plant species to high concentration of ammonia wastewater and nitrogen removal efficiency in constructed wetlands. *Bioresour Technol* 202: 198-205.
- Ward, D. B. 2007. Keys to the flora of florida: 15, *Typha* (Typhaceae). *Phytologia* 89(1): 58-65.
- Wei, Q., Xin X., Du B., Wu D., Han Y., Zhao Y., Cai Y., Li R., Yang M. and Li H. 2010. Electrochemical immunosensor for norethisterone based on signal amplification strategy of graphene sheets and multienzyme functionalized mesoporous silica nanoparticles. *Biosens Bioelectron* 26(2): 723-729.
- Wen, B., Li L., Zhang H., Ma Y., Shan X. Q. and Zhang S. 2014. Field study on the uptake and translocation of perfluoroalkyl acids (PFAAs) by wheat (*Triticum aestivum* L.) grown in biosolids-amended soils. *Environ Pollut* 184: 547-554.
- Wen, Y., Chen Y., Zheng N., Yang D. and Zhou Q. 2010. Effects of plant biomass on nitrate removal and transformation of carbon sources in subsurface-flow constructed wetlands. *Bioresour Technol* 101(19): 7286-7292.
- White, I. N. H. 1980. Structure-activity relationships in the destruction of cytochrome P-450 mediated by certain ethynyl-substituted compounds in rats. *Biochem Pharmacol* 29: 3253-3255.
- White, I. N. H. 1984. Suicidal destruction of cytochrome P-450 by ethynyl substituted compounds. *Pharmaceutical Research* 1(4): 141-148.

- Wianowska, D. 2015. Application of Sea Sand Disruption Method for HPLC Determination of Quercetin in Plants. *Journal of Liquid Chromatography & Related Technologies* 38(10): 1037-1043.
- Wianowska, D., Dawidowicz A. L., Bernacik K. and Typek R. 2016. Determining the true content of quercetin and its derivatives in plants employing SSDM and LC-MS analysis. *European Food Research and Technology* 243(1): 27-40.
- Wianowska, D., Typek R. and Dawidowicz A. L. 2015. How to eliminate the formation of chlorogenic acids artefacts during plants analysis? Sea sand disruption method (SSDM) in the HPLC analysis of chlorogenic acids and their native derivatives in plants. *Phytochemistry* 117: 489-499.
- Wibbels, T. and Crews D. 1995. Steroid-induced sex determination at incubation temperatures producing mixed sex ratios in a turtle with TSD. *Gen Comp Endocrinol* 100(1): 53-60.
- Wießner, A., Kusch P., Stottmeister U., Struckmann D. and Jank M. 1999. Treating a lignite pyrolysis wastewater in a constructed subsurface flow wetland. *Water Res* 33(5): 1296-1302.
- Wistuba, J., Luetjens C. M., Ehmcke J., Redmann K., Damm O. S., Steinhoff A., Sandhowe-Klaverkamp R., Nieschlag E., Simoni M. and Schlatt S. 2013. Experimental endocrine manipulation by contraceptive regimen in the male marmoset (*Callithrix jacchus*). *Reproduction* 145(4): 439-451.
- Wu, C., Spongberg A. L., Witter J. D. and Sridhar B. B. 2012. Transfer of wastewater associated pharmaceuticals and personal care products to crop plants from biosolids treated soil. *Ecotoxicol Environ Saf* 85: 104-109.
- Wu, J., Feng Y., Dai Y., Cui N., Anderson B. and Cheng S. 2016a. Biological mechanisms associated with triazophos (TAP) removal by horizontal subsurface flow constructed wetlands (HSFCW). *Sci Total Environ* 553: 13-19.
- Wu, M., Pan C., Yang M., Xu B., Lei X., Ma J., Cai L. and Chen J. 2016b. Chemical analysis of fish bile extracts for monitoring endocrine disrupting chemical exposure in water: Bisphenol A, alkylphenols, and norethindrone. *Environ Toxicol Chem* 35(1): 182-190.
- Wu, X., Ernst F., Conkle J. L. and Gan J. 2013. Comparative uptake and translocation of pharmaceutical and personal care products (PPCPs) by common vegetables. *Environ Int* 60: 15-22.
- Xie, S., Xiang B., Zhang M. and Deng H. 2010. Determination of medroxyprogesterone in water samples using dispersive liquid-liquid microextraction with low solvent consumption. *Microchimica Acta* 168(3-4): 253-258.
- Xu, J., Li C., Yang F., Dong Z., Zhang J., Zhao Y., Qi P. and Hu Z. 2011a. Typha angustifolia stress tolerance to wastewater with different levels of chemical oxygen demand. *Desalination* 280(1-3): 58-62.
- Xu, W., Shi W., Yan F., Zhang B. and Liang J. 2011b. Mechanisms of cadmium detoxification in cattail (*Typha angustifolia* L.). *Aquatic Botany* 94(1): 37-43.
- Yadav, S. and Chandra R. 2010. Heavy metals accumulation and ecophysiological effect on *Typha angustifolia* L. and *Cyperus esculentus* L. growing in distillery and

- tannery effluent polluted natural wetland site, Unnao, India. *Environmental Earth Sciences* 62(6): 1235-1243.
- Yalcuk, A., Pakdil N. B. and Turan S. Y. 2010. Performance evaluation on the treatment of olive mill waste water in vertical subsurface flow constructed wetlands. *Desalination* 262(1-3): 209-214.
- Yalcuk, A. and Ugurlu A. 2009. Comparison of horizontal and vertical constructed wetland systems for landfill leachate treatment. *Bioresour Technol* 100(9): 2521-2526.
- Yalkowsky, S. H., He Y. and Jain P. 2010. Handbook of Aqueous Solubility Data, Second Edition, CRC Press.
- Yang, Y. Y., Gray J. L., Furlong E. T., Davis J. G., Revello R. C. and Borch T. 2012. Steroid hormone runoff from agricultural test plots applied with municipal biosolids. *Environ Sci Technol* 46(5): 2746-2754.
- Yi, H. and Crowley D. E. 2007. Biostimulation of PAH Degradation with Plants Containing High Concentrations of Linoleic Acid. *Environmental Science & Technology* 41(12): 4382-4388.
- Yoon, Y., Westerhoff P., Snyder S. A., Wert E. C. and Yoon J. 2007. Removal of endocrine disrupting compounds and pharmaceuticals by nanofiltration and ultrafiltration membranes. *Desalination* 202(1-3): 16-23.
- Yotis, W. 1967. In vivo and in vitro action of norethindrone on staphylococci. *Journal of Bacteriology* 94(5): 1353-1358.
- Yotis, W. and Stanke R. 1966. Bacteriostatic action of progesterone on staphylococci and other microorganisms. *Journal of Bacteriology* 92(5): 1285-1289.
- Yuan, K., Kang H., Yue Z., Yang L., Lin L., Wang X. and Luan T. 2015. Determination of 13 endocrine disrupting chemicals in sediments by gas chromatography–mass spectrometry using subcritical water extraction coupled with dispersed liquid–liquid microextraction and derivatization. *Analytica Chimica Acta* 866: 41-47.
- Zacchini, M., Pietrini F., Scarascia Mugnozza G., Iori V., Pietrosanti L. and Massacci A. 2008. Metal Tolerance, Accumulation and Translocation in Poplar and Willow Clones Treated with Cadmium in Hydroponics. *Water, Air, and Soil Pollution* 197(1-4): 23-34.
- Zhang, A., Wang J. and Li Y. 2015. Performance of calcium peroxide for removal of endocrine-disrupting compounds in waste activated sludge and promotion of sludge solubilization. *Water Res* 71: 125-139.
- Zhang, C., Yin Q., Wen Y., Guo W., Liu C. and Zhou Q. 2016. Enhanced nitrate removal in self-supplying carbon source constructed wetlands treating secondary effluent: The roles of plants and plant fermentation broth. *Ecological Engineering* 91: 310-316.
- Zhang, D. Q., Gersberg R. M., Hua T., Zhu J., Goyal M. K., Ng W. J. and Tan S. K. 2013. Fate of pharmaceutical compounds in hydroponic mesocosms planted with *Scirpus validus*. *Environ Pollut* 181: 98-106.

- Zhao, J., Zhao Y., Xu Z., Doherty L. and Liu R. 2016. Highway runoff treatment by hybrid adsorptive media-baffled subsurface flow constructed wetland. *Ecological Engineering* 91: 231-239.
- Zheng, Y., Wang X., Dzakpasu M., Zhao Y., Ngo H. H., Guo W., Ge Y. and Xiong J. 2016. Effects of interspecific competition on the growth of macrophytes and nutrient removal in constructed wetlands: A comparative assessment of free water surface and horizontal subsurface flow systems. *Bioresour Technol* 207: 134-141.
- Zhou, L. J., Zhang B. B., Zhao Y. G. and Wu Q. L. 2016. Occurrence, spatiotemporal distribution, and ecological risks of steroids in a large shallow Chinese lake, Lake Taihu. *Sci Total Environ* 557-558: 68-79.
- Zoeller, R. T., Brown T. R., Doan L. L., Gore A. C., Skakkebaek N. E., Soto A. M., Woodruff T. J. and Vom Saal F. S. 2012. Endocrine-disrupting chemicals and public health protection: a statement of principles from The Endocrine Society. *Endocrinology* 153(9): 4097-4110.
- Zucchi, S., Castiglioni S. and Fent K. 2012. Progestins and antiprogestins affect gene expression in early development in zebrafish (*Danio rerio*) at environmental concentrations. *Environ Sci Technol* 46(9): 5183-5192.