

**SUBSTRATE PREFERENCES AND SEASONAL
VARIATION OF CIGUATERA FISH
POISONING DINOFLAGELLATES IN
KOTA KINABALU COASTAL WATERS**

STEFFIANA J JIPANIN

**THESIS SUBMITTED IN FULFILLMENT FOR
THE DEGREE OF MASTER OF SCIENCE**

**PERPUSTAKAAN
UNIVERSITI MALAYSIA SABAH**

**BORNEO MARINE RESEARCH INSTITUTE
UNIVERSITI MALAYSIA SABAH
2019**



UMS
UNIVERSITI MALAYSIA SABAH

UNIVERSITI MALAYSIA SABAH

BORANG PENGESAHAN STATUS TESIS

JUDUL: **SUBSTRATE PREFERENCES AND SEASONAL VARIATION OF CIGUATERA FISH POISONING DINOFLAGELLATES IN KOTA KINABALU COASTAL WATERS**

IJAZAH: **SARJANA SAINS (SAINS MARIN)**

Saya **STEFFIANA J JIPANIN**, Sesi **2016-2019**, mengaku membenarkan tesis Sarjana ini disimpan di Perpustakaan Universiti Malaysia Sabah dengan syarat-syarat kegunaan seperti berikut:-

1. Tesis ini adalah hak milik Universiti Malaysia Sabah
2. Perpustakaan Universiti Malaysia Sabah dibenarkan membuat salinan untuk tujuan pengajian sahaja.
3. Perpustakaan dibenarkan membuat salinan tesis ini sebagai bahan pertukaran antara institusi pengajian tinggi.
4. Sila tandakan (/):

SULIT

(Mengandungi maklumat yang berdarjah keselamatan atau kepentingan Malaysia seperti yang termaktub di dalam AKTA RAHSIA 1972)

TERHAD

(Mengandungi maklumat TERHAD yang telah ditentukan oleh organisasi/badan di mana penyelidikan dijalankan)

TIDAK TERHAD



Disahkan Oleh,


NURAZLYNNE MOHD. JOHAN @ JAYLYNE
PUSTAKAWAN

UNIVERSITI MALAYSIA SABAH
(Tandatangan Pustakawan)

STEFFIANA J JIPANIN
MY1611003T



Tarikh : 27 September 2019
(Prof. Madya Dr. Sitti Raehanah binti Muhd Shaleh)
Penyelia Utama


(Prof. Madya Dr. Lim Po Teen)
Penyelia Bersama



UMS
UNIVERSITI MALAYSIA SABAH

DECLARATION

I hereby declare that the material in this thesis is my own except for quotations, equations, summaries and references, which have been duly acknowledged and that it has not been submitted for any other degree or professional qualification.

26 September 2019



Steffiana J Jipanin

MY1611003T



CERTIFICATION

NAME : **STEFFIANA J JIPANIN**

MATRIC NO. : **MY1611003T**

TITLE : **SUBSTRATE PREFERENCES AND SEASONAL VARIATION OF CIGUATERA FISH POISONING DINOFLAGELLATES IN KOTA KINABALU COASTAL WATERS**

NAME OF DEGREE : **MASTER OF SCIENCE
(MARINE SCIENCE)**


DATE OF VIVA : **27th AUGUST 2019**

CERTIFIED BY

1. MAIN SUPERVISOR

Assoc. Prof. Dr. Sitti Raehanah binti Muhd Shaleh

Signature



2. CO-SUPERVISOR

Assoc. Prof. Dr. Lim Po Teen



ACKNOWLEDGEMENT

To be honest, my research would not be an interesting one if it is not for them, my lovely and full of passion supervisors, Assoc. Prof. Dr. Sitti Raehanah, Assoc. Prof. Dr. Lim Po Teen and the most caring lecturer Dr. Leaw Chui Pin for all their kind advices, guidance and support in this research work that lead to the completion of this thesis. Without their valuable teachings, I would not be able to complete my master's study. To Dr. Faihanna Ching Abdullah who is the one I constantly whine to whenever I feel like giving up, thank you for not giving up on me. I wish to thank God and my Guardian Angel for always blessing me with wisdom and perseverance throughout my research by whispering to me and listening to my prayers. My beloved parents, Mary and Jinius, my mother in law, Josephine, and my ever-supportive husband, Darryl, you are very understanding and thoughtful by helping me taking care of the children when I needed the time to focus on my research and write-ups. My lovely children Sumi, Zio and Vava who are my main inspiration for me to complete my studies on time. Mr. Ronnie, Mr. Calvin and Mr. Asmarul who have lend their valuable time and effort in assisting me to collect samples. My wonderful superiors Dr. Shuhadah Mustapha who is very understanding and concern about my research, Dr. Ahemad Sade our director who encourages us to further our studies. Friends and colleagues at Likas Fisheries Complex, Ms. Helena, Ms. Laura, Ms. Dorothy and Ms. Dayang Siti Maryam who always encouraging me when I feel miserable and constantly reminding me to find the fun in doing research even though I always cried my lungs out. Not forgetting Mr. Yusdi Ismail and staff of Borneo Marine Research Institute for their assistance in laboratory works. Wonderful and brilliant master students at Bachok Marine Research Station Mr. Lee Li Keat, Mr. Lim Zhen Fei, Ms. Tan Suh Nih, Ms. Hui Huey Er, Ms. Winnie Lau and Ms. Nurin Izzati Mustapa who openly shared their knowledge and experience in benthic harmful algal bloom studies and interns who have made laboratory works easier. Special thanks to Biasiswa Kerajaan Negeri Sabah for supporting part of my tuition fees. Thank you and God bless.

- Steffi -

26 September 2019



ABSTRACT

Ciguatera fish poisoning (CFP) is a food poisoning endemic to tropical and subtropical regions caused by eating fish that have been contaminated with toxins produced by dinoflagellates mainly of the genus *Gambierdiscus*. The toxin may be found in large reef fish, generally barracuda, red snapper, eel, grouper, sea bass, amberjack and Spanish mackerel. It is the most common seafood poisoning affecting the human population worldwide with 50,000 to 500,000 global incidences annually. Since 1984, there were a few suspected cases of CFP in Sabah, but these were never confirmed. This study aims to identify the CFP species with emphasis to the distribution and substrate preferences as well as the seasonal effects on the cell abundance of CFP dinoflagellates in the reefs of Kota Kinabalu, Sabah, Malaysia. Sepanggar Island, Manukan Island and Gaya Island were chosen as the sampling sites. Sampling duration was done in 12 months which involved 24 diving trips in total. A total of 149 fiberglass screens were used as an artificial substrate to collect the benthic epiphytic-CFP associated microalgae for 24 hours. Samples were then brought back to the laboratory for sample processing and cell enumeration. Photoquadrat survey method was adapted for habitat mapping by taking photographs underwater using a 0.25 m² quadrat at the sites of where devices were deployed. Bray-Curtis similarity was used to construct cluster analysis based on relative abundance data and relationships between substrate types were visualized using cluster analysis. One-way ANOVA (Kruskal-Wallis test) and Spearman correlation tests were done using GraphPad Prism 5 (GraphPad Inc.) for data analysis. The results recorded all five genera of CFP causing dinoflagellates *Amphidinium*, *Coolia*, *Gambierdiscus*, *Ostreopsis* and *Prorocentrum* were found in coral reef marine environments in low abundances at Kota Kinabalu coastal waters. *Prorocentrum* was found dominant in Sepanggar Island (62.2%), while *Ostreopsis* dominates in Manukan Island (73.6%) and Gaya Island (75.7%). Four major distinct geomorphic zones (rubbles, sand, dead corals and macroalgae) were indicated from cluster analysis results. The results showed dinoflagellates in this study were epiphytic on dead corals, and low densities were detected on sands. Two identified CFP dinoflagellates species showed statistically significant substrate preferences. For instance, *Gambierdiscus* was mainly founded on sand, *Prorocentrum* mainly encountered at macroalgae, while *Ostreopsis*, *Coolia* and *Amphidinium* showed no preferences on the four main substrate types. Mean total abundance of CFP dinoflagellates varied significantly between seasons (Southwest Monsoon>Northeast Monsoon>Inter-Monsoon). Throughout the study, gradual increases of *Gambierdiscus* cell numbers were observed during Southwest Monsoon (SWM). A significant positive correlation was observed between *Gambierdiscus* cell and sea surface temperature. *Prorocentrum* cell numbers were stable between seasons but showed an increasing pattern during Northeast Monsoon (NEM). *Coolia* and *Ostreopsis* did not show any pattern related to seasonal changes but average cell abundances of both dinoflagellates were significantly different between SWM and Inter-Monsoon (IM), while *Amphidinium* cell numbers showed no significant difference between seasons. The results observed highest average abundances of total CFP dinoflagellates during SWM (65.22 ± 149.5 cells/ 100 cm²), while the lowest abundances were during IM. In conclusion, this study can be used as a reference in studying and managing benthic harmful algal bloom dinoflagellates by safeguarding public health in order to reduce seafood poisoning cases.



ABSTRAK

KEUTAMAAN SUBSTRAT DAN VARIASI MUSIM BAGI DINOFLAGELLAT KERACUNAN IKAN CIGUATERA DI PERAIRAN PANTAI KOTA KINABALU

Keracunan Ikan Ciguatera (CFP) adalah keracunan makanan endemik di kawasan tropika dan subtropika yang disebabkan oleh ikan yang tercemar dengan toksin yang dihasilkan oleh dinoflagellat genus *Gambierdiscus*. Toksin tersebut boleh dijumpai pada ikan karang bersaiz besar, seperti barakuda, kerapu, ikan merah, belut, "amberjack", ikan siakap dan ikan tenggiri Sepanyol. Ia merupakan keracunan makanan laut yang menjelaskan populasi manusia di seluruh dunia dengan jumlah insiden global sebanyak 50,000 hingga 500,000 setiap tahun. Sejak tahun 1984, terdapat beberapa kes yang disyaki CFP di Sabah, tetapi kes tersebut tidak pernah disahkan. Kajian ini bertujuan untuk mengenalpasti spesies CFP dengan menjurus pada taburan dan keutamaan substrat serta kesan musim ke atas kelimpahan sel dinoflagellat CFP di terumbu karang Kota Kinabalu, Sabah, Malaysia. Pulau Sepanggar, Pulau Manukan dan Pulau Gaya telah dipilih sebagai tapak kajian. Pensampelan telah dijalankan selama 12 bulan yang melibatkan 24 selam skuba. Sebanyak 149 skrin gentian kaca digunakan untuk mendapatkan dinoflagellat CFP selama 24 jam. Sampel kemudian dibawa balik ke makmal untuk pemprosesan sampel dan penghitungan sel. Kaedah 'photoquadrat' telah diaplikasi untuk pemetaan habitat dengan mengambil gambar menggunakan kuadrat bersaiz 0.25 m² di dalam air. 'Bray-Curtis similarity' digunakan untuk membina analisis kluster berdasarkan data kelimpahan relatif dan hubungan di antara jenis substrat didapatkan melalui analisis kluster. Semua data analisis dijalankan dengan kaedah ANOVA Sehala (Kruskal-Wallis test) dan 'Spearman correlation test' menggunakan GraphPad Prism 5 (GraphPad Inc.). Hasil kajian merekodkan semua lima genera dinoflagellat CFP *Amphidinium*, *Coolia*, *Gambierdiscus*, *Ostreopsis* and *Prorocentrum* ditemui di persekitaran terumbu karang marin dalam jumlah yang rendah di perairan pantai Kota Kinabalu. *Prorocentrum* didapati dominan di Pulau Sepanggar (62.2%), manakala *Ostreopsis* mendominasi di Pulau Manukan (73.6%) dan Pulau Gaya (75.7%). Hasil analisis kluster menunjukkan terdapat empat zon utama geomorfik yang berbeza (pecahan karang, pasir, karang mati dan makroalga). Hasil kajian mendapati dinoflagellat dalam kajian ini adalah epifit pada karang mati dan kepadatan rendah dikesan pada pasir. Dua spesies dinoflagellat CFP yang dikenal pasti menunjukkan keutamaan substrat yang signifikan. Sebagai contoh, *Gambierdiscus* ditemui pada pasir, *Prorocentrum* didapati di makroalga, manakala *Ostreopsis*, *Coolia* dan *Amphidinium* tidak menunjukkan keutamaan terhadap empat jenis substrat utama. Purata jumlah kelimpahan dinoflagellat CFP berubah secara signifikan antara musim (SWM> NEM> IM). Sepanjang kajian, bilangan *Gambierdiscus* didapati meningkat semasa SWM. Korelasi positif yang kuat diperhatikan antara sel *Gambierdiscus* dengan suhu permukaan laut. Bilangan sel *Prorocentrum* adalah stabil sepanjang musim tetapi menunjukkan corak peningkatan semasa NEM. *Coolia* dan *Ostreopsis* tidak menunjukkan corak yang berkaitan dengan perubahan musim tetapi purata kelimpahan sel bagi kedua-dua dinoflagellat antara SWM dan IM menunjukkan perbezaan signifikan. Bilangan sel *Amphidinium* tidak menunjukkan perbezaan yang signifikan di kalangan musim. Hasil kajian menunjukkan purata kelimpahan tertinggi jumlah dinoflagellat CFP yang direkodkan adalah semasa SWM (65.22 ± 149.5 sel / 100 cm²), manakala yang paling rendah adalah semasa IM. Sebagai kesimpulan, kajian ini boleh digunakan sebagai rujukan



dalam kajian dan pengurusan dinoflagellat ledakan alga bentik berbahaya dengan menjaga kesihatan masyarakat untuk mengurangkan kes-kes keracunan makanan laut.



TABLE OF CONTENTS

	Page
TITLE	i
DECLARATION	ii
CERTIFICATION	iii
ACKNOWLEDGEMENT	iv
ABSTRACT	v
ABSTRAK	vi
TABLE OF CONTENTS	viii
LIST OF TABLES	x
LIST OF FIGURES	xii
LIST OF SYMBOLS AND ABBREVIATIONS	xv
LIST OF APPENDICES	xvi
CHAPTER 1 : INTRODUCTION	
1.1 Overview of ciguatera fish poisoning	1
1.2 Problem statement	3
1.3 Significance of study	4
1.4 Objectives of study	4
CHAPTER 2 : LITERATURE REVIEW	
2.1 Harmful algal blooms in Malaysia	5
2.2 Ciguatera fish poisoning in Malaysia	7
2.3 Dinoflagellates	7
2.4 Ciguatera associated benthic dinoflagellate	8
2.5 Substrate preferences of benthic dinoflagellates	18
2.6 Seasonal factors	20



2.7 Government policy	22
-----------------------	----

CHAPTER 3 : MATERIALS AND METHODS

3.1 Study location	23
3.2 Sampling procedure	25
3.3 Cell observation	27
3.4 Cell enumeration of benthic dinoflagellates	27
3.5 Species identification	28
3.6 Habitat mapping and classification	32
3.7 Environmental parameter and rainfall data	35
3.8 Data analysis	35

CHAPTER 4 : RESULTS

4.1 CFP dinoflagellates of Sepanggar, Manukan and Gaya Islands	36
4.2 Habitat mapping and classification	38
4.3 Relative abundance and cell abundance of CFP dinoflagellates species	42
4.4 Substrate preferences of CFP dinoflagellates	48
4.5 Seasonal effects	52
4.6 Environmental data of Sepanggar, Manukan dan Gaya Islands	56

CHAPTER 5 : DISCUSSION

5.1 CFP dinoflagellates of Sepanggar, Manukan and Gaya Islands	58
5.2 Abundance of CFP Dinoflagellates Species	59
5.3 Substrate preferences	62
5.4 Seasonal effects	65

CHAPTER 6 : CONCLUSION

69

REFERENCES

71

APPENDICES

79



LIST OF TABLES

	Page
Table 2.1: Toxins Produced by Benthic Coral Reef-Associated Dinoflagellates	10
Table 2.2: Summary of studies on substrate preferences	18
Table 3.1: Oligonucleotide primers used for amplification of ITS and 28S rDNA regions	30
Table 3.2: PCR amplification conditions for ITS and LSU rDNA respectively	30
Table 4.1: Benthic biological and physical substratum of Sepanggar, Manukan and Gaya Islands were briefly explained and classified into four major distinct geomorphic zones (Cluster I – IV). Variation of benthic substrata was denoted as type A – K to clearly differentiate their surface features within bottom marine reef environment	40
Table 4.2: CFP dinoflagellates assemblage composition obtained from sampling sites: Average cell composition (cells/100 cm ² ± SD) of each genus; relative abundance (%)	44
Table 4.3: Abundance (cells/100 cm ²) of CFP dinoflagellates species of interest at each classified microhabitat speciation (Cluster I -IV) from June 2017 to May 2018. Values were reported as average ± standard deviation; Average with different letters were significantly different (Kruskal-Wallis test, <i>p</i> -value < 0.05)	50
Table 4.4: Abundance (cells/100 cm ²) of CFP dinoflagellates species of interest at each different season from June 2017 to May 2018. Values were reported as average ± standard deviation; Average with different letters were significantly different (Kruskal-Wallis test, <i>p</i> -value < 0.05)	53
Table 4.5: Details of the environmental data taken recorded for each season from Sepanggar, Manukan and Gaya Islands showing the sampling date, sea surface salinity (PSS), pH, temperature (°C) and dissolved oxygen (mg/L)	57



Table 4.6: Spearman's rank correlation (p -value < 0.05) coefficients 57
between salinity, dissolved oxygen, pH, temperature, rainfall and
total CFP dinoflagellates as well as each genera of CFP
dinoflagellates in coastal waters of Kota Kinabalu, Sabah



LIST OF FIGURES

	Page
Figure 2.1: Morphology of <i>Gambierdiscus</i> adapted from other studies	9
Figure 2.2: Flattened and tear shaped <i>Ostreopsis</i> cells with many golden-brown chloroplasts	12
Figure 2.3: (A) Schematic drawing of a hypothetical <i>Procentrum</i> cell showing the prorocentroid tabulation type, cell orientation convention and naming of morphological features. (B) Desmokont type of flagellation	14
Figure 2.4: Scanning electron micrographs of <i>C. malayensis</i> in (A) dorsal view, (B) epithelial view, (C) ventral/apical view, and (D) ventral/antapical view	16
Figure 2.5: Morphology of <i>Amphidinium carterae</i> cells (light microscopy): A -motile vegetative cells in culture; B -immotile cells after centrifugation. Scale bars: 10 µm	17
Figure 3.1: (A) Map of the three sampling locations in Kota Kinabalu, Sabah (adapted from Google Earth Pro); (B) Gaya Island-GI, (C) Sepanggar Island - SI, (D) Manukan Island-MI; the sampling sites represented with the black coloured circles	24
Figure 3.2: Artificial sampling device that was placed underwater for hours to collect benthic dinoflagellates	26
Figure 3.3: Determination of actual screen surface area considering series of cylindrical structured screen filaments (adopted from Weisstein, 2013)	28
Figure 3.4: Photoquadrat image uploaded in CoralNet with 100 points annotation	33
Figure 3.5: (A) Artificial sampling devices were inserted carefully into a bottle underwater. (B) Photoquadrat survey conducted using PVC frame photoquadrat with area of 0.25 m ² (50 cm x 50 cm) to determine bottom biota and physical substratum coverage in percentage cover (%)	34



- Figure 4.1: Five main genera of CFP dinoflagellates obtained from Sepanggar, Manukan and Gaya Islands were observed under LM. (A) *Amphidinium* sp., (B) *Coolia* sp., (C) *Gambierdiscus* sp., (D) *Ostreopsis* sp., (E) *Prorocentrum lima*, (F) *Prorocentrum* sp1., (G) *Prorocentrum* sp2., (H) *Prorocentrum* sp3. (Scale bar = 10 μm) 37
- Figure 4.2: Clustering dendrogram showing the dissimilarities relationship of each sites based on benthic biological and physical substratum characters cover (Types A – K), separating into four distinguishable geomorphic zones (Cluster I- IV) 41
- Figure 4.3: Relative abundance of CFP dinoflagellates (in percentage, %). *Ostreopsis* abundance represented 59.4% of relative abundance of CFP dinoflagellates assemblages, followed by *Prorocentrum* (31.0%). Less species assemblages were encountered on *Gambierdiscus* (4.1%), *Coolia* (3.6%) and *Amphidinium* (1.9%) 45
- Figure 4.4: Relative abundance of BHAB compositions (in percentage, %) according to each sampling sites 46
- Figure 4.5: Species composition (cells/100 cm²) of CFP dinoflagellates in relation to variation of bottom microhabitats. Dotted line represented four different substrate preferences (Cluster I - IV); Labels of sampling locations: S - Sepanggar Island, M- Manukan Island, G – Gaya Island 47
- Figure 4.6: Tukey plot of each and total CFP dinoflagellates in relation to classified microhabitat speciation (Cluster I - IV). Alphabet indicate significant difference (Kruskal-Wallis test, p -value < 0.05); horizontal line in box showed median; Box ends at the quartiles Q1 and Q3. Whiskers showed the upper and lower extreme; Dots represents outliers 51
- Figure 4.7: Total monthly rainfall (mm) and total CFP dinoflagellates (cells/ 100 cm²) from June 2017 to May 2018 in Kota Kinabalu 54

Figure 4.8: Temporal distribution of CFP dinoflagellate species at 55 Sepanggar, Manukan and Gaya Island sampling sites (cells / 100 cm²). (A) *Gambierdiscus*, (B) *Ostreopsis*, (C) *Coolia*, (D) *Prorocentrum* and (E) *Amphidinium*



LIST OF SYMBOLS AND ABBREVIATIONS

BHAB	:	Benthic harmful algal bloom
CFP	:	Ciguatera fish poisoning
CTX	:	Ciguatoxin
DO	:	Dissolved oxygen
DSP	:	Diarrheic Shellfish Poisoning
DTX	:	Dinophysistoxin
HAB	:	Harmful algal bloom
IM	:	Inter-Monsoon
MOH	:	Ministry of Health
MTX	:	Maitotoxin
NEM	:	Northeast Monsoon
OA	:	Okadaic acid
PAST	:	Paleontological statistics software package
PSP	:	Paralytic Shellfish Poisoning
SWM	:	Southwest Monsoon



LIST OF APPENDICES

	Page
Appendix A: Percentage covers of photoquadrat images extracted from CoralNet software	84
Appendix B: Environmental Parameters in Kota Kinabalu coastal waters from July 2017 to May 2018	89
Appendix C: Average Wind Velocity in Kota Kinabalu from June 2017 to May 2018	91



CHAPTER 1

INTRODUCTION

1.1 Overview of ciguatera fish poisoning

During the Spanish Conquest, there was an occurrence of a food poisoning caused by consumption of a marine snail, *Turbo pica* (termed cigua by the natives in Cuba) in the Caribbean and to illustrate this intoxication, a term ciguatera fish poisoning (CFP) was used (de Fouw *et al.*, 2001). Guzman-Perez *et al.*, (2000) stated that the word CFP is used today to name the intoxication instigated by consuming reef fish which have gained specific toxins from their diet. The fish usually originates from the tropical and subtropical areas of the Caribbean Sea and the Pacific Ocean.

CFP is a seafood illness endemic to tropical and subtropical coral reef areas of the world. Dinoflagellates of the genus *Gambierdiscus* produces lipid-soluble toxins, accumulates in finfish and consumption of certain reef fish causes intoxication (Dickey & Plakas, 2010). According to Mattei *et al.*, (2014), the most common seafood illness worldwide is ciguatera with 50,000 to 500,000 global occurrences per annum and it is mostly restricted to endemic areas which are located between 35° northern and 35° southern latitude.

de Fouw *et al.*, (2001) stated that a lot of countries such as Australia, Bahamas, Canada, China, Haiti, Hawaii, Madagascar, Mexico, Tonga, US, and even



UMS
UNIVERSITI MALAYSIA SABAH

in Germany and the Netherlands have reports of ciguatera intoxication. A theory made by Lehane & Lewis (2000) mentioned that ciguatera may develop at areas which are far from tropical oceans, where alive, fresh or preserved reef fish are exported to other areas and this might lead to a world health problem. Throughout the year, due to the expansion of international seafood trade from tropical fisheries worldwide, ciguatera has become a threat to consumers in non-endemic countries (Kibler *et al.*, 2015). Mattei *et al.*, (2014) suggested that the globalization of fishing industry and tourism in Europe and the lack of awareness among medical personnel which causes under-reporting are the other factors which contributed to the huge increase in the amount of ciguatera cases.

The causative agents of CFP in human are marine biotoxins produced by dinoflagellates (Mattei *et al.*, 2014). Ciguatera causes gastrointestinal and neurological disorders and may cause human death in rare cases (Faust, 2009). According to Xu *et al.*, (2014), *Gambierdiscus* produces gambiertoxins which are the precursors for the ciguatoxins (CTX) and are responsible for ciguatera. The toxins enter the coral reef food web through grazing by herbivores and detritivores, and when these animals are eaten by predators, accumulation and biomodification of these toxins take place. Coral-reefs are nurseries for harmful dinoflagellates and possible sites for ciguatera outbreak world-wide (Faust, 2009). According to Bienfang *et al.*, (2008), the dinoflagellates *Gambierdiscus* spp. produces CTX and these dinoflagellates are commonly found in association with numerous macroalgae in the coral reef ecosystems. Herbivorous fish eat up these dinoflagellates, initiates the bioaccumulation, biomagnification, and biomodification processes in the food web, as the herbivores are eaten by carnivores and, ends up as human consumption.

The symptoms of ciguatera seem to differ geographically, between individuals and incidents and differ temporally within an area. It mostly consists of early-onset (two to six hours) gastrointestinal disturbance, such as nausea, vomiting, and diarrhea, and may be followed by other neurological illnesses (18 hours), including numbness, reversal of temperature sensation, muscle and joint aches, itching,

tachycardia, blurred vision and paralysis. On some rare occasions, ciguatera may cause death (Botana, 2000).

Unfortunately, due to under-reporting and misdiagnosis, the real extent of illness is not well documented (Radke *et al.*, 2015). Berdalet *et al.*, (2015) stated that the cases of CFP often have been traditionally ‘managed’ by native fishermen via their local and traditional knowledge. In tropical areas, CFP is a threat to public health, and it the most widespread, and hence best described, in the island population of the Pacific and the Caribbean. There has been a consistent increase in the number of known harmful species, algal toxins, toxic outbreaks, and fisheries and ecosystem impacts, all at a time when human dependence on the coastal area for food, recreation, and commerce is rapidly growing (Anderson, 1998).

1.2 Problem statement

Department of Fisheries Sabah has been doing monitoring on CFP and based on their unpublished reports from 1984 to 1992, there have been 38 complaints regarding fish poisoning mainly due to red snappers, particularly *Lutjanus bohar*. Most of the samples were collected from the Kota Kinabalu Fish Market, SAFMA Fish Landing Jetty and several places on the West Coast of Sabah. In April 1989, red snapper *Lutjanus bohar* was totally banned either for local or overseas market to safeguard public health and fisheries industry in Sabah. In year 2017, an unpublished report by Ministry of Health (MOH) Sabah stated that there were three suspected cases of CFP. In December of the same year, one person was confirmed by laboratory test and medical practitioner as a case of CFP. All these reports raised a lot of questions such as the origin of the sample and the toxin itself. How were the CFP dinoflagellates distributed in Kota Kinabalu throughout the years? Up until now, there were lack of information on where and what is the CFP pattern in Sabah. Thus, the purpose of this study was to (1) check the presence of other CFP dinoflagellates in coastal waters of Kota Kinabalu, (2) find out if the CFP dinoflagellates show any substrate



preferences and (3) observe if seasonal factors have any effects on the cell abundance of CFP dinoflagellates.

1.3 Significance of study

Seafood poisoning cases in the west coast of Sabah has started since 1970s. To date, monitoring of paralytic shellfish poisoning by Department of Fisheries Sabah (DOFS) was successful in lowering the risk of seafood poisoning. By knowing the pattern and distribution of CFP dinoflagellates, monitoring plan can be carried out effectively in HAB management programs. Therefore, this study can be used as a reference in studying and managing BHAB dinoflagellates by safeguarding public health.

1.4 Objectives of study

The objectives of this study are:

- i. To identify the causative dinoflagellates species responsible to Ciguatera Fish Poisoning (CFP).
- ii. To determine the substrate preferences of CFP Dinoflagellates in Gaya, Manukan and Sepanggar Islands.
- iii. To determine the effects of seasonal factors on cell abundance of CFP Dinoflagellates.



CHAPTER 2

LITERATURE REVIEW

2.1 Harmful algal blooms in Malaysia

Harmful algal blooms (HABs) or commonly known as “red tides” are a worldwide problem which affects not only to human but also to our natural ecosystem. HABs can be categorized to three different types which are (1) dinoflagellate species that produce harmless water discolorations but cause mass mortality of fish due to oxygen depletion, (2) toxic to humans which causes gastrointestinal and neurological illnesses and (3) non-toxic to humans but harmful to fish by clogging their gills (especially fish in cage aquaculture).

The first report of HABs and shellfish toxicity in Malaysia was in 1976 when the *Pyrodinium bahamense* var. *compressum* bloomed in Brunei Bay on the west coast of Sabah and this event poisoned several people (Roy, 1977). The bloom eventually spread to other parts of Sabah but focuses mainly at the west coast of Sabah. Over the following few decades, many poisoning cases have been reported including fatality but with the establishment of monitoring programme, these cases were under control.



In Sabah, HABs caused by *P. bahamense* and *Cochlodinium polykrikoides* took place almost every year on the west coast of Sabah (Wang *et al.*, 2008), especially in coastal areas of Kota Kinabalu. Many poisoning cases have been reported over the years including casualty. Other unidentified toxic dinoflagellates may exist in waters of Sabah and current environmental conditions are not favorable for bloom to happen. The number of HAB species in Sabah are on the increase with toxic *Gymnodinium catenatum* identified off the west coast of Sabah (Mohammad-Noor *et al.*, 2008), *Gonyaulax polygramma* in Kota Kinabalu (Lim *et al.*, 2014) and *Noctiluca scintillans* in Kota Marudu and Kudat (Department of Fisheries Sabah).

Previous research concluded that the waters in Malaysia have a high diversity of marine benthic dinoflagellates flora and the highest species diversity among the sites examined was Sipadan Island (Mohammad-Noor *et al.*, 2007). A survey conducted in Singaporean waters showed that species diversity was low, with *O. ovata* was the dominant species (Holmes *et al.*, 1998). *Gambierdiscus yasumotoi*, a new toxin-producing species was identified in the reef around the Singapore island of Pulau Hantu (Holmes, 1998). In Malaysia waters, several species of benthic dinoflagellates which may be involved in CFP have been identified, they were *G. toxicus*, *O. ovata*, *O. lenticularis*, and *Coolia* sp. Toxicity screening based on hemolytic assay, antimicrobial and mouse bioassay has proven the existence of biological activities in aqueous and organic phase extract of these cultures. However, the toxins compound is yet need to be determined (Lim *et al.*, 2003). Department of Fisheries Sabah took the initiative to conduct a preliminary study on distribution of CFP dinoflagellates in Sabah between 1994 – 1997 with the help of a consultant from the Asean-Canada Cooperative Programme and the results showed that Sepanggar Island, Kota Kinabalu recorded the highest number of *G. toxicus* cells (383 cells/ g FW) (Department of Fisheries Sabah). To date, there were no reported blooms caused by benthic dinoflagellates in coastal waters of Sabah.

2.2 Ciguatera fish poisoning in Malaysia

In Malaysia, based from a report by annual local health conference in 2011, ciguatera was first reported in April 2010 where 6 subjects were hospitalized after consuming red snapper in Keningau, Sabah (Mohd *et al.*, 2011). In September 2010, 22 members from five families at Jeli, Kelantan were involved and red snapper was the cause of these outbreaks according to a review done by Chan (2015). According to Lim *et al.*, (2003), there was an outbreak of CFP in Hong Kong which claimed that the toxic parrot fishes were originated from Southeast Asia although the country was not specified. Screening was carried out using CTX immunological assay and it shows positive results on reef fishes from west coast of Sabah.

An unpublished report by Ministry of Health (MOH) Sabah in year 2017 stated that there were three suspected cases of CFP in April and May. According to their report, both patients had consumed red snapper, *Lutjanus* sp. commonly known as 'Ikan tambak' and 'Ikan merah'. They suffered from diarrhea, neurological symptoms and numbness. However, no samples were retrieved from victims to conduct analysis for confirmation. *Lutjanus* sp. is listed in reports of ciguatera poisoning (Lewis, 1986). In December of the same year, one person was hospitalized and confirmed by the medical practitioner as a case of CFP via clinical symptoms such as numbness, itching, diarrhea and reversal of hot and cold sensation. Samples were sent to Japan for CTX analysis and shows positive result (presence of ciguatoxin, P-CTX-1B). The positive samples were identified as emperor snapper, *Lutjanus sebae* commonly known as 'Ikan merah'. Preliminary investigation found out that the fish was bought in Lido Fishmarket, Kota Kinabalu, Sabah. However, the origin of fish was yet to be confirmed (Department of Fisheries Sabah).

2.3 Dinoflagellates

According to Gaines & Elbrächter (1987), dinoflagellates are natural organisms in all types of aquatic ecosystems. Some of the species in the group are considered



REFERENCES

- Accoroni, S., Romagnoli, T., Colombo, F., Pennesi, C., Di Camillo, C. G., Marini, M., Battocchi, C., Ciminiello, P., Dell'Aversano, C., Dello Iacovo, E., Fattorusso, E., Tartaglione, L., Penna, A. & Totti, C. 2011. *Ostreopsis cf. ovata* bloom in the northern Adriatic Sea during summer 2009: ecology, molecular characterization and toxin profile. *Mar. Pollut. Bull.* 62, 2512-2519.
- Accoroni, S., Glibert, P. M., Pichierri, S., Romagnoli, T., Marini, M. & Totti, C. 2015. A conceptual model of annual *Ostreopsis cf. ovata* blooms in the northern Adriatic Sea based on the synergic effects of hydrodynamics, temperature, and the N:P ratio of water column nutrients. *Harmful Algae*. 45, 14–25.
- Adachi, R. & Fukuyo, Y. 1979. The thecal structure of a marine toxic dinoflagellate *Gambierdiscus toxicus* gen. et sp. nov. collected in a ciguatera-endemic area. *Bull. Jap. Soc. Sci. Fish.* 45, 67–71.
- Adachi, M., Sake, Y. & Ishida, Y. 1996. Analysis of *Alexandrium* (Dinophyceae) species using sequences of the 5.8S ribosomal DNA and internal transcribed spacer regions. *Journal of Phycology*. 32(3), 424-432.
- Aligizaki, K. & Nikolaidis, G. 2006. The presence of the potentially toxic genera *Ostreopsis* and *Coolia* (Dinophyceae) in the North Aegean Sea, Greece. *Harmful Algae*. 5, 717-730.
- Aligizaki, K., Nikolaidis, G., Katikou, P., Baxevanis, A. D. & Abatzopoulos, T. J. 2009. Potentially toxic epiphytic *Prorocentrum* (Dinophyceae) species in Greek coastal waters. *Harmful Algae*. 8, 299–311.
- Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. 1990. Basic local alignment search tool. *Journal of Molecular Biology*. 215(3), 403-410.
- Anderson, D. M. 1998. Physiological ecology of harmful algal blooms. In Anderson, D. M., Cembella, A. D. & Hallegraeff, G. M. (eds.). *Proceedings of the NATO advanced Study Institute on The Physiological Ecology of Harmful Algal Blooms*. Bermuda Biological Station for Research, Bermuda, USA. May 27-June 6, 1996. Germany: Springer-Verlag Berlin Heidelberg.
- Armi, Z. & Turki, S. 2010. First recorded proliferation of *Coolia monotis* (Meunier, 1919) in the North Lake of Tunis (Tunisia) correlation with environmental factors. *Environ. Monit Assess.* 164, 423–433.
- Ateweberhan, M., Bruggemann, J. H. & Breeman, A. M. 2005. Seasonal dynamics of *Sargassum ilicifolium* (Phaeophyta) on a shallow reef flat in the southern Red Sea (Eritrea). *Mar. Ecol. Prog. Ser.* 292, 159–171.
- Baig, H. S., Saifullah, S. M., & Dar, A. 2006. Occurrence and toxicity of *Amphidinium carterae* Hulbert in the North Arabian Sea. *Harmful Algae*. 5, 133-140.

- Ballantine, D. L., Tosteson, T. R. & Bardales, A. T. 1988. Population dynamics and toxicity of natural populations of benthic dinoflagellates in Southwest Puerto Rico. *Journal of Experimental Marine Biology and Ecology*. 119(3), 201-212.
- Barone, R. 2007. Behavioural trait of *Ostreopsis ovata* (Dinophyceae) in Mediterranean rock pools: the spider's strategy. *Harmful Algae News*. 33, 1.
- Battocchi, C., Totti, C., Vila, M., Masó, M., Capellacci, S., Accoroni, S., Reñé, A., Scardi, M. & Penna, A. 2010. Monitoring toxic microalgae *Ostreopsis* (dinoflagellate) species in coastal waters of the Mediterranean Sea using molecular PCR-based assay combined with light microscopy. *Mar. Pollut. Bull.* 60, 1074-1084.
- Berdalet, E., Fleming, L. E., Gowen, R., Davidson, K., Hess, P., Backer, L. C., Moore, S. K., Hoagland, P. & Enevoldsen, H. 2015. Marine Harmful Algal Blooms, Human Health and Wellbeing: Challenges and Opportunities in the 21st Century. *J. Mar. Biol. Assoc. U.K.* 1, 61-91.
- Berdieva, M., Safonov, P. & Matantseva, O. 2019. Ultrastructural aspects of ecdysis in the naked dinoflagellate *Amphidinium carterae*. *Protistology*. 13(2), 57-63.
- Besada, E. G., Loeblich, L. A. & Loeblich III, A. R. 1982. Observations on Tropical, Benthic Dinoflagellates from Ciguatera-endemic Areas: *Coolia*, *Gambierdiscus* and *Ostreopsis*. *Bulletin of Marine Science*. 32(3), 723-735.
- Bianco, I., Sangiorgi, V., Penna, A., Guerrini, F., Pistocchi, R., Zaottini, E. & Congestri, R. 2007. *Ostreopsis ovata* in benthic aggregates along the Latium Coast (middle Tyrrhenian Sea), p. 29 (abstract). *Proc. Int. Symp. On Algal Toxins*. Trieste, Italy.
- Bienfang, P. K., Parsons, M. L., Bidigare, R. R., Laws E. A. & Moeller, P. D. R. 2008. Chapter 4. Ciguatera Fish Poisoning: A Synopsis from Ecology to Toxicity. In Walsh, P. J., Smith, S. L., Fleming, L. E., Solo-Gabriele, H. M. & Gerwick, W. H. (eds.). *Oceans and Human Health: Risks and Remedies from the Seas*, pp. 257-270. USA: Elsevier Inc.
- Bomber, J. W., Norris, D. R. & Mitchell, L. E. 1985. *Benthic dinoflagellates associated with Ciguatera from the Florida Keys. II. Temporal, spatial and substrate heterogeneity of Prorocentrum lima*. New York: Elsevier Science Publishing.
- Bomber, J. W. & Aikman, K. E. 1989. The ciguatera dinoflagellates. *Biological Oceanography*. 6, 291-311.
- Botana, L. M. 2000. *Seafood and freshwater toxins: Pharmacology, physiology, and detection*. New York: Marcel Dekker, Inc.
- Brescianini, C., Grillo, C., Melchiorre, N., Bertolotto, R., Ferrari, A., Vivaldi, B., Icardi, G., Gramaccioni, L., Funari, E. & Scardala, S. 2006. *Ostreopsis ovata* algal blooms affecting human health in Genova, Italy, 2005 and 2006. *Euro Surveill.* 11(36), 3040.
- Carlson, R. D. 1984. *Distribution, periodicity, and culture of benthic/epiphytic*

- dinoflagellates in a ciguatera endemic region of the Caribbean.* PhD thesis. Carbondale: Southern Illinois University. (Unpublished).
- Carlson, R. D. & Tindall, D. R. 1985. Distribution and periodicity of toxic dinoflagellates in the Virgin Islands. In Anderson, D. M., White, A. W. & Baden, D. G. (eds.). *Toxic dinoflagellates*, pp. 171–176. New York: Elsevier Science.
- Castro, P. & Huber, M. E. 2010. *Marine Biology* 8th ed. p. 95. New York: McGraw Hill.
- Catania, D. 2017. *The influence of macroalgae on the proliferation and regulation of the benthic dinoflagellate *Ostreopsis cf. ovata* blooms.* PhD thesis, Agricultural sciences: Université Côte d'Azur. (Unpublished).
- Chan, T. Y. K. 2015. Ciguatera Fish Poisoning in East Asia and Southeast Asia. *Mar. Drugs.* 13, 3466-3478.
- Chang, F. H., Shimizu, Y., Hay, B., Stewart, R., Mackay, G. & Tasker, R. 2000. Three recently recorded *Ostreopsis* spp. (Dinophyceae) in New Zealand: Temporal and regional distribution in the upper North Island from 1995 to 1997. *New Zealand Journal of Marine and Freshwater Research.* 34(1), 29-39.
- Cembella, A. D. 2003. Chemical ecology of eukaryotic microalgae in marine ecosystems. *Phycologia.* 42, 420-447.
- Chinain, M., Germain, M., Deparis, X., Pauillac, S. & Legrand A. M. 1999a. Seasonal abundance and toxicity of the dinoflagellate *Gambierdiscus* spp. (Dinophyceae), the causative agent of ciguatera in Tahiti, French Polynesia. *Marine Biology.* 135, 259-267.
- Chinain, M., Faust, M. A. & Pauillac, S. 1999b. Morphology and Molecular Analyses of Three Toxic Species of *Gambierdiscus* (Dinophyceae): *G. pacificus* sp. nov., *G. australis* sp. nov., and *G. polynesiensis* sp. nov. *J. Phycol.* 35, 1282-1296.
- Cohu, S., Thibaut, T., Mangialajo, L., Labat, J. P., Passafiume, O., Blanfuné, A., Simon, N., Cottalorda, J. M. & Lemée, R. 2011. Occurrence of the toxic dinoflagellate *Ostreopsis cf. ovata* in relation with environmental factors in Monaco (NW Mediterranean). *Mar. Pollut. Bull.* 62, 2681-2691.
- Cohu, S., Mangialajo, L., Thibaut, T., Blanfuné, A., Marro, S. & Lemée, R. 2013. Proliferation of the toxic dinoflagellate *Ostreopsis cf. ovata* in relation to depth, biotic substrate and environmental factors in the North West Mediterranean Sea. *Harmful Algae.* 24, 32–44.
- Cruz-Rivera, E. & Villareal, T. A. 2006. Macroalgal palatability and the flux of ciguatera toxins through marine food webs. *Harmful Algae.* 5, 497–525.
- de Fouw, J.C., van Egmond, H.P. & Speijers, G.J.A. 2001. *Ciguatera Fish Poisoning: A Review.* RIVM report (February), 1–66.
- de Sylva, D. P. 1994. Distribution and ecology of ciguatera fish poisoning in Florida, with emphasis on the Florida Keys. *Bulletin of Marine Science.* 54(3), 944-954.



- Delgado, G., Lechuga-Devéze, C. H., Popowski, G., Troccoli, L. & Salinas, C. D. 2006. Epiphytic dinoflagellates associated with ciguatera in the northwestern coast of Cuba. *Revista de Biología Tropical*. 54, 299-310.
- Dickey, R. W. & Plakas, S. M. 2010. Ciguatera : A Public Health Perspective. *Toxicon*. 56, 123-36.
- Echigoya, R., Rhodes, L., Oshima, Y. & Satake, M. 2005. The structures of five new antifungal and hemolytic amphidinol analogs from *Amphidinium carterae* collected in New Zealand. *Harmful Algae*. 4, 383-389.
- Faust, M. A. 1995. Observation of sand-dwelling toxic dinoflagellates (Dinophyceae) from widely differing sites, including two new species. *J. Phycol.* 31(6), 996-1003.
- Faust, M. A. & Morton, S. L. 1995. Morphology and ecology of the marine dinoflagellate *Ostreopsis labens* sp. nov. (Dinophyceae). *J. Phycol.* 31, 456-463.
- Faust, M. A., Larsen, J. & Moestrup, O. 1999. *Potentially toxic phytoplankton genus Prorocentrum*. ICES identification leaflets for plankton, Copenhagen.
- Faust, M. A. 2009. Ciguatera-Causing Dinoflagellates in a Coral-Reef Mangrove Ecosystem, Belize. *Atoll Research Bulletin*. 569, 1-32.
- Foden, J., Purdie, D. A., Morris, S. & Nascimento, S. 2005. Epiphytic abundance and toxicity of *Prorocentrum lima* populations in the Fleet Lagoon, UK. *Harmful Algae*. 4, 1063-1074.
- Fukuyo, Y. 1981. Taxonomical study on benthic dinoflagellates collected in coral reefs. *Bull. Jap. Soc. Sci. Fish.* 47, 967-978.
- Gaines, G. & Elbrächter, M. 1987. Heterotrophic nutrition. In Taylor, F. J. R. (ed.) *The biology of dinoflagellates, Botanical Monographs Volume 21*. Oxford: Blackwell Scientific Publications.
- Guzman-Perez, S. E. & Park, D. L. 2000. Ciguatera Toxins: Chemistry and Detection. In Botana, L. M. (ed.). *Seafood and freshwater toxins: Pharmacology, physiology, and detection*, pp. 401-418. New York: Marcel Dekker, Inc.
- Hachani, M. A., Dhib, A., Fathalli, A., Ziadi, B., Turki, S. & Aleya, L. 2018. Harmful epiphytic dinoflagellate assemblages on macrophytes in the Gulf of Tunis. *Harmful Algae*. 77, 29-42.
- Hall, T. A. 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series*. 41, 95-98.
- Hammer, Ø., Harper, D. A. T. & Ryan, P. D. 2001. PAST: Paleontological statistics software package for education and data analysis. *Palaeontologia Electronica*. 4(1), 9.

- Hansen, G., Turquet, J., Quod, J-P., Ten-Hage, L., Lugomela, C., Kyewalyanga, M., Hurbungs, M., Wawiye, P., Ogongo, B., Tunje, S. & Rakotoarinjanahary, H. 2001. *Potentially harmful microalgae of the Western Indian Ocean, guide based on preliminary survey*. IOC Manuals and Guides No. 41. Intergovernmental Oceanic Commission of UNESCO, Paris. pp. 105.
- Holmes, M. J., Lewis, R. J., Jones, A. & Hoy, A. W. 1995. Cooliatoxin, the first toxin from *Coolia monotis* (Dinophyceae). *Natural Toxins*. 3(5), 355-362.
- Holmes, M. J. 1998. *Gambierdiscus yasumotoi* sp. nov. (Dinophyceae), a toxic benthic dinoflagellate from Southeastern Asia. *J. Phycol.* 34, 661-668.
- Holmes, M. J., Lee, F. C., Teo, S. L. M. & Khoo, H. W. 1998. A survey of benthic dinoflagellates on Singapore reefs. In Reguera, B., Blanco, J., Fernández, M.L. & Wyatt, T. (eds.). *Harmful Algae*, pp. 50-51. Santiago de Compostela: Xunta de Galicia and Intergovernmental Oceanographic Commission of UNESCO.
- Hoppenrath, M., Chomérat, N., Horiguchi, T., Schweikert, M., Nagahama, Y. & Murray, S. 2013. Taxonomy and phylogeny of the benthic *Prorocentrum* species (Dinophyceae)—A proposal and review. *Harmful Algae*. 27, 1-28.
- Hoshaw, R. W. & Rosowski, J. R. 1973. Methods for microscopic algae. In Stein, J.R. (ed.). *Handbook of Phycological Methods: Culture Methods and Growth Measurements*, pp. 53-68. Cambridge: Cambridge University Press.
- Hu, T., deFreitas, A. S., Curtis, J. M., Oshima, Y., Walter, J. A. & Wright, J. L. 1996. Isolation and structure of prorocentrolide B, a fast-acting toxin from *Prorocentrum maculosum*. *J Nat Prod.* 59(11), 1010-4.
- Illoul, H., Hernandez, F. R., Vila, M., Adjas, N., Younes, A. A., Bournissa, M., Koroghli, A., Marouf, N., Rabia, S. & Ameur, F. L. K. 2012. The genus *Ostreopsis* along the Algerian coastal waters (SW Mediterranean Sea) associated with a human respiratory intoxication episode. *Cryptogam. Algol.* 33, 209-216.
- James, K. J., Bishop, A. G. & Furey, A. 2000. New Toxins on the Horizon. In Botana, L. M. (ed.). *Seafood and freshwater toxins: Pharmacology, physiology, and detection*, pp. 693-714. New York: Marcel Dekker, Inc.
- Karafas, S., York, R. & Tomas, C. 2015. Morphological and genetic analysis of the *Coolia monotis* species complex with the introduction of two new species, *Coolia santacroce* sp. nov. and *Coolia palmyrensis* sp. nov. (Dinophyceae). *Harmful Algae*. 46, 18-33.
- Kibler, S. R., Holland, W. C., Vandersea, M. W., Faust, M. A., Litaker, R. W. & Tester, P. A. 2010. A collection method for Ciguatera-associated dinoflagellates. In Kin-Chung, H., Zhou, M. J. & Qi, Y. Z. (eds.). *Harmful Algae 2008. Proceedings of 13th International Conference on Harmful Algae*. November 3-7, 2008. Hong Kong.
- Kibler, S. R., Tester, P. A., Kunkel, K. E., Moore, S. K. & Litaker, R. W. 2015. Effects of ocean warming on growth and distribution of dinoflagellates associated with ciguatera fish poisoning in the Caribbean. *Ecological Modelling*. 316, 194–210.

- Kita, M., Ohishi, N., Washida, K., Kondo, M., Koyama, T., Yamada, K. & Uemura, D. 2005. Symbioimine and neosymbioimine, amphoteric iminium metabolites from the symbiotic marine dinoflagellate *Symbiodinium* sp. *Bioorganic Med. Chem.* 13, 5253-5258.
- Kohler, S. & C. Kohler. 1992. Dead bleached coral provides new surfaces for dinoflagellates implicated in ciguatera fish poisonings. *Environmental Biology of Fishes.* 35(4), 413-416.
- Landsberg, J. H. 2002. The effects of harmful algal blooms on aquatic organisms. *Rev Fisheries Sci.* 10(2), 113-390.
- Leaw, C. P., Lim, P. T., Ahmad, A. & Usup, G. 2001. Genetic diversity of *Ostreopsis ovata* (Dinophyceae) from Malaysia. *Mar. Biotechnol.* 3, 246-255.
- Leaw, C. P, Lim, P. T., Tan, T. H., Tuan-Halim, T. N., Cheng, K. W., Ng, B. K. & Usup, G. 2011. First report of the benthic dinoflagellate, *Gambierdiscus belizeanus* (Gonyaulacales: Dinophyceae) for the east coast of Sabah, Malaysian Borneo. *Phycological Research.* 59, 143-146.
- Lehane, L. & Lewis, R. J. 2000. Ciguatera: Recent advances but the risk remains. *International Journal of Food Microbiology.* 61(2-3), 91-125.
- Lewis, N. D. 1986. Epidemiology and impact of ciguatera in the Pacific: A review. *Mar. Fish. Rev.* 48(4), 6-13.
- Lewis, R. J., Vernoux, J. P. & Brereton, I. M. 1998. Structure of Caribbean ciguatoxin isolated from *Caranx latus*. *J. Am. Chem. Soc.* 120, 5914-5920.
- Lewis, N. I., Wolny, J. L., Achenbach, J. C., Ellis, L., Pitula, J. S., Rafuse, C., Rosales, D. S. & McCarron, P. 2018. Identification, growth and toxicity assessment of *Coolia Meunier* (Dinophyceae) from Nova Scotia, Canada. *Harmful Algae.* 75, 45-56.
- Lim, P. T., Leaw, C. P. & Usup, G. 2003. *Status of HAB and potential remote sensing application in detection of HAB events in Malaysia water.* Workshop on red tide monitoring in Asian Coastal Waters, Graduate School of Agricultural and Life Sciences, The University of Tokyo, pp. 81-84.
- Lim, H. C., Lim, P. T., Leaw, C. P. & Teng, S. T. 2014. An unprecedent bloom of *Gonyaulax polygramma* in Kota Kinabalu, Sabah, Malaysian Borneo. *Harmful Algae News.* 49, 9.
- Litaker, R. W., Vandersea, M. W., Faust, M. A., Kibler, S. R., Nau, A. W., Holland, W. C., Chinain, M., Holmes, M. J. & Tester, P. A. 2010. Global distribution of ciguatera causing dinoflagellates in the genus *Gambierdiscus*. *Toxicon.* 56, 711-730.
- Lobel, P. S., Anderson, D. M. & Durand-Clement, M. 1988. Assessment of ciguatera dinoflagellate populations: sample variability and algal substrate selection. *Biol. Bull.* 175, 94-101.

- Mabrouk, L., Hamza, A., Mahfoudi, M. & Bradai, M. N. 2012. Spatial and temporal variations of epiphytic *Ostreopsis siamensis* on *Posidonia oceanica* (L.) Delile leaves in Mahdia (Tunisia). *Cah. Biol. Mar.* 53, 419–427.
- Mangialajo, L., Bertolotto, R., Cattaneo-Vietti, R., Chiantore, M., Grillo, C., Lemee, R. & Ruggieri, N. 2008. The toxic benthic dinoflagellate *Ostreopsis ovata*: Quantification of proliferation along the coastline of Genoa, Italy. *Marine Pollution Bulletin*. 56(6), 1209–1214.
- Mangialajo, L., Ganzin, N., Accoroni, S., Asnaghi, V., Blanfuné, A., Cabrini, M., Cattaneo-Vietti, R., Chavanon, F., Chiantore, M., Cohu, S., Costa, E., Fornasaro, D., Grossel, H., Marco-Miralles, F., Masó, M., Reñé, A., Rossi, A.M., Montserrat Sala, M., Thibaut, T., Totti, C., Vila, M. & Lemée, R. 2011. Trends in *Ostreopsis* proliferation along the Northern Mediterranean coasts. *Toxicon*. 57, 408–420.
- Mattei, C., Vetter, I., Eisenblätter, A., Krock, B., Ebbecke, M., Desel, H. & Zimmermann, K. 2014. Ciguatera fish poisoning: A first epidemic in Germany highlights an increasing risk for European countries. *Toxicon*. 91, 76–83.
- Meunier, A. 1919. Microplankton de la mer Flamande. III. Les Péridiens. In Hayez, M. (ed.). *Mém. Mus. R. Hist. Nat. Belgique Bruxelles*. 8(1), 116.
- Mohammad-Noor, N., Daugbjerg, N., Moestrup, Ø. & Anton, A. 2007. Marine epibenthic dinoflagellates from Malaysia – a study of live cultures and preserved samples based on light and scanning electron microscopy. *Nord. J. Bot.* 24, 629–690.
- Mohammad Noor, N., Adam, A., Soler, J. M. F., Anton, A. & Muhamad Shaleh, S. R. 2008. First Record of Toxic *Gymnodinium catenatum* off the West Coast of Sabah, Malaysia. In Ho, K-C., Zhou, M. J., Qi, Y. Z. & Virginia Kai, Y-K. (eds.). *Proceeding of the 13th International Conference on Harmful Algae*, pp. 243–245. Hong Kong: International Society for the Study of Harmful Algae & Environmental Publication House.
- Mohammad-Noor, N., Moestrup, Ø., Lundholm, N., Fraga, S., Adam, A., Holmes, M. & Saleh, E. 2013. Autecology and phylogeny of *Coolia tropicalis* and *Coolia malayensis* (Dinophyceae), with emphasis on taxonomy of *C. tropicalis* based on light microscopy, scanning electron microscopy and LSU rDNA. *J. Phycol.* 49, 536–545.
- Mohammad Noor, M., Adam, A., Saad, S., Khodzori, F. A. & Shaleh, S. R. M. 2019. Substrate Preferences and Effects of Medium with Soil Extract on Growth of Genus *Coolia* (Dinophyceae). *J. Sustainability Science and Management*. 14 (2), 71–81.
- Mohd Shymer, A. H. & Marsudi, M. 2011. Kejadian Keracunan Makanan Ciguatera di Jalan Tenom, Keningau. 7th Kelantan Health Conference. June 15–16, 2011. *Malaysian Journal of Public Health Medicine*. 11 (2), pp. 34.
- Morton, S. L., Norris, D. R. & Bomber, J. W. 1992. Effect of temperature, salinity and light intensity on the growth and seasonality of toxic dinoflagellates associated with ciguatera. *J. Exp. Mar. Biol. Ecol.* 157, 79–90.

- Murakami, Y., Oshima, Y. & Yasumoto, T. 1982. Identification of okadaic acid as a toxic component of a marine dinoflagellate *Prorocentrum lima*. *Nippon Suisan Gakkaishi*. 48, 69–72.
- Nakahara, H., Sakami, T., Chinain, M. & Ishida, Y. 1996. The role of macroalgae in epiphytism of the toxic dinoflagellate *Gambierdiscus toxicus* (Dinophyceae). *Phycological Research*. 44(2), 113–117.
- Norris, D. R., Bomber, J. W. & Balech, E. 1985. Benthic dinoflagellates associated with ciguatera from Florida Keys. I. *Ostreopsis heptagona* sp. nov, pp. 39–44. In Anderson, D. M., White, A. W. & Baden, D. G. (eds.). *Toxic dinoflagellates*. Elsevier.
- Okolodkov, Y. B., Campos-Bautista, G., Gárate-Lizárraga, I., González-González, J. A. G., Hoppenrath, M. & Arenas, V. 2007. Seasonal changes of benthic and epiphytic dinoflagellates in the Veracruz reef zone, Gulf of Mexico. *Aquat. Microb. Ecol.* 47, 223–237.
- Okolodkov, Y. B., Merino-Virgilio, F. D. C., Aké-Castillo, J. A., Aguilar-Trujillo, A. C., Espinosa-Matías, S. & Herrera-Silveira, J. A. 2014. Seasonal changes in epiphytic dinoflagellate assemblages near the northern coast of the Yucatan Peninsula, Gulf of Mexico. *Acta Botanica Mexicana*. 107, 121–151.
- Parsons, M. L. & Preskitt, L. B. 2007. A survey of epiphytic dinoflagellates from the coastal waters of the island of Hawai'i. *Harmful Algae*. 6, 658–669
- Parsons, M. L., Settemier, C. J. & Bienfang, P. K. 2010. A simple model capable of simulating the population dynamics of *Gambierdiscus*, the benthic dinoflagellate responsible for ciguatera fish poisoning. *Harmful Algae*. 10(1), 71–80.
- Parsons, M. L., Settemier, C. J. & Ballauer, J. M. 2011. An examination of the epiphytic nature of *Gambierdiscus toxicus*, a dinoflagellate involved in ciguatera fish poisoning. *Harmf. Algae*. 10, 598–605.
- Parsons, D. L., Aligizaki, K., Bottein, M. D., Fraga, S., Morton, S. L., Penna, A. & Rhodes, L. 2012. *Gambierdiscus* and *Ostreopsis*: Reassessment of the state of knowledge of their taxonomy, geography, ecophysiology, and toxicology. *Harmful Algae*. 14, 107–129.
- Peirano, A., Cocito, S., Banfi, V., Cupido, R., Damasso, V., Farina, G., Lombardi, C., Mauro, R., Morri, C., Roncarolo, I., Saldaña, S., Savini, D., Sgorbini, S., Silvestri, C., Stoppelli, N., Torricelli, L. & Bianchi, C. N. 2011. Phenology of the Mediterranean seagrass *Posidonia oceanica* (L.) Delile: medium and long-term cycles and climate inferences. *Aquat. Bot.* 94, 77–92.
- Penna, A., Vila, M., Fraga, S., Giacobbe, M. G., Andreoni, F., Riobo, P. & Vernezi, C. 2005. Characterization of *Ostreopsis* and *Coolia* (Dinophyceae) isolates in the Western Mediterranean Sea based on morphology, toxicity and Internal Transcribed Spacer 5.8S rDNA sequences. *J. Phycol.* 41, 212–225.
- Popowski, G., Delgado, G., Sánchez, M. & Nodar, R. 2001. *G. toxicus* Adachi y Fukuyo, en el litoral norte de Ciudad de la Habana. *Rev. Invest. Mar.* 21, 69–72.



- Potisarn, N., Lirdwitayaprasit, T., Pholpunthin, P. & Iwataki, M. 2017. Distribution and abundance of benthic dinoflagellates in seagrass areas of the eastern coast of the upper Gulf of Thailand. *Proceedings of the 4th National Meeting on Biodiversity Management in Thailand*. June 21-23, 2017. Udon Thani Province, Thailand.
- Praseno, J. P. & Wiadnyana, N. N. 1996. HAB Organisms in Indonesia Water. In Pennet, R. W. (ed.). Canadian Technical Report of Fisheries and Aquatic Sciences, Canada, pp. 69-75.
- Radke, E. G., Reich, A. & Morris, J. G. 2015. Epidemiology of Ciguatera in Florida. *Am. J. Trop. Med. Hyg.* 93(2), 425-432.
- Rahman Shah, M. M., An, S. J. & Lee, J. B. 2013. Seasonal abundance of epiphytic dinoflagellates around coastal waters of Jeju Island, Korea. *Journal of Marine Science and Technology*. 21, 156-165.
- Rhodes, L. L. & Thomas, A. E. 1997. *Coolia monotis* (Dinophyceae): a toxic epiphytic microalgal species found in New Zealand. *NZ J Mar Fresh Res.* 31, 139–141.
- Rhodes, L., Adamson, J., Suzuki, T., Briggs, L. & Garthwaite, I. 2000. Toxic marine epiphytic dinoflagellates, *Ostreopsis siamensis* and *Coolia monotis* (Dinophyceae), in New Zealand. *NZ J. Mar. Freshwat. Res.* 34, 371–383.
- Richlen, M. L. & Lobel, P. S. 2011. Effects of depth, habitat, and water motion on the abundance and distribution of ciguatera dinoflagellates at Johnston Atoll, Pacific Ocean. *Mar. Ecol. Prog. Ser.* 421, 51–66.
- Ronquist, F. & Huelsenbeck, J. P. 2003. MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics*. 19(12), 1572-1574.
- Roy, R. N. 1977. Red tide and outbreak of paralytic shellfish poisoning in Sabah. *Medical Journal of Malaysia*. 31(3), 247-251.
- Schmidt, J. 1902. Flora of Koh Chang. Contribution to the knowledge of the vegetation in the Gulf of Siam. Part IV. Peridiniales. *J. Botanique*. 23, 212–218.
- Scholin, C. A., Herzog, M., Sogin, M. & Anderson, D. M. 1994. Identification of group- and strain-specific genetic marks for globally distribute *Alexandrium* (Dinophyceae). II. Sequence analysis of a fragment of the LSU rRNA gene. *J. Phycol.* 30, 999-1011.
- Sellner, K. G., Doucette, G. J. & Kirkpatrick, G. J. 2003. Harmful algal blooms: causes, impacts and detection. *J. Ind. Microbiol. Biotechnol.* 3, 383–406.
- Shears, N. T. & Ross, P. M. 2009. Blooms of benthic dinoflagellates of the genus *Ostreopsis*; an increasing and ecologically important phenomenon on temperate reefs in New Zealand and worldwide. *Harmful Algae*. 8(6), 916-926.
- Sidabutar, T., Praseno, D. P. & Fukuyo, Y. 2001. Harmful algae in Indonesian waters. In Hallegraeff, G., et al., (eds.). *Harmful Algal Blooms 2000*. Intergovernmental Oceanographic Commission of UNESCO, pp. 124-128.

- Smayda, T. J. 1997. What is a bloom? A commentary. *Limnol. Oceanogr.* 42, 1132-1136.
- Smith, R. L. 1980. *Ecology and field biology*. 3rd ed. New York: Harper & Row Publishers.
- Steidinger, K. A., 1983. A re-evaluation of toxic dinoflagellate biology and ecology. In Round, F. G. & Taylor, F. J. R. (eds.), *Progress in Phycological Research*, vol. 2. pp. 147-182.
- Steidinger, K. A. & Baden, D. G., 1985. Toxic marine dinoflagellates. In Spector, D.L. (ed.), *Dinoflagellates*, pp. 210-261. Orlando, FL, Academic Press.
- Steidinger, K. A. & Tangen, K. 1996. Dinoflagellates. In Tomas, C.R. (ed.), *Identifying Marine Diatoms and Dinoflagellates*, pp. 387-598. New York, Academic Press.
- Stonik, I. V. 1995. A potentially toxic dinoflagellate, *Prorocentrum minimum*, in AMurskii Bay of the Sea of Japan. *Russ. J. Mar. Biol.* 20, 314-320.
- Swofford, D. L., 2002. PAUP. Phylogenetic Analysis Using Parsimony (and other methods). Version 4.0b10. Sunderland, MA, USA: Sinauer Associates.
- Tan, T. H., Lim, P. T., Mujahid, A., Usup, G., & Leaw, C. P. 2013. Benthic harmful dinoflagellate assemblages in a fringing reef of Sampadi Island, Sarawak, Malaysia. *Marine Research in Indonesia*. 38(2), 77-87.
- Tango P. J., Magnien R., Butler W., Luckett C., Luckenbach M., Lacouture R. & Poukish C. 2005. Impacts and potential effects due to *Prorocentrum minimum* blooms in Chesapeake Bay. *Harmful Algae*. 4, 525- 531.
- Tartaglione, L., Mazzeo, A., Dell'Aversano, C., Forino, M., Giussani, V., Capellacci, S., Penna, A., Asnaghi, V., Faimali, M., Chiantore, M., Yasumoto, T. & Ciminiello, P. 2016. Chemical, molecular, and eco-toxicological investigation of *Ostreopsis* sp. from Cyprus Island: structural insights into four new ovatoxins by LC-HRMS/MS. *Analytical Bioanalytical Chemistry*. 408, 915-932.
- Tawong, W., Yoshimatsu, T., Yamaguchi, H. & Adachi, M. 2015. Effects of temperature, salinity and their interaction on growth of benthic dinoflagellates *Ostreopsis* spp. from Thailand. *Harmful Algae*. 44, 37-45.
- Taylor, F. J. R. 1979. The description of the benthic dinoflagellate associated with maitotoxin and ciguatoxin, including observations on Hawaiian material. In Taylor, D. L. & Seliger, H. H. (eds.). *Toxic dinoflagellate blooms*, pp. 71-77. Elsevier Scientific.
- Taylor, F. J. R. 1985. The distribution of the dinoflagellate *Gambierdiscus toxicus* in the eastern Caribbean. In Salvat, B. (ed.). *Proc. 5th Int. Coral Reef Congr. Antenne Museum*, pp. 423-428. Moorea: EPHE.
- Taylor, F. J. R. 1987. The biology of dinoflagellates. Oxford, Blackwell Scientific.

- Taylor, F. J. R., Hoppenrath, M., & Saldarriaga, J. F. 2008. Dinoflagellate diversity and distribution. *Biodiv. Cons.* 17, 407-418.
- Thompson, J. D., Gibson, T. J., Plewniak, F., Jeanmougin, F. & Higgins, D. G. 1997. The CLUSTAL_X Windows Interface: Flexible Strategies for Multiple Sequence Alignment Aided by Quality Analysis Tools. *Nucleic Acids Research.* 25(24), 4876–4882.
- Terao, K. 2000. Ciguatera toxins: toxicology. In Botana, L. (ed.). *Seafood and Freshwater Toxins: Pharmacology, Physiology and Detection*, pp. 449–472. New York: Marcel Dekker.
- Tester, P. A, Kibler, S. R., Holland, W. C., Usup, G., Vandersea, M. W., Leaw, C. P., Lim, P. T, Larsen, J., Mohammad-Noor, N., Faust, M. A & Litaker, R. W. 2014. Sampling Harmful Benthic Dinoflagellates: Comparison of Artificial and Natural Substrate Methods. *Harmful Algae.* 39, 8-25.
- Throndsen, J. 1995. Estimating cell numbers. In Hallegraeff, G. M., Anderson, D. M. & Cembella, A. D. (eds.). *Manual on Harmful Marine Microalgae, IOC Manuals and Guides No. 33*, pp. 63-80. Paris: UNESCO.
- Tindall, D. R. & Morton, S. L. 1998. Community dynamics and physiology of epiphytic/benthic dinoflagellates associated with ciguatera. In Anderson, D.M., Cembella, A.D. & Hallegraeff, G.M. (eds.). *Physiological Ecology of Harmful Algal Blooms: Proceedings of the NATO-ASI, volume 41*, pp. 293-313. Heidelberg: Springer Verlag.
- Tomas, C. R. & Baden, D. G. 1993. The influence of phosphorus source on the growth and cellular toxin content of the benthic dinoflagellate *Prorocentrum lima*. In Smayda, T.J. & Schimizu, Y. (eds.). *Toxic Phytoplankton Blooms in the Sea*, pp. 1–570. Amsterdam: Elsevier Science Publishers.
- Tosteson, T. R., Ballantine, D. L., Tosteson, C. G., Hensley, V. & Bardales, A. T. 1989. Associated bacterial flora, growth, and toxicity of cultured benthic dinoflagellates *Ostreopsis lenticularis* and *Gambierdiscus toxicus*. *Appl. Environ. Microbiol.* 55, 137-141.
- Totti, C., Accoroni, S., Cerino, F., Cucchiari, E. & Romagnoli, T. 2010. *Ostreopsis ovata* along the Conero Riviera (northern Adriatic Sea): relationships with environmental conditions and substrata. *Harmf. Algae.* 9, 233–239.
- Tubaro, A., Durando, P., Del Favero, G., Ansaldi, F., Icardi, G., Deeds, J. R. & Sosa, S. 2011. Case definitions for human poisonings postulated to palytoxins exposure. *Toxicon.* 57(3), 478-495.
- Turki, S. 2005. Distribution of toxic dinoflagellates along the leaves of seagrass *Posidonia oceanica* and *Cymodocea nodosa* from the Gulf of Tunis. *Cah. Biol. Mar.* 46, 29-34.
- Turki, S., Harzallah, A. & Sammari, C. 2006. Occurrence of harmful dinoflagellates in two different Tunisian ecosystems: the lake of Bizerte and the Gulf of Gabes. *Cah. Biol. Mar.* 47, 253-259.

- Turquet, J., Quod, J. P., Couté, A. & Faust, M. A. 1998. Assemblage of benthic dinoflagellates and monitoring of harmful species in Reunion Island, SW Indian Ocean, 1993-1996. In Reguera, B., Blanco, J., Fernández, M. L., Wyatt, E. (eds.). *Harmful Algae*, pp. 44-47. Xunta de Galicia, Spain and IOC: UNESCO.
- Van Dolah, F. M. 2000. Diversity of Marine and Freshwater Algal Toxins. In Botana, L. M. (ed.). *Seafood and freshwater toxins: Pharmacology, physiology, and detection*, pp 19-43. New York: Marcel Dekker, Inc.
- Vila, M., Garcés, E. & Masó, M. 2001. Potentially toxic epiphytic dinoflagellate assemblages on macroalgae in the NW Mediterranean. *Aquat. Microbiol. Ecol.* 26(1), 51–60.
- Wang, S. F., Tang, D. L., He, F. L., Fukuyo, Y. & Azanza, R. V. 2008. Occurrences of harmful algal blooms (HABs) associated with ocean environments in the South China Sea. *Hydrobiologia*. 596, 79-93.
- Xu, Y. X., Richlen, M. L., Morton, S. L., Mak, Y. M., Chan, L. L., Tekiau, A. & Anderson, D. M. 2014. Distribution, abundance and diversity of *Gambierdiscus* spp. from a ciguatera-endemic area in Marakei, Republic of Kiribati. *Harmful Algae*. 34, 56–68.
- Yasumoto, T., Nakajima, I., Bagnis, R. & Adachi, R. 1977. Finding of a dinoflagellate as a likely culprit of ciguatera. *Jpn. Soc. Sci. Fish.* 43, 1021–1026.
- Yasumoto, T., Inoue, A. & Bagnis, R. 1979. Ecological survey of a toxic dinoflagellate associated with ciguatera. In Taylor, D. L. & Seliger, H. (eds.). *Toxic Dinoflagellate Blooms*, pp. 221–224. New York: Elsevier.
- Yasumoto, T., Oshima, Y., Murakami, Y., Nakajima, I., Bagnis, R. & Fukuyo, Y. 1980a. Toxicity of benthic dinoflagellates. *Bull. Jpn. Soc. Fish.* 46, 327–331.
- Yasumoto, T., Inoue, A., Ochi, T., Fujimoto, K., Oshima, Y., Fukuyo, Y., Adachi, R. & Bagnis, R. A. 1980b. Environmental studies on a toxic dinoflagellate responsible for ciguatera. *Bull. Jpn. Soc. Sci. Fish.* 46, 1397–1404.
- Yasumoto, T., Murata, M., Oshima, Y., Matsumoto, G. K. & Clardy, J. 1984. Diarrhetic Shellfish Poisoning. *Seafood Toxins*. Vol. 262, pp. 2017-214. American Chemical Society
- Yasumoto, T., Seino, N., Murakami, Y. & Murata, M. 1987. Toxins produced by benthic dinoflagellates. *Biol. Bull.* 172, 128-131.
- Yasumoto, T., Underdahl, B., Aune, T., Hormazabal, V., Skulberg, O.M. & Oshima, Y. 1990. Screening for haemolytic and ichthyotoxic components of *Chrysochromulina polylepis* and *Gyrodinium aureolum* from Norwegian coastal waters. In Graneli, E., Sundstrom, B., Edler, L. & Anderson, D. M. (eds.). *Toxic marine phytoplankton*, pp. 436-440. New York: Elsevier Science Publishing Co.
- Yong, H. L., Mustapa, N. I., Lee, L. K., Lim, Z. F., Tan, T. H., Usup, G., Gu, H., Litaker, R. W., Tester, P. A., Lim, P. T. & Leaw, C. P. 2018. Habitat complexity affects

benthic harmful dinoflagellate assemblages in the fringing reef of Rawa Island, Malaysia. *Harmful Algae*. 78, 56-68.

Yoo, K. I., 1991. Population dynamics of dinoflagellate community in Masan bay with a note on the impact of environmental parameters. *Mar. Poll. Bull.* 23, 185-188.