

**POPULATION ECOLOGY OF IRRAWADDY  
DOLPHIN (*Orcaella brevirostris*) IN RAJANG  
AND SARIBAS RIVERS, SARAWAK,  
MALAYSIA**



**JAMES ANAK BALI**

**UMS**  
UNIVERSITI MALAYSIA SABAH

**BORNEO MARINE RESEARCH INSTITUTE  
UNIVERSITI MALAYSIA SABAH  
2016**

**POPULATION ECOLOGY OF IRRAWADDY  
DOLPHIN (*Orcaella brevirostris*) IN RAJANG  
AND SARIBAS RIVERS, SARAWAK,  
MALAYSIA**



**JAMES ANAK BALI**

**UMS**  
UNIVERSITI MALAYSIA SABAH

**THESIS SUBMITTED IN FULFILLMENT FOR  
THE DEGREE OF MASTER OF SCIENCE**

**BORNEO MARINE RESEARCH INSTITUTE  
UNIVERSITI MALAYSIA SABAH  
2016**

## DECLARATION

I hereby declare that the materials in this thesis is the result of my own work, except for the quotations, experts' opinions, equation, summaries and references, which have been duly acknowledged as referenced work.

Date: 13 May 2016

---

James Anak Bali  
PO20098024



UMS  
UNIVERSITI MALAYSIA SABAH

## CERTIFICATION

NAME : **JAMES ANAK BALI**  
MATRIC NO : **PO20098024**  
TITLE : **POPULATION ECOLOGY OF IRRAWADDY DOLPHIN  
(*Orcaella brevirostris*) IN RAJANG AND SARIBAS  
RIVERS, SARAWAK, MALAYSIA**  
DEGREE : **MASTER OF SCIENCE (MARINE SCIENCE)**  
VIVA VOCE DATE : **18 NOVEMBER 2015**

### DECLARED BY

**1. SUPERVISOR**

Assoc. Prof. Dr. Ejria Saleh



Signature

UMS  
UNIVERSITI MALAYSIA SABAH

**2. CO-SUPERVISOR**

Assoc. Prof. Dr. Saifullah Arifin Jaaman

\_\_\_\_\_

## ACKNOWLEDGEMENT

I would like to extend my appreciation and gratitude to a few people who has been with me in completing this academic project.

First and foremost, I would like to dedicate this thesis to my late parents (Bali Anak Tinggi and Sima anak Jimbun) who will always be my source of motivation and inspiration.

To my dearest wife; Nurin Anak Chunggat, my precious daughters; Eunice Sumang, Elly Sara, Emmalyn Selah and Edith Suelin - thank you for being the most supportive women in my life. Without your words of love and encouragement, I will always 'see the bottle as half empty'. My supervisor Assoc. Prof. Dr. Ejria Saleh from the Borneo Marine Research Institute, Universiti Malaysia Sabah (UMS) my co-supervisor Assoc. Prof. Dr. Saifullah Arifin Jaaman from the Faculty of the Institute of Oceanography and Environment, University Malaysia Terengganu (UMT); thank you for your expert technical advice and constructive comments. Your guidance really helped in the process of completing this project.

Funding for the data collection of this thesis was supported by the grant to Sarawak Forest Department and Sarawak Forestry Corporation from Ministry of Resources and Environment (NRE) under the 9<sup>th</sup> Malaysian Plan Development Project. My colleagues from the Sarawak Forestry Corporation; especially Oswald Braken Tisen, Toloy Keripin Munsang, Tonny Ganyai, Christopher Kri Ubang, Het Kaliang, Taha Wahab, Liboy, Rosli Ulis and Dennis Jengga; thank you for your patience and dedication in helping me in terms of data collection throughout the project.

To my best friend and colleague, Toloy Keripin Munsang; thank you for helping me to navigate the thesis proper to be in its present form. Your friendship is my fortitude. To my cousin, Leelajini Jothy Paranjothy from the Academy of Language Studies, UiTM Sarawak; thank you for your help in solidifying the writing process of my thesis.

I also would like to take this opportunity to express my sincere thanks to my former colleagues at the Borneo Marine Research Institute, UMS; Amyra Suryatie Kamaruzzan, Eda Izanie Ahmad Kamil, Hairul Masrini Binti Muhamad, Nur Muna Mohd Nor, Nur Azeyanti Norhashim, Khairulniza Rizwan and Teoh Shu Woan. Not forgetting my good friends; Ajahar Abdul Sahar, Josli@ Roslan, Bujang and Harun.

Last but not least, special thanks to all my family members and relatives in Roban and Saratok, Sarawak and Temburong, Brunei Darussalam. Thank you for giving me strength, against all odds.

James Anak Bali  
13 May 2016

## ABSTRACT

The Irrawaddy dolphin (*Orcaella brevirostris*) is the most common cetacean species inhabiting the coastal waters of the main estuaries of Sarawak. Their habitats are highly overlapping with the areas of intensive use by human activities, which have evolved potential threats to their very survival. The study was conducted in Rajang and Saribas rivers, Sarawak with the objectives to determine their spatial and temporal occurrence and distribution, density and abundance patterns, their association with environmental water parameters, and to identify their 'hot spot' area. Data collection was carried out from April 2009 to October 2010. Surveys were conducted using modified boat strip-transect technique, with the width of the river as the strip width for each transects and each river was divided into segments. Irrawaddy dolphins were found in all segments of both Rajang and Saribas rivers but they were not distributed evenly. Higher mean sighting frequency, mean number of individual sighted per survey, sighting rate, probability of sighting, density and abundance of Irrawaddy dolphins were recorded at the lower river segment of Rajang and Saribas rivers. Probability of sighting was highly significant between the Kuala Rajang-Sarikei and Sarikei-Sibu segments (McNemar:  $\chi^2 = 38.726$ ,  $df=1$ ,  $p=0.000$ ). Probability of sighting was highly significant between the Kuala Saribas and Pusa-Sebemban segments in Saribas River (McNemar:  $\chi^2 = 37.770$ ,  $df=1$ ,  $p=0.000$ ). The density (Mann-Whitney:  $U=28.0$ ,  $n=9$  and  $7$ ,  $p=0.758$ ) and abundance (Mann-Whitney:  $U=24.0$ ,  $n=9$  and  $7$ ,  $p=0.470$ ) of animals between the river segments in Rajang River are not statistically different. There is no statistically significant difference on density (Mann-Whitney:  $U=24.0$ ,  $n=12$  and  $8$ ,  $p=0.069$ ) and abundance (Mann-Whitney:  $U=23.0$ ,  $n=12$  and  $8$ ,  $p=0.57$ ) of animals between the river segments in Saribas River. The probability of sighting between the seasons in both Rajang and Saribas rivers was not statistical significant difference. The abundance of *O. brevirostris* was not statistically different between seasons for both in Rajang River (Kruskall-Wallis:  $H=2.768$ ,  $df=2$ ,  $p=0.251$ ) and Saribas River (Kruskall-Wallis:  $H=0.323$ ,  $df=2$ ,  $p=0.851$ ). Distribution of *O. brevirostris* in Rajang River is significantly associated with salinity with weak correlation, while in Saribas River their distribution is significantly associated with turbidity, surface water temperature and salinity with water temperature has strong negative correlation. The 'hot spot' of *O. brevirostris* in Rajang River was identified between Sungai Sebulal to Sungai Barong while in Saribas River, the area is situated between the waters of Kampung Sepinang-Kampung Beladin Hilir. The conservation and protection of *O. brevirostris* and their habitats in the two river systems need to be managed holistically with specific recommendations on conservation education to raise public awareness about the importance of protecting the new identified 'hot spot' areas, developed the the newly identified 'hot spots' as new eco-tourism tousim site for dolphin watching and continue carry out more detail researches to overcome the knowledge gaps that required to produce a 'Species Conservation Management Plan' for *O. brevirostris* in these areas.

## **ABSTRAK**

### **POPULASI EKOLOGI IKAN LUMBA-LUMBA (*Orcaella Brevirostris*) IRRAWADDY DI SUNGAI RAJANG DAN BATANG, SARAWAK**

*Lumba-lumba Irrawaddy (*Orcaella brevirostris*) merupakan spesies setasean yang paling kerap dijumpai di kawasan persisiran pantai dan kuala sungai utama di Sarawak. Habitatnya selalu bertindih dengan kawasan yang digunakan oleh aktiviti manusia dan berpotensi mendatangkan ancaman kepada kemandiriannya. Kajian telah dijalankan di Sungai Rajang dan Sungai Saribas, Sarawak dengan objektif untuk menentukan corak kehadiran, taburan, kepadatan dan kelimpahan mengikut kawasan dan musim, hubungan kaitnya dengan parameter air persekitaran dan mengenalpasti 'hot spot' atau 'kawasan tumpuan' *O. brevirostris*. Pengumpulan data telah dilakukan dari April 2009 hingga Oktober 2010. Tinjauan telah dilakukan dengan menggunakan teknik bot jalur-transek yang diubahsuai dengan menggunakan kelebaran sungai sebagai jaluran setiap transek. *O. brevirostris* ditemui di kesemua segmen pada kedua-dua sungai tetapi taburannya tidak sekata. Purata frekuensi penemuan, purata jumlah individu ditemui per tinjauan, kadar penemuan, kepadatan dan kelimpahan adalah lebih tinggi pada segmen yang terletak di hilir sungai. Kebarangkalian penemuan secara statistiknya sangat signifikan di antara segmen pada Sungai Rajang (McNemar:  $\chi^2 = 38.726$ ,  $df=1$ ,  $p=0.000$ ). Kebarangkalian penemuan di antara segmen di Sungai Saribas juga secara statistiknya sangat signifikan (McNemar:  $\chi^2 = 37.770$ ,  $df=1$ ,  $p=0.000$ ). Kepadatan (Mann-Whitney:  $U=28.0$ ,  $n=9$  and  $7$ ,  $p=0.758$ ) dan kelimpahan (Mann-Whitney:  $U=24.0$ ,  $n=9$  and  $7$ ,  $p=0.470$ ) *O. brevirostris* di antara segmen di Sungai Rajang adalah secara statistiknya tidak signifikan. Corak yang sama ditunjukkan pada kepadatan (Mann-Whitney:  $U=24.0$ ,  $n=12$  and  $8$ ,  $p=0.069$ ) dan kelimpahan (Mann-Whitney:  $U=23.0$ ,  $n=12$  and  $8$ ,  $p=0.57$ ) *O. brevirostris* di antara segmen di Sungai Saribas. Kebarangkalian penemuan di antara musim adalah secara statistiknya tidak signifikan di Sungai Rajang (Kruskall-Wallis:  $H=2.768$ ,  $df=2$ ,  $p=0.251$ ) dan Saribas (Kruskall-Wallis:  $H=0.323$ ,  $df=2$ ,  $p=0.851$ ). Taburan *O. brevirostris* di Sungai Rajang adalah secara statistiknya signifikan berhubung kait dengan saliniti dengan korelasi yang lemah, manakala di Sungai Saribas taburan mereka secara statistiknya signifikan berhubung kait dengan turbiditi, suhu permukaan air dan saliniti dengan korelasi negatif yang kuat. 'Kawasan tumpuan' *O. brevirostris* di Sungai Rajang telah dikenalpasti di antara Sungai Sebulang dan Sungai Barong manakala, di Sungai Saribas terletak di antara Kampung Sepinang ke Kampung Beladin Hilir. Pemuliharaan dan perlindungan *O. brevirostris* dan habitat mereka di kedua-dua sungai tersebut perlu diuruskan secara menyeluruh dengan saranan yang khusus terhadap: pendidikan konservasi untuk mempertingkatkan kesedaran awam tentang kepentingan menjaga 'kawasan tumpuan', membangunkan 'kawasan tumpuan' baru sebagai tapak eco-pelancongan untuk pemerhatian lumba-lumba dan meneruskan kajian terperinci untuk mengatasi jurang pengetahuan bagi menghasilkan Pelan Pengurusan Pemuliharaan *O. brevirostris* di kawasan-kawasan tersebut.*

# LIST OF CONTENTS

	Page
<b>TITLE</b>	i
<b>DECLARATION</b>	ii
<b>CERTIFICATION</b>	iii
<b>ACKNOWLEDGEMENT</b>	iv
<b>ABSTRACT</b>	v
<b><i>ABSTRAK</i></b>	vi
<b>LIST OF CONTENTS</b>	vii
<b>LIST OF TABLES</b>	x
<b>LIST OF FIGURES</b>	xi
<b>LIST OF ABBREVIATIONS</b>	xii
<b>LIST OF SYMBOLS</b>	xiv
<b>LIST OF APPENDICES</b>	xv
<b>CHAPTER 1: INTRODUCTION</b>	
1.1 Irrawaddy dolphin	1
1.2 Taxonomy and morphology	2
1.3 Research and conservation of marine mammals in Malaysia	3
1.4 Research and conservation of Irrawaddy dolphins in Sarawak	4
1.5 Significance and objectives of the study	6
<b>CHAPTER 2: LITERATURE REVIEW</b>	
2.1 Distribution	8
2.1.1 Distribution in South Asia and Southeast Asia	8
2.1.2 Distribution in Malaysia	10
2.2 Habitat Preference	11
2.3 Behaviour	11
2.3.1 Movement	11
2.3.2 Food and feeding	12
2.4 Population	13



2.5	Threats	15
2.6	Socio-economic status	17
2.7	Association with water parameters	17
2.7.1	Sea surface temperature (SST)	18
2.7.2	Surface water salinity	18
2.7.3	Turbidity	19
2.7.4	Depth	19
2.7.5	Other environmental factors	20
2.8	International, national and state agreements and legislations	21
2.8.1	International Agreement ratified by nation	21
2.8.2	National and state legislations	22

### **CHAPTER 3: METHODOLOGY**

3.1	Study area	24
3.2	Data collection	27
3.2.1	Occurrence and distribution	30
3.2.2	Water parameters	30
3.3	Data analysis	31
3.3.1	Occurrence and distribution	31
	a. Survey efforts	31
	b. Probability of sighting	32
3.3.2	Density and abundance	33
3.3.3	Association of Irrawaddy dolphins with water parameters	34
3.3.4	Determine of 'hot spot' of Irrawaddy dolphins	34

### **CHAPTER 4: RESULTS**

4.1	Survey effort	36
4.2	Occurrence and distribution	37
4.3	Probability of sightings	40
4.3.1	Spatial patterns	41
4.3.2	Temporal patterns	41
4.4	Density and abundance	42

4.4.1	Spatial patterns	42
4.4.2	Temporal patterns	43
4.5	Association of Irrawaddy dolphin with water parameters	44
4.5.1	Rajang River	44
4.5.2	Saribas River	45
4.6	Irrawaddy dolphins 'hot spot'	46
4.6.1	Rajang River	46
4.6.2	Saribas River	47
<b>CHAPTER 5: DISCUSSION</b>		
5.1	Survey effort and limitation	49
5.2	Spatial and temporal patterns	50
5.2.1	Spatial occurrence, distribution, density and abundance patterns	50
5.2.2	Temporal occurrence, distribution, density and abundance patterns	52
5.3	Association with environmental parameters	53
5.4	Irrawaddy dolphins 'hot spot'	54
<b>CHAPTER 6: CONCLUSIONS</b>		
6.1	Spatial and temporal occurrence, distribution, density and abundance patterns	56
6.2	Association with environmental parameters	56
6.3	Irrawaddy dolphins 'hot spot'	56
6.4	Recommendations for conservation management	56
6.4.1	Conservation education	57
6.4.2	Development of eco-tourism product based on <i>O. brevirostris</i>	57
6.4.3	Future research priorities	58
<b>REFERENCES</b>		59
<b>APPENDICES</b>		72

## LIST OF TABLES

	Page
Table 3.1: Category "class" of number of animals sighted in each sighting	32
Table 4.1: Number of survey replicates, total survey efforts, mean survey effort per survey replicates and mean sightings in each segment of Rajang and Saribas rivers	36
Table 4.2: Number of mean sightings frequency, number of individual sighted and sighting rates of Irrawaddy dolphin recorded in each segment for Rajang and Saribas rivers.	38
Table 4.3: Mean probability of sightings, $p$ (detection) with upper and lower values of confidence interval for each segment in Rajang and Saribas rivers	41
Table 4.4: The density and abundance of Irrawaddy dolphins for each segment with lower and upper values of confident interval in each segment of Rajang River	42
Table 4.5: The density and abundance of Irrawaddy dolphins for each segment with lower and upper values of confident interval in each segment of Saribas River	43
Table 4.6: Mean water parameters for every sightings with standard error (SE), confident interval (CI) and coefficient of variance (CV) in Rajang River	44
Table 4.7: Mean water parameters for every sightings with standard error (SE), confident interval (CI) and coefficient of variance (CV) in Saribas River	45
Table 4.8: The values of mean, minimum, maximum, median and standard error of water parameters at Irrawaddy dolphins 'hot spot' between Sungai Sebulal and Sungai Barong in Rajang River	47
Table 4.9: The values of mean, minimum, maximum, median and standard error of water parameters at Irrawaddy dolphins 'hot spot' between Kampung Sepinang and Beladin Hilir in Saribas River	47


## LIST OF FIGURES

	Page
Figure 1.1: Irrawaddy dolphins of Rajang River	1
Figure 2.1: Range of Irrawaddy dolphins, indicated by area shaded in red	9
Figure 3.1: Study site in the box square	25
Figure 3.2: Sampling segments in (a) Rajang River and (b) Saribas River	28
Figure 3.3: Diagram showing how strip boat-transect survey conducted	29
Figure 3.4: Measuring the physical water parameters using H10 HORIBA Water Quality Checker	31
Figure 3.5: Diagram (a) and (b) showing how 'hot spot' was identify based on Confidence Interfals of mean straight distance for all sighting class categories from the river mouth	35
Figure 4.1: Distribution of Irrawaddy dolphins in Rajang River	39
Figure 4.2: Distribution of Irrawaddy dolphins in Saribas River	40
Figure 4.3: Irrawaddy dolphin 'hot spot' in Rajang River	46
Figure 4.4: Irrawaddy dolphin 'hot spot' in Saribas River	48

## LIST OF ABBREVIATIONS

<b>AIC</b>	- Akaike Information Criterion
<b>BEACON</b>	- Biodiversity, Environment and Conservation
<b>CBD</b>	- Convention on Biological Diversity
<b>CeMMS</b>	- Centre for Marine Mammal Studies
<b>CI</b>	- Confidence Intervals
<b>CITES</b>	- Convention on International Trade in Endangered Species of Wild Fauna and Flora
<b>DoF</b>	- Department of Fisheries
<b>EEZ</b>	- Exclusive Economic Zone
<b>FDS</b>	- Forest Department Sarawak
<b>FGRS</b>	- Fundamental Research Grant Scheme
<b>GPS</b>	- Global positioning system
<b>HCVA</b>	- High conservation value area
<b>HiCOE</b>	- Higher Institution Centre of Excellence
<b>ICZM</b>	- Integrated coastal zone management
<b>IOES</b>	- Institute of Ocean and Earth Science
<b>IPPCS</b>	- Institut Penyelidikan Perikanan Cawangan Sarawak
<b>IUCN</b>	- International Union for Conservation of Nature
<b>MareCet</b>	- MareCet Research Organisation
<b>MLNG</b>	- Malaysian Liquefied Natural Gas
<b>MNMSRN</b>	- Malaysian National Marine Mammals Stranding Response Network
<b>MNS</b>	- Malaysian Nature Society
<b>MOHE</b>	- Ministry of Higher Education
<b>MOSTI</b>	- Ministry of Science, Technology and Innovation
<b>MPA</b>	- Marine Protected Area
<b>MPAs</b>	- Marine Protected Areas
<b>NGO</b>	- Non-Governmental Organization
<b>NPBD</b>	- National Policy on Biological Diversity
<b>NRE</b>	- Ministry of Natural Resource and Environment


<b>NTU</b>	- Nephelometric turbidity unit
<b>QGIS</b>	- Quantum Geographic Information System
<b>SCS</b>	- South China Sea
<b>SEAMAM</b>	- Southeast Asian Marine Mammal Symposium
<b>SFC</b>	- Sarawak Forestry Corporation
<b>SHELL</b>	- Sarawak Shell Berhad
<b>SST</b>	- Sea surface temperature
<b>SWD</b>	- Sabah Wildlife Department
<b>TPS</b>	- Totally protected species
<b>UAV</b>	- Unmanned aerial vehicle
<b>UMS</b>	- University Malaysia Sabah
<b>UMT</b>	- University Malaysia Terengganu
<b>WWF</b>	- World Wide Fund for Nature



**UMS**  
UNIVERSITI MALAYSIA SABAH

## LIST OF SYMBOLS

<b>%</b>	-	Percentage
<b>°C</b>	-	Degree Celcius
<b>M</b>	-	Meter
<b>Km</b>	-	Kilometer
<b>km<sup>-2</sup></b>	-	Per square Kilometer
<b>kmh-1</b>	-	Kilometer per hour
<b>Mt</b>	-	Metric tonnes
<b>NTU</b>	-	Nephelometric Turbidity Unit
<b>Nm</b>	-	Nautical mile
<b>Psu</b>	-	Practical Salinity Units
<b>mg/l</b>	-	Milligram per liter
<b>mS/cm</b>	-	Milisiemem per centimeter
<b>day<sup>-1</sup></b>	-	per day
<b>H</b>	-	hour
<b>h<sup>-1</sup></b>	-	per hour
<b>Min</b>	-	minute



UMS  
UNIVERSITI MALAYSIA SABAH

## LIST OF APPENDICES

	Page
Appendix A: Photo of Irrawaddy dolphins	72
Appendix B: Checklist of all confirmed marine mammal species found in East Malaysia, and Brunei	73
Appendix C: Analysis comparison of number of individuals recorded per sightings at different river segments	74
Appendix D: R Statistic Software Workspace analysis work command for probability of sightings according to river segments.	76
Appendix E: Analysis comparison the probability of sighting between river segments	78
Appendix F: Analysis comparison the probability of sighting in different season between river segments	80
Appendix G: Analysis comparison of abundance between river segments	86
Appendix H: Analysis comparison of density between river segments	89
Appendix I: Analysis comparison of seasonal abundance in Rajang River	91
Appendix J: Analysis comparison of seasonal abundance in Saribas River	93
Appendix K: Correlation of physical water parameters with distribution	96
Appendix L: Analysis R Sighting categorization analysis according to class in Saribas River	97
Appendix M: Analysis R sighting categorization analysis according to classification in Saribas River	99
Appendix N: 'Hot spot' range distance analysis	101
Appendix O: 'Hot spot' detection analysis	102
Appendix P: Data sheet for water parameters	103
Appendix Q: Data sheet for sighting	104



# CHAPTER 1

## INTRODUCTION

### 1.1 Irrawaddy dolphin

The earliest mentioned of the Irrawaddy dolphin (*Orcaella brevirostris*) in literature is by Owen (1866), who described a specimen found in 1852 at the mouth of the Vizagapatam (now known as Vishakhapatnam) river along the east coast of India (Stacey and Leatherwood, 1997). The species' name *brevirostris* derives from Latin, literally meaning short-beaked.

The Irrawaddy dolphin is one of the six subspecies of river dolphins which remain extant today (Beasley, 2007). Leatherwood and Reeves (1994) stated that the Irrawaddy dolphins are described as facultative river dolphins due to their ecological flexibility that allows them to inhabit both marine and freshwater environment (Figure 1.1).



**Figure 1.1: Irrawaddy dolphins of Rajang River.**

Irrawaddy dolphins are patchily distributed in shallow nearshore tropical and subtropical marine waters of the Indo-Pacific from the east coast of India in the Bay of Bengal, to Papua New Guinea, and from the northern part of Australia through the coast of Indo-China (Stacey and Arnold, 1999). However, recently the *Orcaella* were split into two species based on genetics, morphology and habitat preferences between Asian and Australia/Papua New Guinea specimens (Beasley et al., 2002a, 2005). The Australian/Papua New Guinea stocks of *Orcaella* were then designated as a separate species named Australian Snubfin dolphin (*O. heinsohni*) (Beasley et al., 2002b).

## **1.2 Taxonomy and morphology**

Once reaching adulthood, the Irrawaddy dolphin is about 2.0 - 2.5 m long. This dolphin has a rounded head that overhangs the mouth and a crescent-shaped blowhole. A neck crease is visible in some individual. Its dorsal fin is small, triangular, and slightly swept back with a blunt tip. Colouration is almost all grey but gets lighter towards the belly (Jefferson et al., 2008). A new born has an average mass between 10 - 12 kg. Adults can be from 2.1 - 2.3 m long and has an average mass of 114 - 133 kg (Arnold and Heinsohn, 1996). It is also equipped with a broad triangular dorsal view. The crescentic blowhole is displaced towards the left. The neck crease is usually distinct in some individual, about half way between the eye and anterior insertion of the flipper (Stacey and Arnold, 1999). The flippers are about 16% of body length, with a smooth curved anterior margin and maximum breadth less than half of the length along the anterior margin (Marsh et al., 1989; Stacey and Arnold, 1999). The dorsal fin, situated over halfway along the back, is rather small for a delphinid. Mean height ranged from 2.9% body length in Southeast Asian animals to 4.4% in Northeast Australian animals (Arnold and Heinsohn, 1996). The species obscure two to three-tone colour patterns from uniformly dark grey on the dorsal and lateral field, with variable shading among individuals, and is much lighter on the ventral field, which extends from the lower chin to the anus (Smith, 2008). The genital aperture is more anterior in males than females, and no other sexual dimorphism has been demonstrated (Stacey and Arnold, 1998). The skull has a short rostrum (43 - 45% of condylobasal length), with a well-developed mesethmoid crest. Its teeth have an expanded crown but are not compressed. The dental formula is

variable; published values include 13 to 17 (upper jaw)/12 to 14 mandible (Anderson, 1879), 16 to 20/15 to 19 (Arnold and Heinsohn, 1996 and 19/15 (Lloze, 1973).

### **1.3 Research and conservation of marine mammals in Malaysia**

The workshop on the Biology and Conservation of Small Cetacean and Dugongs of Southeast Asia (SEAMAM I) held from 27 - 30 June 1995 in Dumaguete, Philippines concluded that the prospects for a long-term persistence of many populations are generally poor and proposed surveys to determine the status and development of national programs in the conservation of inshore cetaceans and Dugongs in the region (Perrin et al., 1996). Malaysia was not represented due to the fact that, before 1996, there was no research establishment dedicated to study or conserve marine mammals in the country (Jaaman, 2006). In July 2002, a second similar workshop (SEAMAM II) was held in the same venue and highlighted the large and growing problems of cetaceans and Dugongs incidental catch in fisheries in Southeast Asia. Malaysia was represented by staff from the Forestry Department of Sarawak (FDS) and University Malaysia Sabah (UMS). The workshop concluded that unless the issues are addressed in an immediate, aggressive manner, major losses of biodiversity are inevitable and has proposed a Regional Action Plan to be developed and implemented in a phase-like manner, beginning with public awareness and an educational phases, (Perrin et al., 2005). The third Southeast Asian Marine Mammal Symposium (SEAMAM III) was organized in Fave Hotel, Langkawi, Malaysia from 4-10 March 2013. Malaysian delegates comprising of staff from Governmental agencies, universities and Non-governmental organizations (NGOs) conferred and discussed on several issues and miscellaneous matters, pertaining to domestic marine mammals.

UMS through the Centre for Marine Mammal Studies, CeMMS (formerly known as the Marine Mammals and Whale Shark Research and Conservation Programme) has initiated the research on marine mammals in Malaysia in 1996. From there, extensive and scientific studies on marine mammals were extended to Sarawak and Peninsular Malaysia. For the past 15 years, conservation and research efforts on marine mammals in Malaysia were focused on documentation of their occurrences and distributions, assessment on their population size, assessment and threats, and community perceptions.

Jamal Hisne et al., (2013) listed a number of initiatives on the marine mammals in Malaysia that was conducted by various management authorities, universities and non-governmental organizations (NGOs) through government grants or sponsorships by private sectors: Eight of postgraduate research projects in East Malaysia (Sabah and Sarawak) by CeMMS of UMS funded by the Ministry of Science, Technology and Innovation (MOSTI) e-Science Fund from 2008-2010); Evaluation on Management and Public Opinions for the Conservation of Marine Mammals in West Coast of Sabah by CeMMS from 2008 to 2010 sponsored by the Ministry of Higher Education(MOHE) under Fundamental Research Grant Scheme(FGRS); by Tioman Dolphin Project in 2010 by Malaysian Nature Society (MNS), and Institute of Ocean and Earth Sciences (IOES) of University Malaya (UM); Marine Mammals Surveys within the proposed Tun Mustapha Park area, Sabah by World Wide Fund for Nature-Malaysia (WWF Malaysia) in collaboration with the University of St. Andrews, Scotland in 2012; Numbers of research programs on Marine Mammals (Dugongs, Whale and Dolphins) by the Universiti Malaysia Terengganu, UMT under the Ministry of Higher Education (MOHE) Higher Institution Centre of Excellence (HiCoE) in Marine Science Project started from 2012; Langkawi Dolphin Research Program in Pulau Langkawi conducted by The MareCet Research Organization (MareCet) and IOES of UM in 2010; A collaboration project between MareCet and UM on an Ecology and Conservation of coastal cetaceans in The Matang Mangrove Forest Reserve and adjacent coastal waters, Perak in 2012; A study of dolphins of the Kinabatangan River in Sabah by the Land Empowerment Animals People and Sabah Forestry Department in collaboration with the University of St. Andrews, Scotland in 2010; and the establishment of the Malaysian National Marine Mammal Stranding Response Network (MNMMSN) involving a collaboration of multiple government agencies, non-governmental agencies, universities and private companies to provide effective response to stranding incidents around the country in 2013.

#### **1.4 Research and conservation of Irrawaddy dolphins in Sarawak**

Bali and Tisen (2008) reported that 15 of the species of marine mammals have been recorded in Sarawak, notably Bryde's whale (*Balaenoptera edeni*), Killer whale (*Orcinus orca*), Pygmy Sperm whale (*Kogia breviceps*), Sperm whale (*Physeter macrocephalus*), Risso's dolphin (*Grampus griseus*), Fraser's dolphin (*Lagenodelphis*

*hosei*), Indo-Pacific Humpback dolphin (*Sousa chinensis*), Irrawaddy dolphin (*Orcaella brevirostris*), Indo-Pacific Finless porpoise (*Neophocaena phocaenoides*), Dugong (*Dugong dugon*), Indo-Pacific Bottlenose dolphin (*Tursiops aduncus*), Common Bottlenose dolphin (*Tursiops truncatus*), Humpback whale (*Megaptera novaeangliae*), Long-snouted Spinner dolphin (*Stenella longirostris*) and Pan-Tropical Spotted dolphin (*Stenella attenuata*).

Although there are some publications on marine mammals from Sarawak were published in 1990s by Lydekker (1901); Weber, (1923); Bank, (1931); Gibson-Hill, (1949;1950); Fraser, (1956); Lewin, (1956); Harrison, (1957;1960); Harrison and Jamuh, (1958); Morzer-Bruyns, (1966); Beasley and Jefferson, (1997), extensive researches and conservation efforts only had begun with the establishment of collaboration studies between UMS and FDS in 2000. Since then, more information on marine mammals in Sarawak have been documented, compiled, reviewed and published in local and international journals; Jaaman et al., (2000; 2001; 2002), Jaaman, (2004; 2006); Jaaman, 2010; Lah-Anyi & Jaaman, 2002; W.Y. Chin, (2008); Bali, (2005); Bali et al., (2008; 2012); Bali and Tisen, (2008); Minton et al., (2019;2011;2013;2014). Past researches and conservation programmes conducted in Sarawak are: Interview surveys from fishermen, village headmen and/or knowledgeable villagers that were conducted along the coast of Sarawak from Lawas to Sematan in 2000 by UMS in collaboration with FDS; Aerial and in-shore boat surveys which were carried out along the coast of Sarawak in 2001 by UMS and FDS. Dialogue between SFC with tourism industries on marine mammals watching guideline in Sarawak was held in Kuching on March 2007; Aerial surveys on marine mammals along the coast of Sarawak conducted by the SFC in collaboration with UMS and Sabah Wildlife Department (SWD) from 19 June to 6 July 2007 (Bali, 2008); aerial surveys to monitor the Dugong population over Kuala Lawas was carried out from the 11-13 April 2008 (Bali and Tisen, 2008); offshore boat survey on marine mammals along the coast of Sarawak was conducted by the SFC in collaboration with UMS, *Institut Penyelidikan Perikanan Cawangan Sarawak* (IPPCS) and FDS from 18th June-2nd July 2008 (Bali, 2008); Number of collaboration studies between SFC and UMS on sea grasses and Dugongs distribution in Lawas, Sarawak from 2000 to 2008; Sarawak Dolphins' Project in 2008 to 2009 by the Sarawak Shell Berhad (Sarawak

SHELL), SFC and Universiti Malaysia Sarawak (UNIMAS); and Studies on the Occurrence and Distribution of Marine Mammals in the Malaysian South China Sea, and Species Composition, Distribution and Abundance of Sea grass and their Relationship with Dugong's population Ecology in Malaysian Bay of Brunei under MOHE HiCoE project by UMT in collaboration with SFC in 2013 to 2015; and Cetacean Survey and Monitoring in Bintulu waters under Biodiversity, Environment and Conservation (BEACON) project by Malaysian Liquefied Natural Gas (MLNG) group of companies and SFC from 2013 to 2018.

### **1.5 Significance and objectives of the study**

There are four species of inshore cetaceans, namely the Irrawaddy dolphin (*Orcaella brevirostris*), Indo-Pacific Bottlenose dolphin (*Tursiops aduncus*), Indo-Pacific Humpback dolphin (*Sousa chinensis*) and Finless porpoise (*Neophocaena phocaenoides*), inhabiting the coastal waters and main estuaries of East Malaysia. (Beasley and Jefferson, 1997; Jaaman et al., 2001; Jaaman et al., 2000; Jamaan et al., 2000b). Out of these four species, the Irrawaddy dolphin is the most common species and is considered as the flagship species in Sarawak.

Despite having laws to protect and conserve this species in Sarawak for more than decades, Irrawaddy dolphin's population size, ecology, abundance and threats to their population have been poorly studied and monitored due to lack of funding and experts. Applied population biology and ecology can contribute greatly to the management of this species, (Primack, 2002; Beasley, 2007).

Occasional sightings, stranding, and death of few of these animals have attracted the media and public attention; hence the awareness of their presence in the state. Based on published and unpublished papers, reports and personal observations, Irrawaddy dolphins are mostly abundant in the Saribas and Rajang rivers as compared to the other river tributaries in Sarawak. There are reasons for concerns about the population ecology of the Irrawaddy dolphins in these two river systems. In addition to their habitats that overlapped with areas of intensive use by human which has invoked potential threats to their survival, this species is also known to get trapped accidentally in fishing gears, threatened by declining fish resources,

loss or degradation of aquatic habitats, and direct hunting for Puffer fish baits, (Beasley and Jefferson 1997; Beasley 1998; Jaaman et al., 2001; Jaaman, 2000; 2001; Jamaan and Lah-Anyi, 2002; Pers. obse.). When managing threats and impacts, it is vital to have robust scientific understanding of a species' ecology, its distinctive characteristics, population status, and the dynamic processes that affect population size and distribution (Slooten and Lad, 1991; Primack, 2002). Primack (2002) stressed that conservation biologist need to answer question about species' environment, distribution, biotic interaction, systematic and morphology, physiology and life history, behavior, and genetic, to implement effective population-level conservation efforts. Thus, the main goal of this study is to provide scientific information as a foundation to plan conservation management strategies of Irrawaddy dolphins in Rajang and Saribas river systems.

To achieve the goal of this study, the following primary objectives have been set up:

- a. To determine the spatial and temporal occurrence, distribution, density and abundance patterns of Irrawaddy dolphin in Rajang and Batang rivers.
- b. To determine the association of Irrawaddy dolphins distribution in Rajang and Saribas rivers with environmental water parameters.
- c. To identify and map the 'hot spot' for Irrawaddy dolphins in Rajang and Saribas rivers

## CHAPTER 2

### LITERATURE REVIEW

#### 2.1 Distribution

Irrawaddy dolphins have been described as facultative river cetaceans, due to their ecological flexibility that allows them to inhabit the marine and freshwater environments (Leatherwood and Reeves 1994). They are apparently riverine, estuarine and coastal, often associated with mangrove forests (Dhandapani, 1992). There are some populations of Irrawaddy dolphin in Asia that are restricted to fresh water, and it is doubtful whether they would venture further offshore. Irrawaddy dolphins are found upstream in the Ayeryarwady (Irrawaddy) river system of Myanmar, the Mahakam river system in East Kalimantan, Indonesia, and the Mekong river system of Laos, Cambodia and Vietnam. The dolphins are present in the Ayeryarwady River to at least 1,287 km upriver from its mouth (Anderson, 1871; 1879; Thien, 1977). Tas'an and Leatherwood (1984) described that the species distributed in coastal waters near the river mouth of the Mahakam and upstream for at least 200km. However, Krebs (2002) revealed that the dolphins sporadically move as far as 80 km and 600 km down-and upstream, respectively. After the Australian/Papua New Guinea stocks of *Orcaella* were designated as a separate species named Australian Snubfin dolphin (*O. heinsohni*), Irrawaddy dolphins' distributions restricted to Indo-Pacific from the east coast of India in the Bay of Bengal (Beasley et al., 2002b). (Figure 2.1).

##### 2.1.1 Distribution in South Asia and Southeast Asia

In Bangladesh, Irrawaddy dolphins prefer coastal areas associated with muddy, brackish water at the river mouth, ranging offshore as far as the extent of freshwater plume-often only a few km but more than 60 km at the Meghna river mouth and waterways of the Sundarbans mangrove forest (IUCN, 2008).