

**CANDIDATE MUTANTS OF *Solanum*  
*Lycopersicum* DEVELOPED USING ETHYL  
METHANE SULFONATE**



**ASWANEEZA BINTI KHAIRUL ASWANDY**

**UMMS**  
UNIVERSITI MALAYSIA SABAH

**BIOTECHNOLOGY RESEARCH INSTITUTE**

**UNIVERSITI MALAYSIA SABAH**

**2019**

**CANDIDATE MUTANTS OF *Solanum*  
*Lycopersicum* DEVELOPED USING ETHYL  
METHANE SULFONATE**

**ASWANEEZA BINTI KHAIRUL ASWANDY**



**UMS**  
UNIVERSITI MALAYSIA SABAH

**THESIS SUBMITTED IN FULFILLMENT FOR  
THE DEGREE OF MASTER OF SCIENCE**

**BIOTECHNOLOGY RESEARCH INSTITUTE**

**UNIVERSITI MALAYSIA SABAH**

**2019**

**UNIVERSITI MALAYSIA SABAH**

**BORANG PENGESAHAN STATUS TESIS**

JUDUL: **CANDIDATE MUTANTS OF *Solanum Lycopersicum* DEVELOPED USING ETHYL METHANE SULFONATE**

IJAZAH: **SARJANA SAINS (BIOTEKNOLOGI)**

Saya **ASWaneeza Binti Khairul Aswandy**, Sesi **2014-2019**, mengaku membenarkan tesis Sarjana ini disimpan di Perpustakaan Universiti Malaysia Sabah dengan syarat-syarat kegunaan seperti berikut:-

1. Tesis ini adalah hak milik Universiti Malaysia Sabah
2. Perpustakaan Universiti Malaysia Sabah dibenarkan membuat salinan untuk tujuan pengajian sahaja.
3. Perpustakaan dibenarkan membuat salinan tesis ini sebagai bahan pertukaran antara institusi pengajian tinggi.
4. Sila tandakan ( / ):

SULIT

(Mengandungi maklumat yang berdarjah keselamatan atau kepentingan Malaysia seperti yang termaktub di dalam AKTA RAHSIA 1972)

TERHAD

(Mengandungi maklumat TERHAD yang telah ditentukan oleh organisasi/badan di mana penyelidikan dijalankan)

TIDAK TERHAD

Disahkan Oleh,

**ASWaneeza Binti Khairul  
Aswandy  
MZ1411006T**

(Tandatangan Pustakawan)

Tarikh : 11 Januari 2019

(Prof. Madya Dr. Kenneth Francis Rodrigues)  
Penyelia

## DECLARATION

The thesis was written by me and additional information has been properly cited in this thesis.

19 January 2018

---

Aswaneeza Binti Khairul Aswandy

MZ1411006T



UMS  
UNIVERSITI MALAYSIA SABAH

## CERTIFICATION

NAME : **ASWANEEZA BINTI KHAIRUL ASWANDY**

MATRIC NO : **MZ1411006T**

TITLE : **CANDIDATE MUTANTS OF *Solanum lycopersicum*  
DEVELOPED USING ETHYL METHANE SULFONATE**

DEGREE : **MASTER OF SCIENCE (BIOTECHNOLOGY)**

DATE OF VIVA VOCE : **MARCH 14, 2018**



**CERTIFIED BY**

**UMMS**

UNIVERSITI MALAYSIA SABAH

Signature

### **SUPERVISOR**

Assoc Prof. Dr. Kenneth Francis Rodrigues

---


## ACKNOWLEDGEMENT

I would like to express my deepest gratitude to my supervisor, Assoc Prof. Dr. Kenneth Francis Rodrigues for his effort in giving guidance and opinion throughout this research. I am grateful for his patient, passion and constant support for this research. I would also like to thank Assoc. Prof. Dr. Vijay Kumar, Prof. Dr. Clemente Michael Wong, Miss Cheong Bo Eng, Dr. Mailin Misson and Mr. Wilson Yong for encouraging me and giving guideline throughout my study in UMS.

I want to thank all the staff and students of Biotechnology Research Institute, UMS for helping me and encourage me during my study. I'm thankful for the lab assistant's madam Azian Ag. Besar, Christina Kungin, MonyMian, Salmi Afidah Shabuddin, Vidarita Maikin for assistant especially in the maintenance of greenhouse facility. Special thanks to all my friends Noor Haniza, Nurul Amzar, Raimi Redwan, Noraini Philip, Lia Natasha, Nur Marlessa, Elvina Clarie, Chan Yi Wei, Dexter Lee and Alex. Thank you to everyone that helped me throughout this study, I can't thank you enough. My best wishes to all of you.

Aswaneeza Binti Khairul Aswandy

19 January 2018



## ABSTRACT

*Solanum lycopersicum* belongs to the genus of *Solanum* with a small genome size of 950Mb of DNA in 91 scaffold aligned to 12 tomato chromosome. *Solanum lycopersicum* which has been known to be originated from Andean Region and widely used as a research material has contributed towards the vast amount of tomato genetic information such as the discovery of 34,727 protein coding gene and a mass of accumulated information of tomato functional genomics in various databases. However, the functional genomics of plant genes has not yet been fully correlated to its specific function which only 18,320 genes were specified. Despite of having a high economical value with other significant attribution and benefits, its fullness can't be achieved if vital gene function has not been known yet. The present study was performed to correlate the scrutinized targeted plant genes to its function in the basis of *S.lycopersicum* mutagenesis via chemical mutagen ethyl methane sulfonate (EMS). Mutant tomato plants were developed up to its second generation mutant line which its genomic DNA were sequenced with the use of nine primers for the study of functional genomics of the specific loci in the tomato genome. All 99 samples were sequenced and screened for the changes in genotypic aspect through TILLING and reference comparison to its control untreated DNA sequence via bioinformatic software. Based on the analyzed results a total of five genes were detected to be mutated through random mutation induced in seeds by ethyl methane sulfonate treatment. The variations in 12 main classes and 24 sub-classes of tomato phenotypes, Brix sugar index, chlorophyll level and tomato height that were observed and recorded from germination stage up to senescence stage of tomato plants were in line with the polymorphism screened in the sequence of the mutated genes. Sugar index were also relatively low in mutant fruits with a mean value of 3.8 %Brix in comparison to control tomato fruits of 6.2 %Brix, indicating HXK gene might take part in the Carbon catabolite repression (CCR). Mutagenic effect of EMS were notably in the variation of mutant and control plant heights, mutant plant were stunted with the shortest to be recorded with a height of 10 cm of its main stem. Chlorophyll levels were relatively lower in leaves of mutant plants. EXP gene was screened to be mutated and caused the growth of shorter stem and nodes of tomato plants as it regulates the loosening of plant cell wall. Ethyl Methanesulfonate has proved its reliability in plant mutagenesis through the screening of newly emerging technology of TILLING for the study of plant functional genomics and further intensive research can be applied on tomato whole genome. This research of tomato as a plant model could contribute in the agriculture sector that emphasized on *Solanum lycopersicum* and mutagenesis via Ethyl methanesulfonate.

## **ABSTRAK**

### **CALON-CALON MUTAN *Solanum lycopersicum* DIBANGUNKAN MENGGUNAKAN ETHYL METHANE SULFONATE**

*Solanum lycopersicum* tergolong dalam genus *Solanum* dengan saiz kromosom 950Mb asid deoksi ribonukleik (DNA) di 91 perancah dijajarkan kepada 12 kromosom tomato. *Solanum lycopersicum* telah diketahui berasal dari Wilayah Andean dan digunakan sebagai bahan penyelidikan secara meluas, ia menyumbang kepada sejumlah besar maklumat genetik tomato seperti penemuan 34,727 gen pengekodan protein dan sejumlah besar maklumat genomic fungsian tomato terkumpul dalam pangkalan data. Walaubagaimanapun, fungsi genomik gen tumbuhan belum dikaitkan sepenuhnya kepada fungsi khusus mereka, hanya 18,320 gen telah dispesifikasikan. Walaupun ia mempunyai nilai ekonomi yang tinggi dan faedah lain yang penting, kebaikan keseluruhannya tidak boleh dikecapi jika fungsi gen penting belum diketahui lagi. Kajian ini telah dijalankan untuk mengaitkan gen tumbuhan yang dikaji kepada fungsinya secara dasar mutagenesis *S. lycopersicum* menggunakan Ethyl methanesulfonate (EMS). Tanaman tomato mutan dibiak sehingga generasi kedua di mana DNA genomic mereka telah melalui proses penjujukan DNA dengan menggunakan sembilan primer bagi mengkaji genomic fungsian lokus tertentu dalam genom tomato. Kesemua 99 sampel telah melalui proses penjujukan DNA dan disaring untuk sebarang perubahan dalam aspek genotip melalui kaedah TILLING dan perbandingan kepada jujukan DNA kumpulan kawalan. Keputusan analisis menunjukkan sejumlah lima gen dikesan telah bermutasi melalui kaedah rawatan Ethyl methanesulfonate. Perbezaan pada 12 kelas utama dan 24 sub-kelas fenotip tomato, indeks gula Brix, paras klorofil dan ketinggian tomato yang direkodkan dari peringkat percambahan sehingga peringkat penuaan tanaman tomato adalah selaras dengan polimorfisme yang disaring di jujukan DNA gen yang bermutasi. Indeks gula buah-buahan mutan adalah rendah dengan nilai rata-rata 3.8 %Brix berbanding dengan buah tomato kumpulan kawalan iaitu sebanyak 6.2 % Brix, menunjukkan gen HXK berkemungkinan mengambil bahagian dalam Karbon Represi Katabolite. Kesan mutagenik EMS adalah ketara terutamanya pada variasi ketinggian tumbuhan mutan dengan kumpulan kawalan, pokok-pokok mutant terbantut dengan ketinggian paling rendah, 10 cm dan paras klorofil juga rendah. Saringan gen EXP menunjukkan ia bermutasi dan menyebabkan pertumbuhan batang dan nod tanaman tomato lebih pendek kerana ia mengawal proses pelonggaran dinding sel tumbuhan. Ethyl Methanesulfonate telah membuktikan kebolehpercayaannya dalam mutagenesis tumbuhan melalui kaedah saringan teknologi TILLING. Kajian genomic fungsian tumbuhan yang lebih intensif boleh dilakukan pada keseluruhan genom tomato. Penyelidikan ini yang menggunakan tomato sebagai model tumbuhan yang boleh menyumbang kepada sector pertanian menekankan penggunaan *Solanum lycopersicum* dan mutagenesis melalui Ethyl methanesulfonate.



## TABLE OF CONTENTS

	Page
<b>TITLE</b>	i
<b>DECLARATION</b>	ii
<b>CERTIFICATION</b>	iii
<b>ACKNOWLEDGEMENT</b>	iv
<b>ABSTRACT</b>	v
<b><i>ABSTRAK</i></b>	vi
<b>TABLE OF CONTENTS</b>	vii
<b>LIST OF FIGURES</b>	x
<b>LIST OF TABLES</b>	xii
<b>LIST OF ABBREVIATIONS</b>	xiii
<b>LIST OF APPENDICES</b>	xv
<b>CHAPTER 1: INTRODUCTION</b>	1
1.1 Research background	1
1.2 Objectives	4
1.3 Significant of study	4
<b>CHAPTER 2: LITERATURE REVIEW</b>	5
2.1 <i>Solanum Lycopersicum</i>	5
2.2 Breeding Program of Tomato	7
2.3 EMS as Chemical Mutagen	8
2.4 Targeting Local Lesion IN Genomes (TILLING)	11
2.5 Targeted Genes	12
2.6 Genotypic fingerprinting of Inter Simple Sequence Repeat (ISSR) for the preliminary mutation detection.	19

2.7	BRIX Index	19
2.8	Chlorophyll	20
2.9	Synopsis of Research Background	21
 <b>CHAPTER 3: MATERIALS AND METHODS</b>		 22
3.1	Brief Flow Chart of Experimental Design	22
3.2	Preparation of Seeds prior to EMS Treatment	23
3.3	Mutagenesis of tomato seeds	23
3.3.1	Response of EMS on First Generation Tomato Plants	24
3.4	Grow-out in Greenhouse	25
3.4.1	Irrigation and Fertigation	25
3.5	DNA Extraction	26
3.5.1	Measuring Quality and Quantity of Extracted DNA	26
3.6	Genomic fingerprinting of ISSR as preliminary mutation detection Step	27
3.6.1	Agarose Gel Electrophoresis of PCR products amplified by ISSR primers	29
3.6.2	Band scoring of ISSR	29
3.7	Growing M2 Population	30
3.8	Primers	30
3.9	Analysis of the mutagenic effect of EMS on phenotypes of mutant Tomato trees	32
3.9.1	Measurement of Total Solid Content in tomato fruits	32
3.9.2	Determination of chlorophyll content	33

3.10	Genotypic Analysis of mutant population	34
3.10.1	Condition of Polymerase Chain Reaction (PCR) for the Amplification of designated primers.	34
3.10.1	Agarose Gel Electrophoresis of PCR products amplified using Nine Primers	34
3.10.2	Purification of PCR products	35
3.10.3	Bioinformatic Analysis	35
3.10.4	Software applied in analysis of DNA sequence on specific genes	35
<b>CHAPTER 4: RESULTS AND DISCUSSION</b>		<b>38</b>
4.1	Study on dosage dependent effect of EMS for the development of mutant population.	38
4.2	Genomic DNA extraction of <i>Solanum lycopersicum</i>	42
4.3	Preliminary results for designed primers	43
4.4	Preliminary mutation detection via genomic fingerprinting of ISSR	46
4.5	List of phenotypes classes and subclasses	49
4.6	Brix index for quantifying total solid contain in fruits	53
4.7	Chlorophyll analysis	60
4.8	Analysis of mutagenic effect of EMS through specific primers	66
<b>CHAPTER 5: CONCLUSION</b>		<b>78</b>
5.1	Conclusion	78
<b>REFERENCES</b>		<b>80</b>
<b>APPENDIX</b>		<b>93</b>

## LIST OF FIGURES

	Page
Figure 2.1: Random Single nucleotide change in the genome of treated organisms	10
Figure 3.1: Flow-chart of Experimental Design	36
Figure 4.1: Control and mutant seedlings 7 days after germination on 0.5% v/v 24 hours EMS treatment.	40
Figure 4.2: Examples of mutant young seedlings, 12 days old from 0.5% EMS 24 hours treatment	41
Figure 4.3: Electrophoretic analysis of total genomic DNA.	42
Figure 4.4: Electrophoretic analysis of nine sets of genes on total genomic DNA.	43
Figure 4.5: Electrophoretic analysis of ISSR-1 on mutant genomic DNAsamples in reference to genomic DNA of control plant	48
Figure 4.6: Examples of changes in phenotype of mutant plants in reference to control plant	51
Figure 4.7: Visible phenotypic categories in M1 and M2 population	53
Figure 4.8: Brix values of M1 and M2 population in comparison to control	59
Figure 4.9: Average Brix value of M1 and M2 population in comparison to average Brix value of control	60
Figure 4.10: DNA sequence analysis of mutant with mutated gene NRP2	67
Figure 4.11: ORF sequence of Mutant M210 on NRP2 gene in reference to control ORF	68
Figure 4.12: DNA sequence analysis of mutant with mutated gene NHXK.	69
Figure 4.13: ORF sequence of Mutant M221 on NHXK gene in reference to control ORF	70

Figure 4.14:	DNA sequence analysis of mutant with mutated gene P2D1	71
Figure 4.15:	ORF sequence of Mutant M204 on P2D1 gene in reference to control ORF.	72
Figure 4.16:	DNA sequence analysis of mutant with mutated protein AGO6	73
Figure 4.17:	ORF sequence of Mutant M224 on AGO gene in reference to control ORF.	74
Figure 4.18:	DNA sequence analysis of mutant with mutated gene Expansin	75
Figure 4.19:	ORF sequence of Mutant M206 on Expansin gene in reference to control ORF	76



UMS  
UNIVERSITI MALAYSIA SABAH

## LIST OF TABLES

	Page
Table 2.1: DNA sequence of targeted genes	18
Table 3.1: Table of ISSR Primers	27
Table 3.2: Nine Design Primers for the study of gene function	30
Table 3.3: List of Bioinformatics Software and Statistical Analysis	36
Table 4.1: Percentage of germinated seedling for EMS treatments across four different duration of exposure	39
Table 4.2: Correlation Table of targeted genes	44
Table 4.3: List of ISSR Primers	46
Table 4.4: List of phenotypes observed on mutant and control plants	49
Table 4.5: Visible phenotypic categories and subclasses across M1 and M2 population	52
Table 4.6: Fruit size, fruit weight and Brix Index of control tomato fruits	54
Table 4.7: Fruit size, fruit weight and Brix Index of M1 tomato fruits	55
Table 4.8: Fruit size, fruit weight and Brix Index of control tomato fruits	56
Table 4.9: Fruit size, fruit weight and Brix Index of M2 tomato fruits	57
Table 4.10: Absorbance of light (mm) and content of chlorophyll A in mutant and control leaves	62
Table 4.11: List of mutated genes	66
Table 4.12: Mutation Density on mutated gene	66

## LIST OF ABBREVIATIONS

<b>AGO</b>	- Argonaute 6
<b>ATP</b>	- Adenosine triphosphate
<b>CTAB</b>	- Cetyltrimethylammonium bromide
<b>CIA</b>	- Chloroform-isoamyl
<b>CCR</b>	- carbon catabolite repression
<b>cDNA</b>	- Complementary DNA
<b>dsDNA</b>	- Double-stranded DNA
<b>DAF</b>	- Days after flowering
<b>DNA</b>	- Deoxyribonucleic acid
<b>EMS</b>	- Ethyl Methanesulfonate
<b>EXP</b>	- Expansin
<b>EtBr</b>	- Ethidium Bromide
<b>GMO</b>	- Genetically modified organism
<b>HEX</b>	- Hexokinase
<b>ISSR</b>	- Simple Sequence Repeat
<b>LD</b>	- Lethal Dose
<b>LTR</b>	- Long Terminal Repeat
<b>Mb</b>	- Megabase
<b>NCBI</b>	- National center for biotechnology information
<b>ORF</b>	- Open reading frame
<b>PHYSN</b>	- Phytoene Synthase 1
<b>PCANH</b>	- Carbonic Anhydrase
<b>P2D1</b>	- Photosystem II D1

<b>PHYSN</b>	- Phytoene Synthase 1
<b>PCANH</b>	- Carbonic Anhydrase
<b>PCR</b>	- Polymerase Chain Reaction
<b>RP2</b>	- RNA Polymerase 2
<b>RNA</b>	- Ribonucleic Acid
<b>SNPs</b>	- Single nucleotide polymorphism
<b>TILLING</b>	- Targeting Induced Local Lesions in Genome
<b>TSS</b>	- Total Soluble Solid
<b>UV</b>	- Ultra Violet



**UMS**  
UNIVERSITI MALAYSIA SABAH



## LIST OF APPENDICES

	Page
APPENDIX A: Photos of Solanum Lycopersicum grown in this study and greenhouse facility	95
APPENDIX B: Photos of fruits and leaves of mutant and control	96
APPENDIX C: ISSR Band Scoring	97
APPENDIX D: Absorbance of light (nm) and content of chlorophyll A in control and mutant leaves	98
APPENDIX E: Media and Solution	101
APPENDIX F: DNA Ladder	104



UMS  
UNIVERSITI MALAYSIA SABAH

# CHAPTER 1

## INTRODUCTION

### 1.1 Research Background

Tomato (*Solanum lycopersicum*) is a fruit-bearing vegetable plant that belongs to the family of Solanaceae which also commonly known as nightshades. Tomato and other related nightshades have been ascertained to be originated from Andean Region of South America including some areas of Ecuador, Peru and Chile (Sims, 1980). The Solanaceae family comprises more than 3000 species and were said to be the third most important plant for economical purposes. Solanaceae sub-family consists of a variety amount of crop species from potato to medicinal plants which are constantly used as plant model system in agricultural and biological field (Fernandez-pozo *et al.*, 2014). Plant breeding of *S. lycopersicum* is essential in ensuring better and improved characteristic of the plants to meet several necessities and requirements of commercial production. Tomato is chosen as a plant model system in this study as the genome sequence of tomato is relatively small with 950 megabases of DNA in 91 scaffold aligned to the 12 chromosomes found in tomato, this is relatively small compared to the other crops within Solanaceae family (Matsukura *et al.*, 2008), making tomato an ideal reference species for sequencing the genome of Solanaceae family (Emmanuel & Levy, 2002). SOL genomic network had managed to discover 34,727 protein coding gene (The tomato genome Consortium, 2012). Most of the plants from Solanaceae family exhibit genomic stability (Doganlar *et al.*, 2002). However, the gene function of tomato has not been fully studied with only 18,320 known gene correlated to its specific function. The development of mutant tomatoes by means of mutagenesis could facilitate the study of plant gene function (Reddaiah *et al.*, 2014) through the approach of newly emerging technology of TILLING (Gilchrist &

Haughn, 2005; Stemple, 2004) for the search of mutation on the specific studied genes.

Tomato is seen to contain high amount of vitamin which is mainly ascorbic acid, vitamin A, potassium, micro and macro nutrients. The high demand of tomato due to it's commercialize value and contribution to medical health benefits such as the presence of relatively high content of antioxidant particularly lycopene has necessitated the need to develop novel, high-yielding varieties of tomato plants (George *et al.*, 2004) and for the purpose of functional genomics (Emmanuel & Levy, 2002). Other than the relatively small genome of tomato and its thousands of discovered genes, *S.lycopersicon* is also specifically chosen for its fairly short life cycle which makes it a convenient plant to study up to their second generation. Tomato tree does not consume a lot of space and suitable to be grown with a constant parameter in greenhouse or transgenic facility. Novel varieties of tomato crop can be breed through four fundamental approaches namely conventional plant breeding, protoplast fusion, genetic engineering and mutation breeding.

Breeding programs of tomatoes such as conventional plant breeding, genetic engineering and protoplast fusion are the approaches to breed better or improved varieties of plants. These various approaches had led to tomato crops with commercially targeted traits such as texture and taste enhancement, enhanced growth, resistance to certain diseases and other useful quality traits. Despite of these approaches in the development of new varieties of crops plants, there are some issues that were consider as drawback specifically the negative prospect in regards to ethical issue.

Regulatory issues pertaining to genetically modified organism has been a main reason for the addition of mutagenesis as an alternative approach in functional genomics of *S. lycopersicum*. The approach of developing new verities of crops in a more rapid and easy way by means of mutation breeding was firstly discovered in 1930s (Schouten & Jacobsen, 2007). Techniques of mutagenesis breeding were then improved after World War II. The research approach adopted in this study focuses on chemical mutagenesis breeding via exposure of Ethyl Methanesulfonate (EMS) which is an alternative approach to genetic engineering.

Chemical mutagen such as Ethyl Methanesulfonate (EMS) which acts on guanine, is capable to produce mutation in diploid DNA with the range of two to 10 mutation per megabase (Mb) (Till *et al.*, 2007). Although mutagenesis breeding is more randomized compared to genetic modification of plants (Lai *et al.*, 2004), mutagenesis is a relatively conventional approach as it needs a fewer amount of plantlets, time and cost reduction in yielding high number of tomato plants (Kim *et al.*, 2006; Clark, 1963; Li *et al.*, 2016). The development of mutant population with significant change in phenotypes and physiological responses is a robust tool in the study of biological function of specific genes in plants. The use of chemical mutagen, EMS aided with new technology of Targeting Induced Local Lesions in Genome (TILLING) are the approaches used for the study of functional genomics in tomato plants. TILLING is the best approach used for the detection of mutation developed through EMS in terms of the reduced amount of reagent consumed, duration in producing a mutant population, high density of mutation to the work load in screening for target genes (Serrat *et al.*, 2014) which firstly tested on *Arabidopsis thaliana* (McCallum *et al.*, 2000) and mutation on *Drosophila melanogaster* (Bentley *et al.*, 2000). The functional genomics of the specific genes will be analyzed through the development of mutant plants via EMS and TILLING approach (Kim *et al.*, 2006) through the analysis in both of the plant's genomic and phenotypic aspect. The basis of TILLING involves the development of mutant population and ends with the screening and mutation discovery on the specific studied gene (Comai & Henikoff, 2006). TILLING has been an established technology in mutagenesis breeding for genetic studies linking genes with their associated functions.

## 1.2 Objectives

The research objectives of this study are:

- i. To develop a mutant population of *S.lycopersicum* via chemical mutagenesis.
- ii. To screen the population for mutant on the basis of phenotype.
- iii. To evaluate the effect of chemical mutagenesis on specific genes.

## 1.3 Significance of study

The discovery of thousands coding protein in tomato plant genome is one of the greatest achievements in plant genetics. Various research papers were published in regards to the number and types of new genes found in tomato, recently more than eighteen thousand genes were known publicly. Although this had lent a tremendous improvement in plant study, there are still a few genes have not yet been link to their specific function. This research approaches focus on the study of functional genomics in tomato as a model plant via TILLING technology. Mutation occurs in the targeted gene will aid in determining the function of studied gene. The study of gene function could bridge the gap between the function of the gene or gene regulation within the plant genome. The function of mutagenesis is proven for the study of reverse genetics through the screening of mutation along the target genes by means of DNA sequencing. The knowledge through this research could be useful for the study of this model plant and the mutagenic effect of chemical mutagen particularly EMS. This will benefit the agriculture sector that emphasize on the study of this model plant and gene regulation.

## CHAPTER 2

### LITERATURE REVIEW

#### 2.1 *Solanum lycopersicum*

Tomato (*Solanum lycopersicum*) originated in South America and the commercialization of this particular plant continues to dominate other countries due to its usefulness as a vegetable crop. The anatomic features of tomato consist of its unique compound leaves, differential texture of its fruit and sympodial shoot branching. Furthermore, the tremendous advantages of tomato also include its short life cycle, self-pollination and non-complex diploid genetics.

The word Tomato was originated from a Spanish word tomate where its species originally grown in the South American Andes. The large amount of tomato consumed made it classified as the second highly intake vegetable (Szczechura *et al.*, 2011). Tomato is one of the crops that has been harvested for the purpose of commercialization and contribute as a model plant for increasing the depth in knowledge for the aspect of growth, maturation and metabolism in the biology of fruits (Giovannoni, 2004; Carrari *et al.*, 2006) and considered as a good plant model system for Solanaceous family due to its considerably small genome size (Reddaiah *et al.*, 2014). The whole tomato genome has been successfully sequenced by The International Solanaceae Genomics Project (SOL) (Muller, 2005a; Muller 2005b). It's genome consisted of more than 75% of heterochromatin (Fernandez-pozo *et al.*, 2014) and 730 Mb of the tomato genome are pericentromeric heterochromatin with 220 Mb of the tomato chromosome made up of distal euchromatic segments (Szczechura *et al.*, 2011). Tomato genome was initially thought to be a huge challenge for full sequencing as its

genome is complicated and larger in size compared to animal model, it was then discovered that tomato genome in the range of 950MB is considered manageable.

The Tomato Genome Sequencing Project has been one of the most important projects in the history of sequencing, this project employed both Sanger's and next-generation sequencing (NGS) technologies for the assembly of plant genome as well as a draft of *Solanum pimpinellifolium* which was released in 2012 (TGC, 2012). The sequence data generated by 150 tomato Genome Consortium has further uncover the natural alleles exist in different genotypes of tomato (Mueller *et al.*, 2009). However, the functional potential of the major fraction of this newly generated resource is still undefined.

Tomato has been used for the study of functional genomics in plants. *Arabidopsis thaliana* was initially the main model plant for this purpose due to its significant characteristics advantages. With the identified similar leverage of small genome size and short life cycle of tomato in comparison to *Arabidopsis thaliana*, makes tomato an ideal plant for gene study (Watanabe *et al.*, 2007).

Research on tomato were previously conducted with various tomato varieties including Arka Vikas for the purpose of investigating the effect of mutagenic treatment on agronomic parameters such as seedling heights and plant height (Laskar *et al.*, 2016), Patharkutchi was used to study the frequency and spectrum of macro-mutations of EMS and its combined treatment (Dutta *et al.*, 2017), Red Setter for the reverse genetic study in creating new traits of specified variety (Minoia *et al.*, 2010) and *Lycopersicon esculentum* Mill. was chosen as the model plant for the study of effects of mutagen concentration (Aliyu & Adamu, 2007). Studies of these tomato varieties focuses on the use of ethyl methanesulfonate as the chemical mutagen however, tomato variety of Yates F1 Hybrid (Grosse Lisse) from Australia has never been studied before. Thus, Yates F1 Hybrid (Grosse Lisse) was chosen in this research to study the mutants developed from this specified variety via EMS mutagenic treatment.

## 2.2 Breeding Program of Tomato

Conventional plant breeding is one of the approaches to breed better or improved varieties of plants by means of selecting desirable traits and cross breeding between varieties. Newly improved next generation population could be made possible through this method which carries the best traits from the parental plants. Commercially targeted traits were texture and taste enhancement, resistance to certain diseases and additional nutrient values. Conventional plant breeding method starts with specifically selected traits possessed crops via pure-line selection and hybridization between crops for the breeding of future generation. Conventional plant breeding relies on sexual recombination (Manshardt, 2004). Although this step has greatly contributed in the development of new variety of crop plants with inherited gene of interest but the good traits of the parental are only passed down through multiple back-crossing between progeny and recurrent parent which is considered as a time consuming plant development method (Hoisington *et al.*, 1999)

Protoplast fusion in plants on the other hand uses the application of fusing two or more protoplast to transfer useful genes. This approach of genetic recombination is able to transfer diseases resistance, enhanced growth, heat and cold resistance and other useful quality traits for forming good varieties of plants. The process of making new plant hybrids through protoplast fusion requires the removal of cell wall and cell fusion via electrofusion as the prime step (Grosser & Gmitter, 2011). The successful fused cells will have to be grown into calluses, plantlets and finally develop into a fully grown crop (Salamiah *et al.*, 2000). This important fundamental approach uses genetic modification method and requires a few successful steps to efficiently transfer genes between protoplast cells, making it a higher chance of failure in transferring the targeted gene, moreover longer period is required for the regeneration of whole plant (Marccone *et al.*, 2010; Guan *et al.*, 2010)

Genetic engineering applies the concept of preserving the genetic material of parent by only inserting minute genetic information for expressing the chosen characteristics (Richard, 2004). Restriction enzyme aids in the separation of purified DNA into fragments in order to isolate the candidate gene (Nicholl &