


**EXPRESSION OF SIX CHLOROPLAST DNA
GENES IN *Jatropha curcas* CALLUS UNDER
LIGHT AND DARK CONDITIONS**

CASSANDRA CHEW SZE YII

UNIVERSITI MALAYSIA SABAH
**DISSERTATION SUBMITTED IN
FULFILLMENT FOR THE DEGREE OF MASTER
OF MOLECULAR BIOLOGY**

**BIOTECHNOLOGY RESEARCH INSTITUTE
UNIVERSITI MALAYSIA SABAH**

2011

UNIVERSITI MALAYSIA SABAH
BORANG PENGESAHAN STATUS THESIS@

JUDUL: *Expression of six chloroplast DNA genes in Jatropha curcas callus under light and dark conditions*

IJAZAH: *Master of Molecular Biology*

SAYA *Cassandra Chew Sze Fii*

SESI PENGAJIAN: *2009-2011*

mengaku membenarkan tesis (~~LPSM/ Sarjana/ Doktor Falsafah~~) ini disimpan di Perpustakaan Universiti Malaysia Sabah dengan syarat-syarat kegunaan seperti berikut:-

1. Tesis adalah hakmilik Universiti Malaysia Sabah.
2. Perpustakaan Universiti Malaysia Sabah dibenarkan membuat salinan untuk tujuan pengajian sahaja.
3. Perpustakaan dibenarkan membuat salinan tesis ini sebagai bahan pertukaran antara institusi pengajian tinggi.

4. Sila tandakan (/)

SULIT

TERHAD

TIDAK TERHAD

(Mengandungi maklumat yang berdarjah keselamatan atau Kepentingan Malaysia seperti yang termaktub di dalam AKTA RAHSIA RASMI 1972)

(Mengandungi maklumat TERHAD yang telah ditentukan oleh organisasi/badan di mana penyelidikan dijalankan)



(TANDATANGAN PENULIS)

Alamat Tetap: _____

Tarikh: 26/08/11

UNIVERSITI MALAYSIA SABAH
PENYARAH
UNIVERSITI MALAYSIA SABAH



(TANDATANGAN PUSTAKAWAN)

Penyarah
Institut Penyelidikan Bioteknologi
Universiti Malaysia Sabah

Nama Penyelia

Tarikh: 13/09/2011

CATATAN:- *Potong yang tidak berkenaan.

** Jika tesis in SULIT atau TERHAD, sila lampirkan surat daripada pihak berkuasa/organisasi berkenaan dengan menyatakan sekali sebab dan tempoh tesis ini perlu dikelaskan sebagai SULIT dan TERHAD.

@ Tesis dimaksudkan sebagai tesis bagi Ijazah Doktor Falsafah dan Sarjana secara penyelidikan atau disertai bagi pengajian secara kerja kursus dan Laporan Projek Sarjana Muda (LPSM).

DECLARATION

I hereby declare that the material on this thesis is my own except for quotations, excerpts, equations, summaries and references, which have been duly acknowledged.

25th Aug 2011



Cassandra Chew Sze Yii
PB2009-8039



UMS
UNIVERSITI MALAYSIA SABAH

CERTIFICATION

NAME : **CASSANDRA CHEW SZE YII**

MATRIC NO. : **PB2009-8039**

TITLE : **EXPRESSION OF SIX CHLOROPLAST DNA GENES IN
Jatropha curcas CALLUS UNDER LIGHT AND DARK
CONDITIONS**

DEGREE : **MASTER OF SCIENCE (MOLECULAR BIOLOGY)**

VIVA DATE : **04 AUGUST 2011**



UMS
UNIVERSITI MALAYSIA SABAH

DECLARED BY

1. SUPERVISOR

Mr. Wilson Yong Thau Lym

2. CO-SUPERVISOR


Dr. Kenneth Rodrigues

ACKNOWLEDGEMENT

I wish to express my deepest gratitude and appreciation to my supervisor, Mr Wilson Yong who has been patience enough to guide and advice me during this study. His assistance truly assisted me to solve all my problems and doubts during the study then let me able to complete my thesis on time. My special sense of gratitude to my co-supervisor, Dr Kenneth for being helpful and guided me in all my works. His continuous encouragement provided me the necessary impetus to complete the research on time. Most of all, I would like to thank Prof. Datin Dr. Ann Anton, Director of Biotechnology Research Institute, for her huge support.

I am sincerely feeling thankful and greatly appreciative for all the help, encouragement, guidance and friendship that I have obtained during working with this project from the staff of Biotechnology Research Institute (BRI). I hereby knowledge, with gratitude, Kak Vida, Kak Azian, Mony, and Richard and all the master students from BRI for their help on my research.

With love, I really give thousand thanks to my mummy and daddy for enduring, with patience and understanding and the financial support in order for me to successfully complete my project. Also not forgetting my lovely friends (Joveen Neoh, Kalnisha Devi, Jesica Seng, Daniel Yeoh and last but not least Joseph Koh) for their mentally support, patience and assistance.




UMS
UNIVERSITI MALAYSIA SABAH

ABSTRACT

EXPRESSION OF SIX CHLOROPLAST DNA GENES IN *Jatropha curcas* CALLUS UNDER LIGHT AND DARK CONDITIONS

The expression of genes encoded in the open reading frames of chloroplast genomes have been posited to be induced by light. The current study focused on the induction of ORFs encoded in the chloroplast genome of *Jatropha curcas* (Accession number FJ695500). *J. curcas* is an important non-edible oil seed crop, which produces oil with high calorific value and is regarded as a potential fuel substitute. By understanding the gene expression of chloroplast DNA under light stress, potential genes candidate can be used for callus transformation. Gene induction was characterized in leaves of the plant, green callus and white callus cultivated under condition of light and darkness. A total of six ORFs representing the genes *YCF 1*, *YCF 2*, *psbD* (photosystem II), *rbcL* (Rubisco), *matK* (Maturase K) and *rpoC1* (RNA polymerase) were targeted by designing specific primers for their characterization. Specificity of primers was tested against the genomic DNA. Transcripts of six targeted genes were detected in all three replicates of the green and white callus under light and dark, except for *ycf 2* gene in green callus under light. A part of gene *ycf 2* was no transcribed using reverse transcription PCR, was then validated using real-time PCR assay. The reverse transcription PCR did not detect the gene amplified by YCFD primer in the green callus under photoperiod treatment. The *ycf 2* gene located in the region of region 94050 to 95483 was post-transcriptional modified. In the real-time PCR assay, the amplification curves were produced from the amplification of all callus under different treatments, except for green callus under light.



UNIVERSITI MALAYSIA SABAH

ABSTRAK

*Ekspresi gen dikodkan dalam 'opening reading frames' (ORFs) dari genom kloroplas telah posited diinduksi oleh cahaya. Penyelidikan ini memfokuskan pada induksi ORFs dalam genom kloroplas *Jatropha curcas* (Aksesori nombor FJ695500). *J. curcas* adalah non-minyak biji tanaman yang penting dan menghasilkan minyak dengan kalori tinggi dan juga dianggap sebagai pengganti bahan bakar yang berpotensi. Dengan memahami gen DNA kloroplas di bawah tekanan ringan, potensi gen calon boleh digunakan untuk transformasi kalus. Gene induksi dipercirikan dalam daun tanaman, kalus putih dan kalus hijau ditanamkan dalam keadaan cahaya dan kegelapan. Sebanyak enam ORFs mewakili gen YCF 1, YCF 2, psbD (fotosistem II), rbcL (Rubisco), matK (Maturase K) and rpoC1 (RNA polymerase) menjadi sasaran untuk merancang primer khusus untuk ciri-ciri gen tersebut. Spesifisitas primer diuji terhadap DNA genom. Transkrip enam gen yang disasarkan telah dikesan di semua kalus hijau dan putih di bawah terang dan gelap, kecuali bagi kalus hijau di bawah cahaya. Sebahagian dari gen YCF 2 yang diuji dengan PCR konvensional telah 'switch off', kemudian disahkan menggunakan PCR 'real-time'. PCR konvensional tidak mengesahkan gen diamplifikasi dengan primer YCFD dalam kalus hijau bawah fotoperiodik. Transkripsi berbalik PCR tidak mengesan gen yang dikuatkan oleh primer YCFD dalam kalus hijau di bawah rawatan photoperiod. Ycf 2 gen yang terletak di rantau rantau 94050 untuk 95483 pasca-transkripsi diubahsuai. Dalam uji PCR real-time, kurva amplifikasi dihasilkan dari amplifikasi di semua kalus bawah perlakuan yang berbeza, kecuali untuk kalus hijau di bawah fotoperiode.*



UMS
UNIVERSITI MALAYSIA SABAH

TABLE OF CONTENTS

TITLE	Page
DECLARATION	i
CERTIFICATION	ii
ACKNOWLEDGEMENT	iii
ABSTRACT	iv
ABSTRAK	v
TABLE OF CONTENTS	vi
LIST OF TABLES	vii
LIST OF FIGURES	x
LIST OF SYMBOLS AND ABBREVIATIONS	xi
LIST OF APPENDIX	xiii
	xiv

CHAPTER 1: INTRODUCTION

1.1	Introduction of <i>Jatropha</i>	1
	1.1.1 Challenges for Commercial Scale <i>Jatropha</i> Plantation	2
	1.1.2 Future Perspective of <i>Jatropha curcas</i>	3
1.2	Significance of Study	3
1.3	Objectives	4
1.4	Research Approach	4

CHAPTER 2: LITERATURE REVIEW

2.1	<i>Jatropha curcas</i>	5
	2.1.1 Habitat and Ecology	6
	2.1.2 Botanical Characteristics	6
	2.1.3 Importance of <i>Jatropha curcas</i>	8
2.2	Callus	9
	2.2.1 Media and Culture Environment of Callus	10
	2.2.2 Advantages of Callus Culture	10
	2.2.3 Significance of Molecular Markers Associated With Light Stress	12
2.3	Chloroplast DNA	13
	2.3.1 <i>ycfs</i> (Hypothetical Chloroplast Open Reading Frames)	15
	2.3.2 Photosystem I and II	16
	2.3.3 Other Functional Genes in Chloroplast Genome	18
2.4	Expression of Chloroplast Genes	19
2.5	Reverse Transcription PCR and Real-time PCR	20

CHAPTER 3: MATERIALS AND METHODS

3.1	Plant Materials and Growth Conditions	22
3.2	DNA Extraction from Leaf Sample	23
	3.2.1 Sample Collection	23
	3.2.2 DNA Extraction	23
	3.2.3 Quantitation of DNA	24
3.3	Isolation of Total RNA from Callus	24
	3.3.1 Sample Preparation	24

3.3.2	Total RNA Isolation	25
3.3.3	Analysis of Quality, Stability and Quantification of Extracted Total RNA	25
3.4	Primer Design	27
3.5	Determination of Primer Specificity	27
3.6	First Strand Synthesis of cDNA from Callus RNA	38
3.7	Polymerase Chain Reaction Amplification	28
3.8	PCR Product Purification	30
3.9	Cloning of PCR Products	33
3.10	Transformation of TOP10 Competent Cells using Ligation Mix	37
3.11	Analysis of Recombinant Clones	35
3.12	Plasmid Miniprep	35
3.13	DNA Sequencing	36
3.14	Data Analysis	36
3.15	Real-time PCR	36

CHAPTER 4: RESULTS AND DISCUSSION

4.1	Reverse Transcription PCR Assay	
4.1.1	DNA Extraction and Amplification for Testing The Specificity of the Designed Primers	38
4.1.2	Callus Total RNA Extraction	41
4.1.3	Detection of DNA Contamination in RNA	43
4.1.4	cDNA Amplification using Gene Specific Primers	44
4.1.5	PCR Products Purification	50
4.1.6	Cloning of PCR Products	51
4.1.7	Analysis of Sequence Data	54
4.2	Real-time PCR Assay	
4.2.1	Confirmation of Primer Specificity	61
4.2.2	Detection of Chloroplast Genes under Photoperiod Using Real-time PCR	61

CHAPTER 5: CONCLUSION 72

REFERENCES 74

APPENDICES 81

LIST OF TABLES

		Page
Table 3.1	First component mixture for first strand cDNA Synthesis	28
Table 3.2	Second component mixture for first strand cDNA synthesis	28
Table 3.3	Set primers used for PCR amplification	29
Table 3.4	Component mixture for digestion	32
Table 3.5	Component mixture for blunting reaction	32
Table 3.6	PCR master mix for colony PCR	35
Table 4.1	Summary of reverse transcription PCR amplification of Chloroplast <i>ycf</i> gene against cDNA of callus and leaf of <i>J. curcas</i>	46
Table 4.2	Summary of reverse transcription PCR amplification of chloroplast <i>psbD</i> and reference genes against cDNA of callus and leaf of <i>J. curcas</i>	48
Table 4.3	Comparison of nucleotide amplified by leaf against specific primers using BLASTn	60
Table 4.4	Comparison of nucleotide amplified by callus against Using BLASTn	60
Table 4.5	BLASTn sequence alignment result for cDNA amplified by RBCL primers	61

LIST OF FIGURES

		Page
Figure 2.1	<i>Jatropha curcas</i> plant	5
Figure 2.2	Different plant parts of <i>Jatropha curcas</i>	7
Figure 2.3	Gene map of the <i>Jatropha curcas</i> chloroplast genome showing the large single copy (LSC), small single copy (SSC) and inverted repeat (IRa/IRb)	14
Figure 3.1	Callus of <i>Jatropha curcas</i> cultured in MS media (a) Green callus (b) White Callus	22
Figure 3.2	Leaf of <i>Jatropha curcas</i> used for DNA extraction	23
Figure 3.3	Total RNA extraction procedure	26
Figure 3.4	Amplification of <i>ycf2</i> gene targeted at different region In <i>Jatropha curcas</i> chloroplast genome	30
Figure 3.5	General procedure for gel extraction method	31
Figure 3.6	Schematic diagram for preparation of competent cells using CaCl_2 method	34
Figure 4.1	Electrophoresis of extracted DNA from leaf using CTAB method on a 1.5% TBE agarose gel	39
Figure 4.2	Electrophoresis of PCR products of DNA amplified on a 1.5% TBE agarose gel	39
Figure 4.3	Electrophoresis of total RNA on a 1.5% TBE agarose gel	42
Figure 4.4	Electrophoresis of PCR products of RNA amplified by RPOC1 primer on a 1.5% TBE agarose gel	43
Figure 4.5	Electrophoresis of PCR products of cDNA amplified by RPOC1 housekeeping primer on a 1.5% TBE agarose gel	44
Figure 4.6	Electrophoresis of PCR products of cDNA amplified by YCFD primer on a 1.5% TBE agarose gel	45
Figure 4.7	Electrophoresis of gel excised product on 1.5% TBE agarose gel	55
Figure 4.8	The transformed colonies formed on the LB-ampicillin plate after 16 hours of incubation at 37°C	57

Figure 4.9	Electrophoresis of colony PCR- screen for transformants transformed with ampicillin-resistant gene	58
Figure 4.10	Electrophoresis of recombinant clones purified using GeneJET™ Miniprep Kit on 1.5% TBE agarose gel	59
Figure 4.11	The DNA sequencing chromatogram of gel excised product	63
Figure 4.12	The short and poor trace of DNA sequencing chromatogram of cloning product	63
Figure 4.14	Real-time PCR amplification plot of cDNA from four treatments of <i>J. curcas</i> callus amplified by YCFD primer using SYBR green dye	68
Figure 4.15	Melting curve profile for real-time PCR amplification of cDNA from four treatments of <i>J. curcas</i> callus primers with YCFD	69
Figure 4.16	Electrophoresis of real-time PCR products with approximately 188 bp on 1.5% TBE agarose gel	70
Figure 4.17	Real-time PCR amplification plot of cDNA from four treatments of <i>J. curcas</i> callus amplified by housekeeping primer rpoC1, using SYBR green dye	71
Figure 4.18	Melting curve profile for real-time PCR amplification of cDNA from four treatments of <i>J. curcas</i> callus primers with rpoC1	72
Figure 4.19	Electrophoresis of real-time PCR products with on 1.5% TBE agarose gel	73

LIST OF SYMBOLS AND ABBREVIATIONS

%	percent
°C	degree of Celsius
µg/ml	microgram per milliliter
µl	microliter
BLASTn	basic local alignment search tool for nucleotide
bp	base pair
CaCl ₂	calcium chloride
cDNA	complementary DNA
dNTP	deoxynucleoside-5'-triphosphate
<i>E. coli</i>	<i>Escherichia coli</i>
g	gram
h	hour
HCN	hydrogen cyanide
kb	kilo base
kg	kilogram
LB	Luria Bertani
<i>matK</i>	maturase
M	molar
mg/ml	milligram per milliliter
mg	milligram
MgCl ₂	magnesium chloride
ml	milliliter
mm	millimeter
mM	millimolar
ORF	open reading frame
PCR	polymerase chain reaction
pmol/µl	picomole per microliter
RNA	ribonucleic acid
<i>rbcl</i>	rubisco
rcf	g-force
<i>rpoC</i>	RNA polymerase beta' chain
rpm	resolution per minute
s	seconds
SDS	sodium dodecyl sulphate
Ta	annealing temperature
TE	tris-ethylenediaminetetraacetic acid buffer
V	voltage
xg	times gravity
<i>ycf</i>	hypothetical chloroplast open reading frame

LIST OF APPENDIX

	Page
Appendix A: Composition of Murashige and Skoog (1962) Medium	86
Appendix B: Preparation of Solutions for CTAB DNA Extraction	87
Appendix C: Preparation of Solutions for Electrophoresis Purposes	88
Appendix D: Preparation of Media and Solutions	89
Appendix E: BLASTn Sequence Alignment result for cDNA amplified by rpoC1 primer	90
Appendix F: BLASTn Sequence Alignment result for cDNA amplified by YCFD primer	91
Appendix G: BLASTn Sequence Alignment result for cDNA amplified by KYCF2 primer	93
Appendix H: BLASTn Sequence Alignment result for cDNA amplified by PS2D2 primer	95
Appendix I: BLASTn Sequence Alignment result for cDNA amplified by RBCL primer	96
Appendix J: BLASTn Sequence Alignment result for cDNA amplified by MATK primer	97
Appendix K: BLASTn Sequence Alignment result for cDNA amplified by YCB primer	98
Appendix L: BLASTn Sequence Alignment result for cDNA amplified by YCF1 primer	100
Appendix M: BLASTn Sequence Alignment result for cDNA amplified by YCF2 primer	101

CHAPTER 1

INTRODUCTION

1.1 Introduction to *Jatropha*

Jatropha is a genus of approximately 175 succulent plants, shrubs and trees, from the family Euphorbiaceae. The *Jatropha* species includes *Jatropha cuneata*, *Jatropha curcas*, *Jatropha gossypifolia*, *Jatropha podagrica* and others. Recently, Goldman Sachs cited *J. curcas* as one of the best candidates for future biodiesel production (Barta, 2007). It is resistant to drought and pests, and produces seeds containing 27 to 40% oil. Unlike other biodiesel crops, *Jatropha* can be grown almost anywhere, including deserts, trash dumps, and rock piles. It does not need much water or fertilizer, and it is not edible.

The fact that *Jatropha* oil cannot be used for nutritional purposes without detoxification makes its use as energy or fuel source very attractive as biodiesel. In Madagascar, Cape Verde and Benin, *Jatropha* oil is used as a mineral diesel substitute during the Second World War (Akbar *et al.*, 2009).

In 2007, the Malaysian government cooperated with the Plantation Industries and Commodities Ministry and launched a *Jatropha* pilot project in Kota Marudu, Sabah. The ministry pledged to expand cultivation, especially in poorer areas, if the crops proved to be viable. Besides Malaysia, *Jatropha* projects have also been piloted in China, the Philippines, Rwanda, and the United State of America. In addition, the Indian Government implements the cultivation of *Jatropha* plants as the increased of *Jatropha* oil production delivers economic benefits to India at the macroeconomic or national levels as it reduces the nation's fuel import bill for diesel production.

1.1.1 Challenges for Commercial Scale *Jatropha* Plantation

The cultivation of *Jatropha* faces several challenges. *Jatropha* encounter various disease problems, such as the spider mite, white fly and the mealy bugs, especially on the surface of the leaves. It is reported that, in some areas of Zimbabwe, the golden flea beetle (*Podagrica* spp) can harm *Jatropha* and plays host to the 'frog eyes' fungus (*Cercospera* spp) (Openshaw, 2000).

Beside disease, toxicity of *Jatropha* is another challenge faced. The main toxic compounds are curcun, a protein that synthesis inhibitor, and phorbol ester (cancer promoting compound). Phorbol esters usually are ubiquitous while curcun stays mostly in seeds (Rakshit *et al.*, 2008). However, this is not a significant problem in traditional planting and uses. More concerns should be emphasized for large scale farming, for example the safety of workers, safety of grazing animals, possible adulteration of edible oil and other ecological concerns.

The key challenge for building *Jatropha* oil markets is coordinating feedstock supply and demand when neither exists for lack of the other. The smallholder farmers are unwilling to take risk of paying for, planting, and maintaining *Jatropha* trees. Nonetheless, the refiners are unwilling to make longer-term investments in refining capacity unless they have a secure source of adequate supply. Therefore, by developing institutional innovation, including contractual arrangements, fiscal and other incentives, blend ratio requirements, and other policy tools can overcome the obstacles in *Jatropha* biodiesel markets. Institutions will also be the key in determining the fairness of and allocation of risk in these markets (Weyerhaeuser *et al.*, 2007).

Uncertainty in how much *Jatropha* will cost to grow and the process of *Jatropha* into biodiesel seems to become another challenge for the exploitation of *Jatropha*. Some of this uncertainty is undoubtedly related to scale; once *Jatropha* growing begins on a commercial scale, costs are likely to fall at some level. However, increasing *Jatropha* acreage does not guarantee improvements in oil content and seed yields. If these remain low, unit production costs for *Jatropha* and

thus the subsidies required to make *Jatropha* biodiesel cost competitive will remain high (Weyerhaeuser *et al.*, 2007).

1.1.2 Future Perspective of *Jatropha curcas*

The development of *J. curcas* into a high-yielding and efficient new biofuel source is still at a relatively early stage. Therefore, in order to ensure the market of *J. curcas* moves forward, some tactical strategies need to be taken. For example, enhance priority in breeding either the traditional or molecular and distribution of uniform and superior planting materials. One of the key short to medium term research objectives is to search superior species aimed at higher overall yields of *Jatropha*. This will be achieved through systematic seed selection from different regions and supported by a scientific seed-breeding programme (Achten *et al.*, 2010). Other than that, more researches need to be carried out to address ecological concerns. Tree propagation techniques for specific climates and a wide variety of environmental conditions have to be refined (Vries, 2007).

1.2 Significance of Study

Now, not many studies have been done on the expression of genes encoded in the open reading frames of *J. curcas* chloroplast genome. In order to carry out preliminary study on the differential expression of chloroplast genes under condition of photoperiod and darkness, a total of seven ORFs representing the genes *ycf 1*, *ycf 2*, *psaB* (photosystem I), *psbD* (photosystem II), *rbcl*(RubisCO), *rpoC1* (RNA polymerase) and *matK* (Maturase K) were targeted by designing specific primers for their characterization. These primers were tested against transcriptome under different environment conditions. The molecular perspective of understanding the photosynthesis of *J. curcas*, is the key to productivity, hence, the regulatory mechanism of the gene responsible for photosynthesis is vital for future research work involving *Jatropha*. In addition, by studying the gene expression of chloroplast DNA of *J. curcas*, potential genes response to light stress can be potential candidate for callus transformation and become molecular marker for callus regeneration.

1.3 Objectives

The objectives of this study are:

- a. To identify the chloroplast genes which are induced under conditions of light and complete darkness in *Jatropha curcas*.
- b. To validate expression of genes using real-time PCR.

1.4 Research Approach

The research approach involves the callus culture conditions, RNA extraction, amplification and gel extraction or cloning. One of the approaches, which is through the design of conserved or gene specific primers which will target the transcripts to be induced by light and dark condition. Validation of gene expression was carried out using quantitative real-time PCR.



UMS
UNIVERSITI MALAYSIA SABAH

CHAPTER 2

LITERATURE REVIEW

2.1 *Jatropha curcas*

J. curcas (Figure 2.1), which belongs to the Family Euphorbiaceae have different local names, for example Barbados Nut, Jarak pagar (Indonesia), Physic Nut and Puring Nut. The genus name of *Jatropha* derives from the Greek *jatrós* (doctor), *trophé* (food), which implies medical uses. *Curcas* is the common name for physic nut in Malabar, India (Hanna-Jones and Csurches, 2008).



Figure 2.1: *Jatropha curcas* plant
(source:<http://www.lijunoilacnepimple.com/lijun01/jatrophaE.htm>)

2.1.1 Habitat and Ecology

The plant is native to the American tropics, mostly in Mexico and Central America. Currently it is cultivated in almost all tropical and subtropical countries as protection hedges gardens and fields (Verm and Gaur, 2009).

Jatropha is a poisonous bush or small tree, which can reach the height of five meters and grows well with more than 600 mm of rainfall per year and withstands long drought periods (Verm and Gaur, 2009).

2.1.2 Botanical Characteristics

J. curcas's branches exude whitish colored watery latex, upon cut. It has yellow-green flowers and large heart-shaped (pale) green leaves, arranged alternately. The inflorescence is formed in the leaf axel while the flowers are formed terminally, individually, with female flowers usually slightly larger (Verma and Gaur, 2009).

After pollination, a trilocular ellipsoidal fruit is formed. The exocarp remains fleshy until the seeds are mature. The seeds are black and in the average 18 mm long and 10 mm wide. The seed weight (per 1000) is about 727 g, this are 1375 seeds per kg in the average (Verma and Gaur, 2009). Different plant parts of *J. curcas* are shown in Figure 2.2.




Figure 2.2:

Different plant parts of *Jatropha curcas*, (a) whole plant, (b) bark, (c) leaf, (d) flower, (e) flower bud, (f) internal of fruit (g) fruits

(Source: Verma and Gaur, 2009)

2.1.3 Importance of *J. curcas*

J. curcas is considered as an ideal plant for wetland and its non-edible oil is a prime choice for biodiesel production. Different parts of *J. curcas* have traditionally been used for various purposes and have considerable potential. The importance of *Jatropha* can be categorized into two major parts which are fuel use and non-fuel use.

The oil from *Jatropha* is regarded as a potential fuel substitute, with a hydrocarbon of 16 to 18 carbon atoms per molecule. Compared with diesel oil, *Jatropha* biodiesel molecules are simple hydrocarbon chains, containing no sulphur, or aromatic substances associated with fossil fuels. They contain higher amount of oxygen (up to 10%) that ensures more complete combustion of hydrocarbons. The first successful trial run of a passenger train was conducted on December 31, 2002 when the Delhi-Amritsar Shatabdi Express used 5% biodiesel as fuel (Pahl, 2008).

The fuels can be obtained directly from different parts of *Jatropha* plant, such as wood, the whole fruit and parts of the fruit burnt separately or in combination, namely the exocarp (coat), the nut shell and the kernel (Openshaw, 2000).

Also, *Jatropha* could be used as a substitute for wire fencing for posts round fields, along roadsides and railway tracks. In countries like Zimbabwe, Mali, *Jatropha* is mostly used by farmers as fences around their homesteads and gardens, sometimes also around their fields, to protect crops against roaming animals. In addition, by planting *J. curcas*, it functions to reclaim eroded land and *Jatropha* hedges and shelterbelts can assist other crops on all land types by keeping out animals, improving the microclimate and providing humus to the soil (Openshaw, 2000).

Jatropha seeds contain fatty acid or viscous oil and one of its uses is as a raw material for soap making in cosmetic industries. In India, it is used by a large industry (Hindustan Lever). In Zimbabwe, there is a high demand of *Jatropha* oil

although soap is produced by small informal industries in rural areas (Openshaw, 2000).

According to Osoniyi and Onajobi (2003), *J. curcas* is a medicinal plant, traditionally used as a haemostatic. Some of the ethnomedical uses of the extracts of the leaves and roots are remedy for cancer, as an abortifacient, antiseptic, diuretic, and purgative. The nut of plants has also been used traditionally for the treatment of many ailments including burns, convulsions, fever and inflammation. Esimone *et al.* (2009) demonstrated a significant wound-healing activity in *J. curcas* extracts by applying the latex directly, which is an alkaloid known as jatrophine, or the crushed leaf of the plant to cuts and bleeding wounds.

The ashes of *J. curcas* can also be used as salt substitute, whereas the bark which contains HCN, can be used as fish poison. Besides that, the latex can strongly inhibit the watermelon mosaic virus, which is a kind of virus that can infect all commercial cucurbit crops. In Mexico, the people grow the shrub as a host for the lac insect, which is used in medicine as hepatoprotective and antiobesity drug and also for erosion control (Tee, 2009).

2.2 Callus

Evans *et al.* (2003) defined callus as an amorphous mass of unorganized thin-walled parenchyma cells. Callus is formed at the cut surfaces when the plant is wounded. It is thought to be a protective response by the plant to seal off damaged tissue. According to Purohit (2003), callus is the results when a cell from any part of the plant, such as shoot apex, bud, leaf, mesophyll cells, epidermis, cambium, anthers, pollen or fruit is inoculated in a suitable medium under aseptic laboratory conditions. They are able to differentiate and survive.

Callus varies widely in its general appearance and in other physical features. The variation usually depends on the parent tissue, the age of the callus and the growth conditions. Callus can be in white, green or highly coloured due to the presence of anthocyanin pigments (Evans *et al.*, 2003).

2.2.1 Media and Culture Environment of Callus

Culture media used for the *in vitro* cultivation of plant tissues are composed of three basic components: (i) essential elements, or mineral ions, supplied as a complex mixture of salts; (ii) an organic supplement supplying vitamins and/or amino acids; and (iii) a source of fixed carbon which is usually supplied as the sugar sucrose. The composition of the nutrient solution plays a key role in plant nutrition. A defined nutrient medium consists of inorganic salts, a carbon source, vitamins and growth regulators. Generally, the media can be divided into micronutrients and macronutrients which are necessary for plant tissue or cell culture.

The MS medium was published in 1962 (Murashige and Skoog, 1962) and was used to support rapid growth of tobacco tissue culture and it is the most widely used salt composition, especially in procedures where plant regeneration is the objective. There are over 20 variants based on MS basic medium embracing a large number of specialty cultures (Evans *et al.* 2003). The B5, N6, Nitsch and Nitsch, and derivatives of these media have been applied widely for many plant species and for different objectives.

When cultured *in vitro*, all the needs, both chemical and physical, of the plants cells have to be met by the culture vessel, the growth medium has to supply all the essential mineral ions required for growth and development. In many cases, it must also supply additional organic supplement such as amino acid and vitamins. Many plant cell cultures, as they are not photosynthetic, also require the addition of a fixed carbon source in the form of sugar, for example sucrose. Physical factors, such as temperature, pH, the gaseous environment, light in the term of quality and duration, and osmotic pressure, also have to be maintained within acceptable limits (Vasil, 1994).

2.2.2 Advantages of Callus Culture

The callus culture is a technique of tissue culture, which is usually carried out on solidified gel medium in the presence of growth regulators and initiated by inoculation of small explants. Callus cultures are being widely used for study of expression of race-specific and non-host resistance and offer several advantages